

Zhen Fan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5977640/publications.pdf>

Version: 2024-02-01

43
papers

3,510
citations

172457

29
h-index

254184

43
g-index

43
all docs

43
docs citations

43
times ranked

5425
citing authors

#	ARTICLE	IF	CITATIONS
1	Redirecting host preexisting influenza A virus immunity for cancer immunotherapy. <i>Cancer Immunology, Immunotherapy</i> , 2022, 71, 1611-1623.	4.2	2
2	ASCT2 overexpression is associated with poor survival of OSCC patients and ASCT2 knockdown inhibited growth of glutamine-addicted OSCC cells. <i>Cancer Medicine</i> , 2020, 9, 3489-3499.	2.8	20
3	Rational combination with PDK1 inhibition overcomes cetuximab resistance in head and neck squamous cell carcinoma. <i>JCI Insight</i> , 2019, 4, .	5.0	25
4	Functional cooperation between HIF-1 α and c-Jun in mediating primary and acquired resistance to gefitinib in NSCLC cells with activating mutation of EGFR. <i>Lung Cancer</i> , 2018, 121, 82-90.	2.0	21
5	Trastuzumab upregulates PD-L1 as a potential mechanism of trastuzumab resistance through engagement of immune effector cells and stimulation of IFN γ secretion. <i>Cancer Letters</i> , 2018, 430, 47-56.	7.2	117
6	AP1G1 is involved in cetuximab-mediated downregulation of ASCT2-EGFR complex and sensitization of human head and neck squamous cell carcinoma cells to ROS-induced apoptosis. <i>Cancer Letters</i> , 2017, 408, 33-42.	7.2	31
7	Acetyl-CoA carboxylase rewires cancer metabolism to allow cancer cells to survive inhibition of the Warburg effect by cetuximab. <i>Cancer Letters</i> , 2017, 384, 39-49.	7.2	63
8	ASCT2 (SLC1A5) is an EGFR-associated protein that can be co-targeted by cetuximab to sensitize cancer cells to ROS-induced apoptosis. <i>Cancer Letters</i> , 2016, 381, 23-30.	7.2	51
9	Overcoming cisplatin resistance of ovarian cancer cells by targeting HIF-1-regulated cancer metabolism. <i>Cancer Letters</i> , 2016, 373, 36-44.	7.2	135
10	Trastuzumab upregulates expression of HLA-ABC and T cell costimulatory molecules through engagement of natural killer cells and stimulation of IFN γ secretion. <i>Oncolmmunology</i> , 2016, 5, e1100790.	4.6	46
11	Identification and validation of COX-2 as a co-target for overcoming cetuximab resistance in colorectal cancer cells. <i>Oncotarget</i> , 2016, 7, 64766-64777.	1.8	22
12	AMPK-mediated energy homeostasis and associated metabolic effects on cancer cell response and resistance to cetuximab. <i>Oncotarget</i> , 2015, 6, 11507-11518.	1.8	29
13	Autocrine/paracrine erythropoietin regulates migration and invasion potential and the stemness of human breast cancer cells. <i>Cancer Biology and Therapy</i> , 2014, 15, 89-98.	3.4	12
14	A novel role of EMMPRIN/CD147 in transformation of quiescent fibroblasts to cancer-associated fibroblasts by breast cancer cells. <i>Cancer Letters</i> , 2013, 335, 380-386.	7.2	33
15	HER2 regulates Brk/PTK6 stability via upregulating calpastatin, an inhibitor of calpain. <i>Cellular Signalling</i> , 2013, 25, 1754-1761.	3.6	16
16	Brk/PTK6 cooperates with HER2 and Src in regulating breast cancer cell survival and epithelial-to-mesenchymal transition. <i>Cancer Biology and Therapy</i> , 2013, 14, 237-245.	3.4	32
17	Cetuximab Reverses the Warburg Effect by Inhibiting HIF-1 α -Regulated LDH-A. <i>Molecular Cancer Therapeutics</i> , 2013, 12, 2187-2199.	4.1	67
18	The anti-EGFR antibody cetuximab sensitizes human head and neck squamous cell carcinoma cells to radiation in part through inhibiting radiation-induced upregulation of HIF-1 α . <i>Cancer Letters</i> , 2012, 322, 78-85.	7.2	47

#	ARTICLE	IF	CITATIONS
19	Resveratrol-Activated AMPK/SIRT1/Autophagy in Cellular Models of Parkinson's Disease. <i>NeuroSignals</i> , 2011, 19, 163-174.	0.9	405
20	Constitutively active Harvey Ras confers resistance to epidermal growth factor receptor-targeted therapy with cetuximab and gefitinib. <i>Cancer Letters</i> , 2011, 306, 85-91.	7.2	10
21	Differential Turnover of Myosin Chaperone UNC-45A Isoforms Increases in Metastatic Human Breast Cancer. <i>Journal of Molecular Biology</i> , 2011, 412, 365-378.	4.2	27
22	Recombinant Human Erythropoietin Antagonizes Trastuzumab Treatment of Breast Cancer Cells via Jak2-Mediated Src Activation and PTEN Inactivation. <i>Cancer Cell</i> , 2010, 18, 423-435.	16.8	129
23	The Epidermal Growth Factor Receptor Antibody Cetuximab Induces Autophagy in Cancer Cells by Downregulating HIF-1 α and Bcl-2 and Activating the Beclin 1/hVps34 Complex. <i>Cancer Research</i> , 2010, 70, 5942-5952.	0.9	172
24	Roles of autophagy in cetuximab-mediated cancer therapy against EGFR. <i>Autophagy</i> , 2010, 6, 1066-1077.	9.1	87
25	1, 9-Pyrazoloanthrones Downregulate HIF-1 α and Sensitize Cancer Cells to Cetuximab-Mediated Anti-EGFR Therapy. <i>PLoS ONE</i> , 2010, 5, e15823.	2.5	16
26	Requirement of hypoxia-inducible factor-1 α down-regulation in mediating the antitumor activity of the anti-epidermal growth factor receptor monoclonal antibody cetuximab. <i>Molecular Cancer Therapeutics</i> , 2008, 7, 1207-1217.	4.1	59
27	Epidermal Growth Factor Receptor (EGFR) Ubiquitination as a Mechanism of Acquired Resistance Escaping Treatment by the Anti-EGFR Monoclonal Antibody Cetuximab. <i>Cancer Research</i> , 2007, 67, 8240-8247.	0.9	149
28	Responses of cancer cells with wild-type or tyrosine kinase domain-mutated epidermal growth factor receptor (EGFR) to EGFR-targeted therapy are linked to downregulation of hypoxia-inducible factor-1 α . <i>Molecular Cancer</i> , 2007, 6, 63.	19.2	55
29	Differential Roles of Phosphoinositide-Dependent Protein Kinase-1 and Akt1 Expression and Phosphorylation in Breast Cancer Cell Resistance to Paclitaxel, Doxorubicin, and Gemcitabine. <i>Molecular Pharmacology</i> , 2006, 70, 1045-1052.	2.3	48
30	Autophosphorylation of Akt at Threonine 72 and Serine 246. <i>Journal of Biological Chemistry</i> , 2006, 281, 13837-13843.	3.4	25
31	The anti-epidermal growth factor receptor monoclonal antibody cetuximab/C225 reduces hypoxia-inducible factor-1 alpha, leading to transcriptional inhibition of vascular endothelial growth factor expression. <i>Oncogene</i> , 2005, 24, 4433-4441.	5.9	120
32	Differential responses to doxorubicin-induced phosphorylation and activation of Akt in human breast cancer cells. <i>Breast Cancer Research</i> , 2005, 7, R589-97.	5.0	75
33	C225 anti-epidermal growth factor receptor antibody enhances the efficacy of docetaxel chemoradiotherapy. <i>International Journal of Radiation Oncology Biology Physics</i> , 2004, 59, 1163-1173.	0.8	49
34	The epidermal growth factor receptor mediates radioresistance. <i>International Journal of Radiation Oncology Biology Physics</i> , 2003, 57, 246-254.	0.8	272
35	HER2/PI-3K/Akt activation leads to a multidrug resistance in human breast adenocarcinoma cells. <i>Oncogene</i> , 2003, 22, 3205-3212.	5.9	406
36	Sensitization of breast cancer cells to radiation by trastuzumab. <i>Molecular Cancer Therapeutics</i> , 2003, 2, 1113-20.	4.1	189

#	ARTICLE	IF	CITATIONS
37	Inhibition of angiogenesis by the antiepidermal growth factor receptor antibody ImClone C225 in androgen-independent prostate cancer growing orthotopically in nude mice. <i>Clinical Cancer Research</i> , 2002, 8, 1253-64.	7.0	70
38	C225 antiepidermal growth factor receptor antibody enhances tumor radiocurability. <i>International Journal of Radiation Oncology Biology Physics</i> , 2001, 51, 474-477.	0.8	136
39	Antitumor effect of an HER2-specific antibody-toxin fusion protein on human prostate cancer cells. <i>Prostate</i> , 2001, 47, 21-28.	2.3	32
40	Fibroblast growth factor and insulin-like growth factor differentially modulate the apoptosis and G1 arrest induced by anti-epidermal growth factor receptor monoclonal antibody. <i>Oncogene</i> , 2001, 20, 1913-1922.	5.9	107
41	The monoclonal antibody 225 activates caspase-8 and induces apoptosis through a tumor necrosis factor receptor family-independent pathway. <i>Oncogene</i> , 2001, 20, 3726-3734.	5.9	40
42	Antitumor effect of an HER2-specific antibody-toxin fusion protein on human prostate cancer cells. <i>Prostate</i> , 2001, 47, 21-28.	2.3	2
43	Differential modulation of paclitaxel-mediated apoptosis by p21Waf1 and p27Kip1. <i>Oncogene</i> , 2000, 19, 2423-2429.	5.9	61