

# G Fonseca-Camarillo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5979272/publications.pdf>

Version: 2024-02-01

58  
papers

840  
citations

516710

16  
h-index

501196

28  
g-index

62  
all docs

62  
docs citations

62  
times ranked

1545  
citing authors

#	ARTICLE	IF	CITATIONS
1	Expression of TOB/BTG family members in patients with inflammatory bowel disease. <i>Scandinavian Journal of Immunology</i> , 2021, 93, e13004.	2.7	11
2	Increased expression of extracellular matrix metalloproteinase inducer (EMMPRIN) and MMP10, MMP23 in inflammatory bowel disease: Cross-sectional study. <i>Scandinavian Journal of Immunology</i> , 2021, 93, e12962.	2.7	12
3	Intestinal production of secreted protein acidic and rich in cysteine (SPARC) in patients with ulcerative colitis. <i>Immunobiology</i> , 2021, 226, 152095.	1.9	6
4	Gene expression profiling of inflammatory cytokines in esophageal biopsies of different phenotypes of gastroesophageal reflux disease: a cross-sectional study. <i>BMC Gastroenterology</i> , 2021, 21, 201.	2.0	14
5	AKAP12/Gravin is over-expressed in patients with ulcerative colitis. <i>Immunologic Research</i> , 2021, 69, 429-435.	2.9	3
6	Differential Cytokine Expression in the Duodenum and Rectum of Children with Non-Immunoglobulin E-Mediated Cow's Milk Protein Allergy. <i>Digestive Diseases and Sciences</i> , 2021, 66, 3769-3775.	2.3	6
7	Synthesis of Interleukin-10 in Patients with Ulcerative Colitis and <i>Helicobacter pylori</i> Infection. <i>Gastroenterology Research and Practice</i> , 2020, 2020, 1-7.	1.5	4
8	TRPV Subfamily (TRPV2, TRPV3, TRPV4, TRPV5, and TRPV6) Gene and Protein Expression in Patients with Ulcerative Colitis. <i>Journal of Immunology Research</i> , 2020, 2020, 1-11.	2.2	14
9	Gene Expression Profiling of Mediators Associated with the Inflammatory Pathways in the Intestinal Tissue from Patients with Ulcerative Colitis. <i>Mediators of Inflammation</i> , 2020, 2020, 1-11.	3.0	23
10	Canonical and non-canonical Wnt signaling are simultaneously activated by Wnts in colon cancer cells. <i>Cellular Signalling</i> , 2020, 72, 109636.	3.6	59
11	Differential Expression of IL-36 Family Members and IL-38 by Immune and Nonimmune Cells in Patients with Active Inflammatory Bowel Disease. <i>BioMed Research International</i> , 2018, 2018, 1-12.	1.9	47
12	The Transient Receptor Potential Vanilloid 1 Is Associated with Active Inflammation in Ulcerative Colitis. <i>Mediators of Inflammation</i> , 2018, 2018, 1-7.	3.0	13
13	Caspase recruitment domain (CARD) family (CARD9, CARD10, CARD11, CARD14 and CARD15) are increased during active inflammation in patients with inflammatory bowel disease. <i>Journal of Inflammation</i> , 2018, 15, 13.	3.4	19
14	T $\beta$ 2RIII is induced by TCR signaling and downregulated in FoxP3+ regulatory T cells. <i>Biochemical and Biophysical Research Communications</i> , 2017, 494, 82-87.	2.1	9
15	Expression of Genes Associated with Inflammation in Biopsies of Esophageal Mucosa of different Phenotypes of Gastroesophageal Reflux Disease. <i>Gastroenterology</i> , 2017, 152, S237.	1.3	0
16	Role of IL-38 and its Antagonist in Patients with Inflammatory Bowel Disease. <i>Gastroenterology</i> , 2017, 152, S762.	1.3	1
17	P164 FOXO4 gene expression in colonic mucosa is a potential marker associated with histological remission in patients with ulcerative colitis. <i>Journal of Crohn's and Colitis</i> , 2017, 11, S160-S161.	1.3	0
18	Sa1848 IL34 and IL36 Family Expressing Cytotoxic T cells and Plasmacytoid Dendritic Cells are Increased in Patients With Active Inflammatory Bowel Disease. <i>Gastroenterology</i> , 2016, 150, S379-S380.	1.3	0

#	ARTICLE	IF	CITATIONS
19	Mo1930 Transcriptome Analysis of Immune Innate Response Genes in the Colonic Mucosa from Patients With Ulcerative Colitis. <i>Gastroenterology</i> , 2016, 150, S819.	1.3	0
20	Interleukin 27 is up-regulated in patients with active inflammatory bowel disease. <i>Immunologic Research</i> , 2016, 64, 901-907.	2.9	23
21	Tu1981 The Oxido-reductases Enzymes (TDO2 and SOD2) in Colonic Mucosa are Markers Associated with Histological Activity and Clinical Course in Ulcerative Colitis. <i>Gastroenterology</i> , 2016, 150, S996.	1.3	0
22	Genetic Markers Associated with Clinical Outcomes in Patients with Inflammatory Bowel Disease. <i>Inflammatory Bowel Diseases</i> , 2015, 21, 2683-2695.	1.9	14
23	Immunoregulatory Pathways Involved in Inflammatory Bowel Disease. <i>Inflammatory Bowel Diseases</i> , 2015, 21, 2188-2193.	1.9	83
24	Differential Expression of MUC12, MUC16, and MUC20 in Patients with Active and Remission Ulcerative Colitis. <i>Mediators of Inflammation</i> , 2015, 2015, 1-8.	3.0	29
25	Interleukin 35 (IL-35) and IL-37: Intestinal and peripheral expression by T and B regulatory cells in patients with Inflammatory Bowel Disease. <i>Cytokine</i> , 2015, 75, 389-402.	3.2	66
26	Sa1243 Defensins Alpha 5 and 6 Are Up-Regulated in the Colonic Mucosa From Patients With Ulcerative Colitis and Are Associated With Clinical Outcomes. <i>Gastroenterology</i> , 2015, 148, S-267-S-268.	1.3	0
27	Su1257 The Gene Expression of SPARC in the Colonic Mucosa Is Associated With Histological Activity in Patients With Ulcerative Colitis. <i>Gastroenterology</i> , 2015, 148, S-453.	1.3	1
28	Mo1711 Interleukin 35 and 37 Intestinal Expression and Peripheral Synthesis by Subsets of Regulatory Cells in Patients With Inflammatory Bowel Disease. <i>Gastroenterology</i> , 2015, 148, S-692.	1.3	0
29	Mo1712 Role of Interleukin 27 (IL-27) in the Colonic Mucosa of Patients With Inflammatory Bowel Disease. <i>Gastroenterology</i> , 2015, 148, S-692.	1.3	1
30	Peroxisome Proliferator-Activated Receptors Family Is Involved in the Response to Treatment and Mild Clinical Course in Patients with Ulcerative Colitis. <i>Disease Markers</i> , 2014, 2014, 1-7.	1.3	22
31	Expression of interleukin (IL)-19 and IL-24 in inflammatory bowel disease patients: a cross-sectional study. <i>Clinical and Experimental Immunology</i> , 2014, 177, 64-75.	2.6	58
32	P173 Peroxisome proliferator-activated receptors family is involved in the response to treatment and clinical course in patients with ulcerative colitis. <i>Journal of Crohn's and Colitis</i> , 2014, 8, S134.	1.3	0
33	Tu1714 Interleukins 19 and 24 Are Highly Expressed in the Intestine and Peripheral Blood Cells From Crohn's Disease Patients. <i>Gastroenterology</i> , 2014, 146, S-824.	1.3	0
34	P004 Transcriptome of unfolded protein response (UPR) genes in the colonic mucosa of patients with ulcerative colitis. <i>Journal of Crohn's and Colitis</i> , 2014, 8, S66.	1.3	0
35	Su1364 Peroxisome Proliferator-Activated Receptors Alpha and Gamma Are Associated With Favorable Response to 5 Aminosallylates and Mild Clinical Course of Ulcerative Colitis. <i>Gastroenterology</i> , 2014, 146, S-447.	1.3	0
36	P185 Mucin 16 (MUC16) and mucin 20 (MUC20) over-expression in colonic mucosa is associated with histological remission in patients with ulcerative colitis. <i>Journal of Crohn's and Colitis</i> , 2014, 8, S139.	1.3	1

#	ARTICLE	IF	CITATIONS
37	P005 Transcriptome of intestinal epithelial barrier genes in the colonic mucosa from patients with ulcerative colitis. <i>Journal of Crohn's and Colitis</i> , 2014, 8, S66-S67.	1.3	0
38	High Gene Expression of CXCL8 Is Associated with the Presence of Extraintestinal Manifestations and Long-term Disease in Patients with Ulcerative Colitis. <i>Inflammatory Bowel Diseases</i> , 2013, 19, E22-E23.	1.9	6
39	IL-10 <sup>+</sup> and IL-20 <sup>+</sup> Expressing Epithelial and Inflammatory Cells are Increased in Patients with Ulcerative Colitis. <i>Journal of Clinical Immunology</i> , 2013, 33, 640-648.	3.8	58
40	Mo1330 Transcriptome of Unfolded Protein Response (UPR) Genes in the Colonic Mucosa From Patients With Ulcerative Colitis. <i>Gastroenterology</i> , 2013, 144, S-638-S-639.	1.3	0
41	Tu1129 Expression of Centaurin Beta 1 (CENTB1) Is up-Regulated in Patients With Ulcerative Colitis. <i>Gastroenterology</i> , 2013, 144, S-770.	1.3	0
42	Gene and protein expression of centaurin beta 1 (CENTB1) are up-regulated in patients with ulcerative colitis. <i>Journal of Crohn's and Colitis</i> , 2013, 7, e238-e239.	1.3	3
43	Indoleamine 2,3-Dioxygenase: Expressing Cells in Inflammatory Bowel Disease—A Cross-Sectional Study. <i>Clinical and Developmental Immunology</i> , 2013, 2013, 1-14.	3.3	17
44	Interleukins Involved in Inflammatory Bowel Disease as New Therapeutic Targets. <i>Current Immunology Reviews</i> , 2013, 9, 86-92.	1.2	4
45	Sa1287 Up Regulation of Interleukin 24 in Colonic Tissue and Peripheral Blood Mononuclear Cells is Associated With the Presence of Extraintestinal Manifestations in Patients With Ulcerative Colitis. <i>Gastroenterology</i> , 2012, 142, S-263-S-264.	1.3	0
46	Tu1893 Increased Expression of IL-20 and Its Receptors Alpha and Beta in Patients With Ulcerative Colitis. <i>Gastroenterology</i> , 2012, 142, S-871.	1.3	0
47	Differential expression of occludin in patients with ulcerative colitis and healthy controls. <i>Inflammatory Bowel Diseases</i> , 2012, 18, E1999.	1.9	30
48	Gene expression of solute carrier family 9 (Sodium/Hydrogen Exchanger) 3, (SLC9A3) is downregulated in patients with ulcerative colitis. <i>Inflammatory Bowel Diseases</i> , 2012, 18, 1197-1198.	1.9	9
49	Reduced expression of mucin 9 (MUC9) in patients with ulcerative colitis. <i>Inflammatory Bowel Diseases</i> , 2012, 18, E601.	1.9	9
50	Differential Expression of Disc Large Homologue 5 (DLG5) and Carnitine Organic Cation Transporter (OCTN) Genes in Patients With Ulcerative Colitis. <i>Gastroenterology</i> , 2011, 140, S-422.	1.3	0
51	Transcript levels of Toll-Like receptors 5, 8 and 9 correlate with inflammatory activity in Ulcerative Colitis. <i>BMC Gastroenterology</i> , 2011, 11, 138.	2.0	58
52	Increased expression of discs large homolog 5 gene (DLG5) in ulcerative colitis patients compared to healthy individuals. <i>Inflammatory Bowel Diseases</i> , 2011, 17, 1639.	1.9	2
53	Gene expression of carnitine organic cation transporters 1 and 2 (OCTN) is downregulated in patients with ulcerative colitis. <i>Inflammatory Bowel Diseases</i> , 2011, 17, 2205-2206.	1.9	11
54	Interleukin 17 gene and protein expression are increased in patients with ulcerative colitis. <i>Inflammatory Bowel Diseases</i> , 2011, 17, E135-E136.	1.9	26

#	ARTICLE	IF	CITATIONS
55	High Gene Expression of MDR1 (ABCB1) is Associated with Medical Treatment Response and Long-Term Remission in Patients with Ulcerative Colitis. <i>Inflammatory Bowel Diseases</i> , 2010, 16, 541-542.	1.9	11
56	Interleukin 21 Expression is Increased in Rectal Biopsies from Patients with Ulcerative Colitis. <i>Inflammatory Bowel Diseases</i> , 2010, 16, 1090.	1.9	16
57	TLR9 mRNA expression is upregulated in patients with active ulcerative colitis. <i>Inflammatory Bowel Diseases</i> , 2010, 16, 1267-1268.	1.9	16
58	Colonic epithelial upregulation of interleukin 22 (IL-22) in patients with ulcerative colitis. <i>Inflammatory Bowel Diseases</i> , 2010, 16, 1823.	1.9	25