

Ian R Kelsall

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5985152/publications.pdf>

Version: 2024-02-01

16
papers

1,158
citations

687363

13
h-index

940533

16
g-index

18
all docs

18
docs citations

18
times ranked

2099
citing authors

#	ARTICLE	IF	CITATIONS
1	HOIL-1 ubiquitin ligase activity targets unbranched glucosaccharides and is required to prevent polyglucosan accumulation. <i>EMBO Journal</i> , 2022, 41, e109700.	7.8	51
2	HOIL-1, an atypical E3 ligase that controls MyD88 signalling by forming ester bonds between ubiquitin and components of the Myddosome. <i>Advances in Biological Regulation</i> , 2020, 75, 100666.	2.3	14
3	Coupled monoubiquitylation of the co-E3 ligase DCNL1 by Ariadne-RBR E3 ubiquitin ligases promotes cullin-RING ligase complex remodeling. <i>Journal of Biological Chemistry</i> , 2019, 294, 2651-5314.	3.4	13
4	The E3 ligase HOIL-1 catalyses ester bond formation between ubiquitin and components of the Myddosome in mammalian cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 13293-13298.	7.1	102
5	IFI16 and cGAS cooperate in the activation of STING during DNA sensing in human keratinocytes. <i>Nature Communications</i> , 2017, 8, 14392.	12.8	251
6	Roles of the TRAF6 and Pellino E3 ligases in MyD88 and RANKL signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E3481-E3489.	7.1	88
7	Blocking an N-terminal acetylation-dependent protein interaction inhibits an E3 ligase. <i>Nature Chemical Biology</i> , 2017, 13, 850-857.	8.0	80
8	Two Distinct Types of E3 Ligases Work in Unison to Regulate Substrate Ubiquitylation. <i>Cell</i> , 2016, 166, 1198-1214.e24.	28.9	172
9	Lys63/Met1-hybrid ubiquitin chains are commonly formed during the activation of innate immune signalling. <i>Biochemical and Biophysical Research Communications</i> , 2016, 474, 452-461.	2.1	77
10	TRIAD1 and HHARI bind to and are activated by distinct neddylated Cullin-RING ligase complexes. <i>EMBO Journal</i> , 2013, 32, 2848-2860.	7.8	84
11	The Fanconi Anaemia Components UBE2T and FANCM Are Functionally Linked to Nucleotide Excision Repair. <i>PLoS ONE</i> , 2012, 7, e36970.	2.5	38
12	R3F, a novel membrane-associated glycogen targeting subunit of protein phosphatase 1 regulates glycogen synthase in astrocytoma cells in response to glucose and extracellular signals. <i>Journal of Neurochemistry</i> , 2011, 118, 596-610.	3.9	17
13	Ppm1E is an in cellulo AMP-activated protein kinase phosphatase. <i>Cellular Signalling</i> , 2011, 23, 114-124.	3.6	98
14	Disruption of the allosteric phosphorylase a regulation of the hepatic glycogen-targeted protein phosphatase 1 improves glucose tolerance in vivo. <i>Cellular Signalling</i> , 2009, 21, 1123-1134.	3.6	34
15	Disruption of the striated muscle glycogen-targeting subunit of protein phosphatase 1: influence of the genetic background. <i>Journal of Molecular Endocrinology</i> , 2008, 40, 47-59.	2.5	7
16	The hepatic PP1 glycogen-targeting subunit interaction with phosphorylase <i>α</i> can be blocked by C-terminal tyrosine deletion or an indole drug. <i>FEBS Letters</i> , 2007, 581, 4749-4753.	2.8	26