Gustavo Dominguez-Bernal

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6077378/publications.pdf

Version: 2024-02-01

44 papers 4,889

18 h-index

430754

243529 44 g-index

44 all docs

44 docs citations

44 times ranked 4109 citing authors

#	Article	IF	CITATIONS
1	Transcriptomic Profile of Canine DH82 Macrophages Infected by Leishmania infantum Promastigotes with Different Virulence Behavior. International Journal of Molecular Sciences, 2022, 23, 1466.	1.8	4
2	Properties of virulence emergence of <i>Leishmania infantum </i> isolates from <i>Phlebotomus perniciosus </i> collected during the human leishmaniosis outbreak in Madrid, Spain. Hepatic histopathology and immunological parameters as virulence markers in the mouse model. Transboundary and Emerging Diseases, 2021, 68, 704-714.	1.3	9
3	Raccoons (Procyon lotor) in the Madrid region of Spain are carriers of antimicrobialâ€resistant Escherichia coli and enteropathogenic E.Âcoli. Zoonoses and Public Health, 2021, 68, 69-78.	0.9	4
4	Epitope Selection for Fighting Visceral Leishmaniosis: Not All Peptides Function the Same Way. Vaccines, 2020, 8, 352.	2.1	2
5	A further investigation of the leishmaniosis outbreak in Madrid (Spain): low-infectivity phenotype of the Leishmania infantum BOS1FL1 isolate to establish infection in canine cells. Veterinary Immunology and Immunopathology, 2020, 230, 110148.	0.5	4
6	Strength and medium-term impact of HisAK70 immunization in dogs: Vaccine safety and biomarkers of effectiveness for ex vivo Leishmania infantum infection. Comparative Immunology, Microbiology and Infectious Diseases, 2019, 65, 137-143.	0.7	7
7	Labelâ€free bioanalysis of <i>Leishmania infantum</i> using refractive index tomography with partially coherent illumination. Journal of Biophotonics, 2019, 12, e201900030.	1.1	6
8	Immunization with the HisAK70 DNA Vaccine Induces Resistance against Leishmania Amazonensis Infection in BALB/c Mice. Vaccines, 2019, 7, 183.	2.1	13
9	Alternative strategy for visceral leishmaniosis control: HisAK70-Salmonella Choleraesuis-pulsed dendritic cells. Comparative Immunology, Microbiology and Infectious Diseases, 2017, 54, 13-19.	0.7	12
10	Short communication: Isolation frequency of bacteria causing lymphadenitis and abscesses in small ruminants in central Spain. Small Ruminant Research, 2017, 154, 5-8.	0.6	3
11	Streptococcus ovuberis sp. nov., isolated from a subcutaneous abscess in the udder of a sheep. International Journal of Systematic and Evolutionary Microbiology, 2017, 67, 4340-4344.	0.8	9
12	Ruminants are not a reservoir of enteroaggregative Escherichia coli. Austral Journal of Veterinary Sciences, 2017, 49, 25-26.	0.2	1
13	Subtilase cytotoxin-encoding subAB2 variants in verotoxin-producing Escherichia coli strains isolated from goats and sheep. Research in Veterinary Science, 2016, 105, 74-76.	0.9	3
14	Engineering of a live Salmonella enterica serovar Choleraesuis negative-marker strain that allows serological differentiation between immunised and infected animals. Veterinary Journal, 2016, 213, 53-58.	0.6	4
15	HisAK70: progress towards a vaccine against different forms of leishmaniosis. Parasites and Vectors, 2015, 8, 629.	1.0	19
16	Characterisation of the ex vivo virulence of Leishmania infantum isolates from Phlebotomus perniciosus from an outbreak of human leishmaniosis in Madrid, Spain. Parasites and Vectors, 2014, 7, 499.	1.0	20
17	UCP2 Deficiency Helps to Restrict the Pathogenesis of Experimental Cutaneous and Visceral Leishmaniosis in Mice. PLoS Neglected Tropical Diseases, 2013, 7, e2077.	1.3	15
18	Differences in virulence gene expression between atypical enteropathogenic Escherichia coli strains isolated from diarrheic and healthy ruminants. Canadian Journal of Veterinary Research, 2013, 77, 158-60.	0.2	2

#	Article	IF	CITATIONS
19	Mitigating an undesirable immune response of inherent susceptibility to cutaneous leishmaniosis in a mouse model: the role of the pathoantigenic HISA70 DNA vaccine. Veterinary Research, 2012, 43, 59.	1.1	12
20	Comparison of ruminant and human attaching and effacing Escherichia coli (AEEC) strains. Veterinary Microbiology, 2012, 155, 341-348.	0.8	13
21	Staphylococcus aureus subsp. anaerobius isolates from different countries are clonal in nature. Veterinary Microbiology, 2011, 150, 198-202.	0.8	16
22	Mechanisms of resistance and susceptibility to experimental visceral leishmaniosis: BALB/c mouse versus syrian hamster model. Veterinary Research, 2011, 42, 39.	1.1	82
23	Phenotypic and Genotypic Characterization of Antimicrobial Resistance in Enterohemorrhagic <i>Escherichia Coli</i> and Atypical Enteropathogenic <i>E. Coli Strains</i> from Ruminants. Journal of Veterinary Diagnostic Investigation, 2011, 23, 91-95.	0.5	34
24	Subtilase Cytotoxin-Coding Genes in Verotoxin-Producing Escherichia coli Strains from Sheep and Goats Differ from Those from Cattle. Applied and Environmental Microbiology, 2011, 77, 8259-8264.	1.4	14
25	Salmonella enterica serovar Choleraesuis derivatives harbouring deletions in rpoS and phoP regulatory genes as vehicles for DNA vaccines. Veterinary Microbiology, 2010, 141, 81-88.	0.8	10
26	Association of Vt1C with Verotoxin-Producing Escherichia Coli from Goats and Sheep. Journal of Veterinary Diagnostic Investigation, 2010, 22, 332-334.	0.5	8
27	Restoring catalase activity in <i>Staphylococcus aureus </i> subsp. <i>anaerobius </i> leads to loss of pathogenicity for lambs. Veterinary Research, 2010, 41, 41.	1.1	10
28	Simultaneous lack of catalase and beta-toxin in Staphylococcus aureus leads to increased intracellular survival in macrophages and epithelial cells and to attenuated virulence in murine and ovine models. Microbiology (United Kingdom), 2009, 155, 1505-1515.	0.7	27
29	Salmonella enterica serovar Choleraesuis derivatives harbouring deletions in rpoS and phoP regulatory genes are attenuated in pigs, and survive and multiply in porcine intestinal macrophages and fibroblasts, respectively. Veterinary Microbiology, 2008, 130, 298-311.	0.8	18
30	A longitudinal study of verotoxin-producing Escherichia coli in two dairy goat herds. Veterinary Microbiology, 2008, 132, 428-434.	0.8	25
31	Characterization of Fluoroquinolone Resistance in <i>Escherichia Coli</i> Strains from Ruminants. Journal of Veterinary Diagnostic Investigation, 2008, 20, 342-345.	0.5	17
32	A spontaneous genomic deletion inListeria ivanoviiidentifies LIPI-2, a species-specific pathogenicity island encoding sphingomyelinase and numerous internalins. Molecular Microbiology, 2006, 59, 415-432.	1.2	58
33	Repression of the RcsC-YojN-RcsB phosphorelay by the IgaA protein is a requisite for Salmonella virulence. Molecular Microbiology, 2004, 53, 1437-1449.	1.2	85
34	Hpt, a bacterial homolog of the microsomal glucose- 6-phosphate translocase, mediates rapid intracellular proliferation in Listeria. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 431-436.	3.3	232
35	Regulation of Capsule Synthesis and Cell Motility in <i>Salmonella enterica</i> by the Essential Gene <i>ipaa</i> . Genetics, 2002, 162, 1513-1523.	1.2	107
36	Pathogenicity islands and virulence evolution in Listeria. Microbes and Infection, 2001, 3, 571-584.	1.0	207

#	Article	lF	CITATIONS
37	Listeria Pathogenesis and Molecular Virulence Determinants. Clinical Microbiology Reviews, 2001, 14, 584-640.	5.7	1,892
38	SmcL, a novel membrane-damaging virulence factor in Listeria. International Journal of Medical Microbiology, 2000, 290, 369-374.	1.5	30
39	The smcL gene of Listeria ivanovii encodes a sphingomyelinase C that mediates bacterial escape from the phagocytic vacuole. Molecular Microbiology, 1999, 33, 510-523.	1.2	80
40	A novel PrfA-regulated chromosomal locus, which is specific forListeria ivanovii, encodes two small, secreted internalins and contributes to virulence in mice. Molecular Microbiology, 1998, 30, 405-417.	1.2	43
41	A Gly145Ser substitution in the transcriptional activator PrfA causes constitutive overexpression of virulence factors in Listeria monocytogenes. Journal of Bacteriology, 1997, 179, 1533-1540.	1.0	155
42	Transcriptional activation of virulence genes in wild-type strains of Listeria monocytogenes in response to a change in the extracellular medium composition. Research in Microbiology, 1996, 147, 371-384.	1.0	136
43	The sulphydryl-activated cytolysin and a sphingomyelinase C are the major membrane-damaging factors involved in cooperative (CAMP-like) haemolysis of Listeria spp Research in Microbiology, 1995, 146, 303-313.	1.0	26
44	Lysergic acid diethylamide- and mescaline-induced attenuation of the effect of punishment in the rat. Science, 1976, 192, 801-803.	6.0	1,415