
Lawrence B Smart

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6134212/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Increased Accumulation of Cuticular Wax and Expression of Lipid Transfer Protein in Response to Periodic Drying Events in Leaves of Tree Tobacco. Plant Physiology, 2006, 140, 176-183.	4.8	357
2	The development of short-rotation willow in the northeastern United States for bioenergy and bioproducts, agroforestry and phytoremediation. Biomass and Bioenergy, 2006, 30, 715-727.	5.7	237
3	Genes Involved in Osmoregulation during Turgor-Driven Cell Expansion of Developing Cotton Fibers Are Differentially Regulated1. Plant Physiology, 1998, 116, 1539-1549.	4.8	177
4	MIP Genes are Down-regulated Under Drought Stress in Nicotiana glauca. Plant and Cell Physiology, 2001, 42, 686-693.	3.1	134
5	Breeding progress and preparedness for massâ€scale deployment of perennial lignocellulosic biomass crops switchgrass, miscanthus, willow and poplar. GCB Bioenergy, 2019, 11, 118-151.	5.6	116
6	Phylogenomics of the genus <i>Populus</i> reveals extensive interspecific gene flow and balancing selection. New Phytologist, 2020, 225, 1370-1382.	7.3	93
7	Yield and Woody Biomass Traits of Novel Shrub Willow Hybrids at Two Contrasting Sites. Bioenergy Research, 2013, 6, 533-546.	3.9	92
8	Untapped Potential: Opportunities and Challenges for Sustainable Bioenergy Production from Marginal Lands in the Northeast USA. Bioenergy Research, 2015, 8, 482-501.	3.9	79
9	Development and validation of genetic markers for sex and cannabinoid chemotype in <i>Cannabis sativa</i> L. GCB Bioenergy, 2020, 12, 213-222.	5.6	77
10	A willow sex chromosome reveals convergent evolution of complex palindromic repeats. Genome Biology, 2020, 21, 38.	8.8	74
11	Absence of PsaC subunit allows assembly of photosystem I core but prevents the binding of PsaD and PsaE in Synechocystis sp. PCC6803. Plant Molecular Biology, 1995, 29, 331-342.	3.9	70
12	Analysis of Biomass Composition Using High-Resolution Thermogravimetric Analysis and Percent Bark Content for the Selection of Shrub Willow Bioenergy Crop Varieties. Bioenergy Research, 2009, 2, 1-9.	3.9	61
13	Characterization of a large sex determination region in Salix purpurea L. (Salicaceae). Molecular Genetics and Genomics, 2018, 293, 1437-1452.	2.1	61
14	Expression of photosynthesis genes in the cyanobacteriumSynechocystis sp. PCC 6803:psaA-psaB andpsbA transcripts accumulate in dark-grown cells. Plant Molecular Biology, 1991, 17, 959-971.	3.9	55
15	Mutational analysis of the structure and biogenesis of the photosystem I reaction center in the cyanobacterium Synechocystis sp. PCC 6803 Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 1132-1136.	7.1	54
16	Poplar and shrub willow energy crops in the United States: field trial results from the multiyear regional feedstock partnership and yield potential maps based on the PRISMâ€ELM model. GCB Bioenergy, 2018, 10, 735-751.	5.6	54
17	Pathways to sex determination in plants: how many roads lead to Rome?. Current Opinion in Plant Biology, 2020, 54, 61-68.	7.1	54
18	A General Model to Explain Repeated Turnovers of Sex Determination in the Salicaceae. Molecular Biology and Evolution, 2021, 38, 968-980.	8.9	53

#	Article	IF	CITATIONS
19	Diversity of cuticular wax among Salix species and Populus species hybrids. Phytochemistry, 2002, 60, 715-725.	2.9	52
20	Early selection of novel triploid hybrids of shrub willow with improved biomass yield relative to diploids. BMC Plant Biology, 2014, 14, 74.	3.6	50
21	Hardwood Tree Genomics: Unlocking Woody Plant Biology. Frontiers in Plant Science, 2018, 9, 1799.	3.6	50
22	Seasonâ€long characterization of highâ€cannabinoid hemp (<i>Cannabis sativa</i> L.) reveals variation in cannabinoid accumulation, flowering time, and disease resistance. GCB Bioenergy, 2021, 13, 546-561.	5.6	50
23	Site-directed conversion of cysteine-565 to serine in PsaB of photosystem I results in the assembly of iron-sulfur [3Fe-4S] and iron-sulfur [4Fe-4S] clusters in Fx. A mixed-ligand iron-sulfur [4Fe-4S] cluster is capable of electron transfer to FA and FB. Biochemistry, 1993, 32, 4411-4419.	2.5	49
24	Ploidy Level Affects Important Biomass Traits of Novel Shrub Willow (Salix) Hybrids. Bioenergy Research, 2015, 8, 259-269.	3.9	47
25	Hydroponic Screening of Shrub Willow (<i>Salix</i> Spp.) for Arsenic Tolerance and Uptake. International Journal of Phytoremediation, 2008, 10, 515-528.	3.1	44
26	A mixed-ligand iron-sulfur cluster (C556SPaB or C565SPsaB) in the Fx-binding site leads to a decreased quantum efficiency of electron transfer in photosystem I. Biophysical Journal, 1995, 69, 1544-1553.	0.5	42
27	Differential expression of genes encoding phosphate transporters contributes to arsenic tolerance and accumulation in shrub willow (Salix spp.). Environmental and Experimental Botany, 2012, 75, 248-257.	4.2	41
28	Genotype × environment interaction analysis of North American shrub willow yield trials confirms superior performance of triploid hybrids. GCB Bioenergy, 2017, 9, 445-459.	5.6	41
29	Enzymatic saccharification of shrub willow genotypes with differing biomass composition for biofuel production. Frontiers in Plant Science, 2013, 4, 57.	3.6	39
30	Joint linkage and association mapping of complex traits in shrub willow (Salix purpurea L.). Annals of Botany, 2019, 124, 701-715.	2.9	37
31	High-resolution Thermogravimetric Analysis For Rapid Characterization of Biomass Composition and Selection of Shrub Willow Varieties. Applied Biochemistry and Biotechnology, 2008, 145, 3-11.	2.9	36
32	Population genetic structure of native versus naturalized sympatric shrub willows (<i>Salix</i> ;) Tj ETQq0 0 0 rgB	T (Qverloc	:k 10 Tf 50 2
33	Dominance and Sexual Dimorphism Pervade the Salix purpurea L. Transcriptome. Genome Biology and Evolution, 2017, 9, 2377-2394.	2.5	35
34	Effects of nitrogen fertilization in shrub willow short rotation coppice production – a quantitative review. GCB Bioenergy, 2018, 10, 548-564.	5.6	34
35	Limited effect of environmental stress on cannabinoid profiles in highâ€cannabidiol hemp (<i>Cannabis) Tj ETQq1</i>	1.0.7843 5.6	314 rgBT /Ov 32
36	Genetic inactivation of the psaB gene in Synechocystis sp. PCC 6803 disrupts assembly of photosystem I. Plant Molecular Biology, 1993, 21, 177-180.	3.9	31

#	Article	IF	CITATIONS
37	Cross-Infectivity of Powdery Mildew Isolates Originating from Hemp (<i>Cannabis sativa</i>) and Japanese Hop (<i>Humulus japonicus</i>) in New York. Plant Health Progress, 2020, 21, 47-53.	1.4	30
38	Contributions of environment and genotype to variation in shrub willow biomass composition. Industrial Crops and Products, 2017, 108, 149-161.	5.2	29
39	Biological conversion assay using Clostridium phytofermentans to estimate plant feedstock quality. Biotechnology for Biofuels, 2012, 5, 5.	6.2	28
40	Sex determination through X–Y heterogamety in Salix nigra. Heredity, 2021, 126, 630-639.	2.6	26
41	Genetic Improvement of Willow (Salix spp.) as a Dedicated Bioenergy Crop. , 2008, , 377-396.		26
42	Whole-Genome Sequences of 13 Endophytic Bacteria Isolated from Shrub Willow (Salix) Grown in Geneva, New York. Genome Announcements, 2014, 2, .	0.8	25
43	Predicting within-family variability in juvenile height growth of Salix based upon similarity among parental AFLP fingerprints. Theoretical and Applied Genetics, 2002, 105, 106-112.	3.6	21
44	Quantitative Genetics of Traits Indicative of Biomass Production and Heterosis in 34 Full-sib F1 Salix eriocephala Families. Bioenergy Research, 2008, 1, 80-90.	3.9	21
45	Collection and storage of pollen from <i>Salix</i> (Salicaceae). American Journal of Botany, 2002, 89, 248-252.	1.7	19
46	A mixed model approach for evaluating yield improvements in interspecific hybrids of shrub willow, a dedicated bioenergy crop. Industrial Crops and Products, 2017, 96, 57-70.	5.2	19
47	Morphometric relationships and their contribution to biomass and cannabinoid yield in hybrids of hemp (<i>Cannabis sativa</i>). Journal of Experimental Botany, 2021, 72, 7694-7709.	4.8	18
48	Cuticular wax composition of Salix varieties in relation to biomass productivity. Phytochemistry, 2008, 69, 396-402.	2.9	17
49	Correlations of expression of cell wall biosynthesis genes with variation in biomass composition in shrub willow (Salix spp.) biomass crops. Tree Genetics and Genomes, 2012, 8, 775-788.	1.6	17
50	Isolation of genes predominantly expressed in guard cells and epidermal cells of Nicotiana glauca. Plant Molecular Biology, 2000, 42, 857-869.	3.9	16
51	Sexual dimorphism in the dioecious willow <i>Salix purpurea</i> . American Journal of Botany, 2021, 108, 1374-1387.	1.7	14
52	Variability in pyrolysis product yield from novel shrub willow genotypes. Biomass and Bioenergy, 2015, 72, 74-84.	5.7	13
53	Genetic diversity and population structure of native, naturalized, and cultivated Salix purpurea. Tree Genetics and Genomes, 2019, 15, 1.	1.6	13
54	Genetic evidence for three discrete taxa of Melampsora (Pucciniales) affecting willows (Salix spp.) in New York State. Fungal Biology, 2014, 118, 704-720.	2.5	12

#	Article	IF	CITATIONS
55	Tolerance of novel inter-specific shrub willow hybrids to water stress. Trees - Structure and Function, 2019, 33, 1015-1026.	1.9	12
56	Evaluating the Microbiome of Hemp. Phytobiomes Journal, 2020, 4, 351-363.	2.7	12
57	Integrative genomics reveals paths to sex dimorphism in <i>Salix purpurea</i> L. Horticulture Research, 2021, 8, 170.	6.3	12
58	Shrub Willow. , 2012, , 687-708.		11
59	A Molecular and Fitness Evaluation of Commercially Available versus Locally Collected Blue Lupine <i>Lupinus perennis</i> L. Seeds for Use in Ecosystem Restoration Efforts. Restoration Ecology, 2012, 20, 456-461.	2.9	10
60	Electrical capacitance as a predictor of root dry weight in shrub willow (Salix ; Salicaceae) parents and progeny. Applications in Plant Sciences, 2016, 4, 1600031.	2.1	10
61	Developing Production Guidelines for Baby Leaf Hemp (Cannabis sativa L.) as an Edible Salad Green: Cultivar, Sowing Density and Seed Size. Agriculture (Switzerland), 2020, 10, 617.	3.1	10
62	Differential Susceptibility of Diverse <i>Salix</i> spp. to <i>Melampsora americana</i> and <i>Melampsora paradoxa</i> . Plant Disease, 2020, 104, 2949-2957.	1.4	9
63	Microbiome of Field Grown Hemp Reveals Potential Microbial Interactions With Root and Rhizosphere Soil. Frontiers in Microbiology, 2021, 12, 741597.	3.5	9
64	Mapping the sex determination region in the <i>Salix</i> F1 hybrid common parent population confirms a ZW system in six diverse species. G3: Genes, Genomes, Genetics, 2022, 12, .	1.8	9
65	Differential growth response to fertilization of ten elite shrub willow (Salix spp.) bioenergy cultivars. Trees - Structure and Function, 2018, 32, 1061-1072.	1.9	8
66	Genetic mapping of sexually dimorphic volatile and non-volatile floral secondary chemistry of a dioecious willow. Journal of Experimental Botany, 2022, 73, 6352-6366.	4.8	8
67	Discovery of Geographically Robust Hybrid Poplar Clones. Silvae Genetica, 2019, 68, 101-110.	0.8	7
68	Effects of Cold Temperature and Acclimation on Cold Tolerance and Cannabinoid Profiles of Cannabis sativa L. (Hemp). Horticulturae, 2022, 8, 531.	2.8	7
69	A second member of the Nicotiana glauca lipid transfer protein gene family, NgLTP2, encodes a divergent and differentially expressed protein. Functional Plant Biology, 2006, 33, 141.	2.1	6
70	Genotypic diversity in willow (Salix spp.) is associated with chemical and morphological polymorphism, suggesting human-assisted dissemination in the Eastern Mediterranean. Biochemical Systematics and Ecology, 2020, 91, 104081.	1.3	6
71	Heterosis for Biomass-Related Traits in Interspecific Triploid Hybrids of Willow (Salix spp.). Bioenergy Research, 2022, 15, 1042-1056.	3.9	5
72	Comparative transcriptomics and eQTL mapping of response to Melampsora americana in selected Salix purpurea F2 progeny. BMC Genomics, 2022, 23, 71.	2.8	5

#	Article	IF	CITATIONS
73	Isolation of RNA and Protein from Guard Cells of Nicotiana glauca. Plant Molecular Biology Reporter, 1999, 17, 371-383.	1.8	4
74	Transcriptome analysis of contrasting resistance to herbivory by Empoasca fabae in two shrub willow species and their hybrid progeny. PLoS ONE, 2020, 15, e0236586.	2.5	4
75	Genetic and Environmental Influences on First Rotation Shrub Willow (Salix spp.) Bark and Wood Elemental Composition. Bioenergy Research, 2020, 13, 797-809.	3.9	4
76	Yield and biomass quality of shrub willow hybrids in differing rotation lengths and spacing designs. Biomass and Bioenergy, 2021, 146, 105977.	5.7	4
77	A semiâ€commercial case study of willow biomass production in the northeastern United States. Agronomy Journal, 2021, 113, 1287-1302.	1.8	3
78	Nonadditive gene expression is correlated with nonadditive phenotypic expression in interspecific triploid hybrids of willow (<i>Salix</i> spp.). G3: Genes, Genomes, Genetics, 2022, 12, .	1.8	3
79	The Melampsora americana population on Salix purpurea in the Great Lakes region is highly diverse with a contributory influence of clonality. Phytopathology, 2021, , .	2.2	2
80	Introduction: United States Hemp Research and Education Conference special issue. GCB Bioenergy, 2021, 13, 516-516.	5.6	0