

Anna Solini

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6154791/publications.pdf>

Version: 2024-02-01

201
papers

7,307
citations

47006

47
h-index

74163

75
g-index

205
all docs

205
docs citations

205
times ranked

9940
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------|
| 1 | MG53 marks poor beta cell performance and predicts onset of type 2 diabetes in subjects with different degrees of glucose tolerance.. Diabetes and Metabolism, 2022, 48, 101292. | 2.9 | 4 |
| 2 | Challenges and opportunities in real-world evidence on the renal effects of sodium-glucose cotransporter-2 inhibitors. Diabetes, Obesity and Metabolism, 2022, 24, 177-186. | 4.4 | 11 |
| 3 | Molecular Characterization of Peritoneal Involvement in Primary Colon and Ovary Neoplasm: The Possible Clinical Meaning of the P2X7 Receptor-Inflammasome Complex. European Surgical Research, 2022, 63, 114-122. | 1.3 | 5 |
| 4 | The IGFBP3/TMEM219 pathway regulates beta cell homeostasis. Nature Communications, 2022, 13, 684. | 12.8 | 16 |
| 5 | P2X7 Receptor and Heart Function in a Mouse Model of Systemic Inflammation Due to High Fat Diet. Journal of Inflammation Research, 2022, Volume 15, 2425-2439. | 3.5 | 8 |
| 6 | The P2X7R-NLRP3 and AIM2 Inflammasome Platforms Mark the Complexity/Severity of Viral or Metabolic Liver Damage. International Journal of Molecular Sciences, 2022, 23, 7447. | 4.1 | 5 |
| 7 | Differential metabolomic signatures of declining renal function in Types 1 and 2 diabetes. Nephrology Dialysis Transplantation, 2021, 36, 1859-1866. | 0.7 | 4 |
| 8 | SGLT2 inhibitors and thiazide enhance excretion of DEHP toxic metabolites in subjects with type 2 diabetes: A randomized clinical trial. Environmental Research, 2021, 192, 110316. | 7.5 | 9 |
| 9 | Short-term impact of COVID-19 lockdown on metabolic control of patients with well-controlled type 2 diabetes: a single-centre observational study. Acta Diabetologica, 2021, 58, 431-436. | 2.5 | 47 |
| 10 | Effect of Treatment of Periodontitis on Incretin Axis in Obese and Nonobese Individuals: A Cohort Study. Journal of Clinical Endocrinology and Metabolism, 2021, 106, e74-e82. | 3.6 | 7 |
| 11 | Independent association of atherogenic dyslipidaemia with all-cause mortality in individuals with type 2 diabetes and modifying effect of gender: a prospective cohort study. Cardiovascular Diabetology, 2021, 20, 28. | 6.8 | 6 |
| 12 | Mitochondrial P2X7 Receptor Localization Modulates Energy Metabolism Enhancing Physical Performance. Function, 2021, 2, zqab005. | 2.3 | 29 |
| 13 | Normal Versus Slowly Processed Pasta and Post-Prandial Glucose Homeostasis in Healthy Subjects: A Pilot Study. Nutrients, 2021, 13, 678. | 4.1 | 0 |
| 14 | Metformin Benefits: Another Example for Alternative Energy Substrate Mechanism?. Diabetes Care, 2021, 44, 647-654. | 8.6 | 31 |
| 15 | Insulin resistance, diabetic kidney disease, and all-cause mortality in individuals with type 2 diabetes: a prospective cohort study. BMC Medicine, 2021, 19, 66. | 5.5 | 32 |
| 16 | All-cause mortality prediction models in type 2 diabetes: applicability in the early stage of disease. Acta Diabetologica, 2021, 58, 1425-1428. | 2.5 | 0 |
| 17 | miR-21 antagonism reprograms macrophage metabolism and abrogates chronic allograft vasculopathy. American Journal of Transplantation, 2021, 21, 3280-3295. | 4.7 | 14 |
| 18 | Clinical and epigenetic determinants of edentulism in type 2 diabetic subjects referring to a tertiary center. Journal of Diabetes and Its Complications, 2021, 35, 107910. | 2.3 | 2 |

| # | ARTICLE | IF | CITATIONS |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | Remdesivir, Renal Function and Short-Term Clinical Outcomes in Elderly COVID-19 Pneumonia Patients: A Single-Centre Study. <i>Clinical Interventions in Aging</i> , 2021, Volume 16, 1037-1046. | 2.9 | 10 |
| 20 | P2X7 receptor/NLRP3 inflammasome complex and α -synuclein in peripheral blood mononuclear cells: a prospective study in newly-diagnosed, treatment-naïve Parkinson's disease. <i>European Journal of Neurology</i> , 2021, 28, 2648-2656. | 3.3 | 12 |
| 21 | The P2X7 Receptor: A Promising Pharmacological Target in Diabetic Retinopathy. <i>International Journal of Molecular Sciences</i> , 2021, 22, 7110. | 4.1 | 17 |
| 22 | Physical activity as a proxy to ameliorate inflammation in patients with type 2 diabetes and periodontal disease at high cardiovascular risk. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 2199-2209. | 2.6 | 7 |
| 23 | A long-term nationwide study on chronic kidney disease-related mortality in Italy: trends and associated comorbidity. <i>Journal of Nephrology</i> , 2021, , 1. | 2.0 | 4 |
| 24 | Antiangiogenic Drugs in NASH: Evidence of a Possible New Therapeutic Approach. <i>Pharmaceuticals</i> , 2021, 14, 995. | 3.8 | 9 |
| 25 | Protein and amino acids in nonalcoholic fatty liver disease. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2021, 24, 96-101. | 2.5 | 42 |
| 26 | Glucagon-Like Peptide-1 Receptor Agonists' Use in Clinical Practice. <i>Advances in Chronic Kidney Disease</i> , 2021, 28, 328-336. | 1.4 | 5 |
| 27 | Renal hyperfiltration is independently associated with increased all-cause mortality in individuals with type 2 diabetes: a prospective cohort study. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001481. | 2.8 | 22 |
| 28 | Renoprotection with SGLT2 inhibitors in type 2 diabetes over a spectrum of cardiovascular and renal risk. <i>Cardiovascular Diabetology</i> , 2020, 19, 196. | 6.8 | 52 |
| 29 | Phenotyping individuals with newly-diagnosed type 2 diabetes at risk for all-cause mortality: a single centre observational, prospective study. <i>Diabetology and Metabolic Syndrome</i> , 2020, 12, 47. | 2.7 | 2 |
| 30 | miR-130a and Tgfb ² Content in Extracellular Vesicles Derived from the Serum of Subjects at High Cardiovascular Risk Predicts their In-Vivo Angiogenic Potential. <i>Scientific Reports</i> , 2020, 10, 706. | 3.3 | 13 |
| 31 | Association between On-Treatment Haemoglobin A1c and All-Cause Mortality in Individuals with Type 2 Diabetes: Importance of Personalized Goals and Type of Anti-Hyperglycaemic Treatment. <i>Journal of Clinical Medicine</i> , 2020, 9, 246. | 2.4 | 2 |
| 32 | Cardiovascular protection with sodium-glucose cotransporter 2 inhibitors in type 2 diabetes: Does it apply to all patients?. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 1481-1495. | 4.4 | 17 |
| 33 | Sodium-glucose cotransporter 2 inhibitors antagonize lipotoxicity in human myeloid angiogenic cells and ADP-dependent activation in human platelets: potential relevance to prevention of cardiovascular events. <i>Cardiovascular Diabetology</i> , 2020, 19, 46. | 6.8 | 43 |
| 34 | Diabetes and the Kidney. <i>Endocrinology</i> , 2020, , 203-230. | 0.1 | 0 |
| 35 | Effectiveness of dapagliflozin versus comparators on renal endpoints in the real world: A multicentre retrospective study. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 252-260. | 4.4 | 33 |
| 36 | The Effects of Dapagliflozin on Systemic and Renal Vascular Function Display an Epigenetic Signature. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 4253-4263. | 3.6 | 57 |

| # | ARTICLE | IF | CITATIONS |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 37 | Similar effectiveness of dapagliflozin and GLP-1 receptor agonists concerning combined endpoints in routine clinical practice: A multicentre retrospective study. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 1886-1894. | 4.4 | 17 |
| 38 | Is resistant hypertension an independent predictor of all-cause mortality in individuals with type 2 diabetes? A prospective cohort study. <i>BMC Medicine</i> , 2019, 17, 83. | 5.5 | 9 |
| 39 | Phthalates Exposure as Determinant of Albuminuria in Subjects With Type 2 Diabetes: A Cross-Sectional Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 1491-1499. | 3.6 | 16 |
| 40 | Role of the P2X7 receptor in the pathogenesis of type 2 diabetes and its microvascular complications. <i>Current Opinion in Pharmacology</i> , 2019, 47, 75-81. | 3.5 | 35 |
| 41 | Pharmacological blockade of the P2X7 receptor reverses retinal damage in a rat model of type 1 diabetes. <i>Acta Diabetologica</i> , 2019, 56, 1031-1036. | 2.5 | 30 |
| 42 | Alterations in Carotid Parameters in ApoE ^{-/-} Mice Treated with a High-Fat Diet: A Micro-ultrasound Analysis. <i>Ultrasound in Medicine and Biology</i> , 2019, 45, 980-988. | 1.5 | 2 |
| 43 | Periodontitis affects glucoregulatory hormones in severely obese individuals. <i>International Journal of Obesity</i> , 2019, 43, 1125-1129. | 3.4 | 12 |
| 44 | Renal Resistive Index Predicts Post-Bariatric Surgery Renal Outcome in Nondiabetic Individuals with Severe Obesity. <i>Obesity</i> , 2019, 27, 68-74. | 3.0 | 10 |
| 45 | DNA methylation of genes regulating appetite and prediction of weight loss after bariatric surgery in obese individuals. <i>Journal of Endocrinological Investigation</i> , 2019, 42, 37-44. | 3.3 | 6 |
| 46 | Diabetes and the Kidney. <i>Endocrinology</i> , 2019, , 1-28. | 0.1 | 0 |
| 47 | P2X receptor-ion channels in the inflammatory response in adipose tissue and pancreas – potential triggers in onset of type 2 diabetes?. <i>Current Opinion in Immunology</i> , 2018, 52, 1-7. | 5.5 | 30 |
| 48 | Haemoglobin A1c variability is a strong, independent predictor of all-cause mortality in patients with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2018, 20, 1885-1893. | 4.4 | 45 |
| 49 | PDGF-BB Carried by Endothelial Cell-Derived Extracellular Vesicles Reduces Vascular Smooth Muscle Cell Apoptosis in Diabetes. <i>Diabetes</i> , 2018, 67, 704-716. | 0.6 | 38 |
| 50 | The level of physical training modulates cytokine levels through P2X7 receptor in healthy subjects. <i>European Journal of Clinical Investigation</i> , 2018, 48, e12880. | 3.4 | 13 |
| 51 | A systematic review and meta-analysis of epidemiologic observational evidence on the effect of periodontitis on diabetes An update of the <sc>EFP</sc> review. <i>Journal of Clinical Periodontology</i> , 2018, 45, 167-187. | 4.9 | 245 |
| 52 | Defining the contribution of chronic kidney disease to all-cause mortality in patients with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicenter Study. <i>Acta Diabetologica</i> , 2018, 55, 603-612. | 2.5 | 33 |
| 53 | Future Perspectives on GLP-1 Receptor Agonists and GLP-1/glucagon Receptor Co-agonists in the Treatment of NAFLD. <i>Frontiers in Endocrinology</i> , 2018, 9, 649. | 3.5 | 65 |
| 54 | Phenotyping normal kidney function in elderly patients with type 2 diabetes: a cross-sectional multicentre study. <i>Acta Diabetologica</i> , 2018, 55, 1121-1129. | 2.5 | 2 |

| # | ARTICLE | IF | CITATIONS |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 55 | Ultrasonographic Characterization of the <i>db/db</i> Mouse: An Animal Model of Metabolic Abnormalities. <i>Journal of Diabetes Research</i> , 2018, 2018, 1-9. | 2.3 | 12 |
| 56 | Islet-Derived eATP Fuels Autoreactive CD8+ T Cells and Facilitates the Onset of Type 1 Diabetes. <i>Diabetes</i> , 2018, 67, 2038-2053. | 0.6 | 17 |
| 57 | Diabetes and the Kidney. <i>Endocrinology</i> , 2018, , 1-27. | 0.1 | 0 |
| 58 | Non-albuminuric renal impairment is a strong predictor of mortality in individuals with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian multicentre study. <i>Diabetologia</i> , 2018, 61, 2277-2289. | 6.3 | 83 |
| 59 | A renal genetic risk score (GRS) is associated with kidney dysfunction in people with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018, 144, 137-143. | 2.8 | 5 |
| 60 | Diabetes and the Kidney. <i>Endocrinology</i> , 2018, , 203-229. | 0.1 | 0 |
| 61 | Clozapine as the most efficacious antipsychotic for activating ERK 1/2 kinases: Role of 5-HT 2A receptor agonism. <i>European Neuropsychopharmacology</i> , 2017, 27, 383-398. | 0.7 | 44 |
| 62 | Sodium-glucose co-transporter (SGLT)2 and SGLT1 renal expression in patients with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 1289-1294. | 4.4 | 66 |
| 63 | Dapagliflozin modulates glucagon secretion in an SGLT2-independent manner in murine alpha cells. <i>Diabetes and Metabolism</i> , 2017, 43, 512-520. | 2.9 | 51 |
| 64 | The P2X7 receptor-NLRP3 inflammasome complex predicts the development of non-Hodgkin's lymphoma in Sjogren's syndrome: a prospective, observational, single-centre study. <i>Journal of Internal Medicine</i> , 2017, 282, 175-186. | 6.0 | 49 |
| 65 | Determinants of glomerular filtration rate following bariatric surgery in individuals with severe, otherwise uncomplicated, obesity: an observational, prospective study. <i>Acta Diabetologica</i> , 2017, 54, 593-598. | 2.5 | 7 |
| 66 | Silent coronary heart disease in patients with type 2 diabetes: application of a screening approach in a follow-up study. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 952-957. | 2.3 | 5 |
| 67 | SGLT inhibition in T1DM – definite benefit with manageable risk. <i>Nature Reviews Endocrinology</i> , 2017, 13, 698-699. | 9.6 | 6 |
| 68 | Rationale and design of the DARWIN-T2D (Dapagliflozin Real World evidence in Type 2 Diabetes). <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2017, 27, 1089-1097. | 2.6 | 26 |
| 69 | Heterozygosity for the rs696217 SNP in the Preproghrelin Gene Predicts Weight Loss After Bariatric Surgery in Severely Obese Individuals. <i>Obesity Surgery</i> , 2017, 27, 961-967. | 2.1 | 18 |
| 70 | Deficiency of the Purinergic Receptor 2X ₇ Attenuates Nonalcoholic Steatohepatitis Induced by High-Fat Diet: Possible Role of the NLRP3 Inflammasome. <i>Oxidative Medicine and Cellular Longevity</i> , 2017, 2017, 1-14. | 4.0 | 23 |
| 71 | Dapagliflozin acutely improves endothelial dysfunction, reduces aortic stiffness and renal resistive index in type 2 diabetic patients: a pilot study. <i>Cardiovascular Diabetology</i> , 2017, 16, 138. | 6.8 | 274 |
| 72 | P2X7 Receptor and APOE Polymorphisms and Survival from Heart Failure: A Prospective Study in Frail Patients in a Geriatric Unit. , 2017, 8, 434. | | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 73 | The EMPA-REG outcome study: critical appraisal and potential clinical implications. <i>Cardiovascular Diabetology</i> , 2016, 15, 85. | 6.8 | 16 |
| 74 | Role of SGLT2 inhibitors in the treatment of type 2 diabetes mellitus. <i>Acta Diabetologica</i> , 2016, 53, 863-870. | 2.5 | 18 |
| 75 | Syphilis iridocyclitis in a patient with type 1 diabetes. <i>Journal of Diabetes Investigation</i> , 2016, 7, 641-644. | 2.4 | 2 |
| 76 | What Should Be the Target Blood Pressure in Elderly Patients With Diabetes?. <i>Diabetes Care</i> , 2016, 39, S234-S243. | 8.6 | 21 |
| 77 | Metabolic and Hormonal Determinants of Glomerular Filtration Rate and Renal Hemodynamics in Severely Obese Individuals. <i>Obesity Facts</i> , 2016, 9, 310-320. | 3.4 | 15 |
| 78 | [OP.7B.10] EFFECT OF BARIATRIC SURGERY ON VASCULAR AND RENAL BIOMARKERS IN MORBIDLY OBESE, NORMOTENSIVE, NON-DIABETIC PATIENTS. <i>Journal of Hypertension</i> , 2016, 34, e88-e89. | 0.5 | 2 |
| 79 | Extra-glycaemic properties of empagliflozin. <i>Diabetes/Metabolism Research and Reviews</i> , 2016, 32, 230-237. | 4.0 | 7 |
| 80 | Prediction of Declining Renal Function and Albuminuria in Patients With Type 2 Diabetes by Metabolomics. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016, 101, 696-704. | 3.6 | 62 |
| 81 | Saxagliptin prevents vascular remodeling and oxidative stress in db/db mice. Role of endothelial nitric oxide synthase uncoupling and cyclooxygenase. <i>Vascular Pharmacology</i> , 2016, 76, 62-71. | 2.1 | 25 |
| 82 | Obesity reduces the pro-angiogenic potential of adipose tissue stem cell-derived extracellular vesicles (EVs) by impairing miR-126 content: impact on clinical applications. <i>International Journal of Obesity</i> , 2016, 40, 102-111. | 3.4 | 95 |
| 83 | Early treatment with hydroxychloroquine prevents the development of endothelial dysfunction in a murine model of systemic lupus erythematosus. <i>Arthritis Research and Therapy</i> , 2015, 17, 277. | 3.5 | 55 |
| 84 | Predictive value of dynamic renal resistive index (drin) for renal outcome in type 2 diabetes and essential hypertension: a prospective study. <i>Cardiovascular Diabetology</i> , 2015, 14, 63. | 6.8 | 22 |
| 85 | Tumour necrosis factor-alpha participates on the endothelin-1/nitric oxide imbalance in small arteries from obese patients: role of perivascular adipose tissue. <i>European Heart Journal</i> , 2015, 36, 784-794. | 2.2 | 127 |
| 86 | Hormone replacement therapy, renal function and heart ultrasonographic parameters in postmenopausal women: an observational study. <i>International Journal of Clinical Practice</i> , 2015, 69, 632-637. | 1.7 | 7 |
| 87 | Independent correlates of urinary albumin excretion within the normoalbuminuric range in patients with type 2 diabetes: The Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicentre Study. <i>Acta Diabetologica</i> , 2015, 52, 971-981. | 2.5 | 8 |
| 88 | The Dark Side of Extracellular ATP in Kidney Diseases. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 1007-1016. | 6.1 | 72 |
| 89 | Acute effects of different degrees of ultra-endurance exercise on systemic inflammatory responses. <i>Internal Medicine Journal</i> , 2015, 45, 74-79. | 0.8 | 41 |
| 90 | Hypertriglyceridemia Is Independently Associated with Renal, but Not Retinal Complications in Subjects with Type 2 Diabetes: A Cross-Sectional Analysis of the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicenter Study. <i>PLoS ONE</i> , 2015, 10, e0125512. | 2.5 | 30 |

| # | ARTICLE | IF | CITATIONS |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------|
| 91 | Genetic interaction of P2X7 receptor and VEGFR-2 polymorphisms identifies a favorable prognostic profile in prostate cancer patients. <i>Oncotarget</i> , 2015, 6, 28743-28754. | 1.8 | 21 |
| 92 | The complex P2X7 receptor/inflammasome in perivascular fat tissue of heavy smokers. <i>European Journal of Clinical Investigation</i> , 2014, 44, 295-302. | 3.4 | 32 |
| 93 | Role of Podocyte B7-1 in Diabetic Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2014, 25, 1415-1429. | 6.1 | 114 |
| 94 | Resistant hypertension in patients with type 2 diabetes. <i>Journal of Hypertension</i> , 2014, 32, 2401-2410. | 0.5 | 35 |
| 95 | Distribution of cardiovascular disease and retinopathy in patients with type 2 diabetes according to different classification systems for chronic kidney disease: a cross-sectional analysis of the renal insufficiency and cardiovascular events (RIACE) Italian multicenter study. <i>Cardiovascular Diabetology</i> , 2014, 13, 59. | 6.8 | 24 |
| 96 | The ideal blood pressure target to prevent cardiovascular disease in type 2 diabetes: A neutral viewpoint. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014, 24, 577-584. | 2.6 | 4 |
| 97 | Blood pressure variability: A new target to slow the progression of vascular damage in type 2 diabetes?. <i>Journal of Diabetes and Its Complications</i> , 2014, 28, 117-118. | 2.3 | 1 |
| 98 | Chronic kidney disease in type 2 diabetes: Lessons from the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicentre Study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014, 24, 815-822. | 2.6 | 51 |
| 99 | Impact of mild to moderate reductions of glomerular filtration rate on coronary artery disease severity. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014, 24, 681-688. | 2.6 | 18 |
| 100 | Comparison of Agents That Affect Aldosterone Action. <i>Seminars in Nephrology</i> , 2014, 34, 285-306. | 1.6 | 28 |
| 101 | HbA1c Variability as an Independent Correlate of Nephropathy, but Not Retinopathy, in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2013, 36, 2301-2310. | 8.6 | 130 |
| 102 | Gender differences in cardiovascular disease risk factors, treatments and complications in patients with type 2 diabetes: the RIACE Italian multicentre study. <i>Journal of Internal Medicine</i> , 2013, 274, 176-191. | 6.0 | 111 |
| 103 | Hemoglobin A1c variability as an independent correlate of cardiovascular disease in patients with type 2 diabetes: a cross-sectional analysis of the Renal Insufficiency and Cardiovascular Events (RIACE) Italian Multicenter Study. <i>Cardiovascular Diabetology</i> , 2013, 12, 98. | 6.8 | 61 |
| 104 | Adipocytokines mark insulin sensitivity in euthyroid Hashimoto's patients. <i>Acta Diabetologica</i> , 2013, 50, 73-80. | 2.5 | 4 |
| 105 | How can resistant hypertension be identified and prevented?. <i>Nature Reviews Cardiology</i> , 2013, 10, 293-296. | 13.7 | 9 |
| 106 | P2X7 receptor polymorphisms do not influence endothelial function and vascular tone in neo-diagnosed, treatment-naive essential hypertensive patients. <i>Journal of Hypertension</i> , 2013, 31, 2362-2369. | 0.5 | 11 |
| 107 | The purinergic P2X7 receptor participates in renal inflammation and injury induced by high-fat diet: possible role of NLRP3 inflammasome activation. <i>Journal of Pathology</i> , 2013, 231, 342-353. | 4.5 | 99 |
| 108 | The P2X7 receptor-inflammasome complex has a role in modulating the inflammatory response in primary Sjögren's syndrome. <i>Journal of Internal Medicine</i> , 2013, 274, 480-489. | 6.0 | 74 |

| # | ARTICLE | IF | CITATIONS |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 109 | VEGF-A polymorphisms predict progression-free survival among advanced castration-resistant prostate cancer patients treated with metronomic cyclophosphamide. <i>British Journal of Cancer</i> , 2013, 109, 957-964. | 6.4 | 41 |
| 110 | Age, Renal Dysfunction, Cardiovascular Disease, and Antihyperglycemic Treatment in Type 2 Diabetes Mellitus: Findings from the Renal Insufficiency and Cardiovascular Events Italian Multicenter Study. <i>Journal of the American Geriatrics Society</i> , 2013, 61, 1253-1261. | 2.6 | 65 |
| 111 | THU0304â€¦Gross Cystic Disease Fluid Protein-15(GCDFP-15)/Prolactin-Inducible Protein (PIP): A Functional Salivary Biomarker for Primary Sjögren's Syndrome?. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A268.4-A269. | 0.9 | 0 |
| 112 | Diverging Association of Reduced Glomerular Filtration Rate and Albuminuria With Coronary and Noncoronary Events in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 143-149. | 8.6 | 107 |
| 113 | Rate and Determinants of Association Between Advanced Retinopathy and Chronic Kidney Disease in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 2317-2323. | 8.6 | 106 |
| 114 | Resistance artery mechanics and composition in angiotensin II-infused mice: effects of cyclooxygenase-1 inhibition. <i>European Heart Journal</i> , 2012, 33, 2225-2234. | 2.2 | 28 |
| 115 | SGLT2 inhibition in diabetes mellitus: rationale and clinical prospects. <i>Nature Reviews Endocrinology</i> , 2012, 8, 495-502. | 9.6 | 364 |
| 116 | Adipocytokine levels mark endothelial function in normotensive individuals. <i>Cardiovascular Diabetology</i> , 2012, 11, 103. | 6.8 | 25 |
| 117 | High prevalence of advanced retinopathy in patients with type 2 diabetes from the Renal Insufficiency And Cardiovascular Events (RIACE) Italian Multicenter Study. <i>Diabetes Research and Clinical Practice</i> , 2012, 98, 329-337. | 2.8 | 29 |
| 118 | Combination Therapy in Hypertension. <i>Current Cardiovascular Risk Reports</i> , 2012, 6, 291-298. | 2.0 | 1 |
| 119 | The Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation provides a better definition of cardiovascular burden associated with CKD than the Modification of Diet in Renal Disease (MDRD) Study formula in subjects with type 2 diabetes. <i>Atherosclerosis</i> , 2011, 218, 194-199. | 0.8 | 55 |
| 120 | Adipocyte P2X7 receptors expression: A role in modulating inflammatory response in subjects with metabolic syndrome?. <i>Atherosclerosis</i> , 2011, 219, 552-558. | 0.8 | 43 |
| 121 | Clinical significance of nonalbuminuric renal impairment in type 2 diabetes. <i>Journal of Hypertension</i> , 2011, 29, 1802-1809. | 0.5 | 198 |
| 122 | Angiotensin-II and rosuvastatin influence matrix remodeling in human mesangial cells via metalloproteinase modulation. <i>Journal of Hypertension</i> , 2011, 29, 1930-1939. | 0.5 | 15 |
| 123 | Reproducibility of albuminuria in type 2 diabetic subjects. Findings from the Renal Insufficiency And Cardiovascular Events (RIACE) study. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 3950-3954. | 0.7 | 65 |
| 124 | Pathophysiology, Prevention and Management of Chronic Kidney Disease in the Hypertensive Patient With Diabetes Mellitus. <i>Journal of Clinical Hypertension</i> , 2011, 13, 252-257. | 2.0 | 12 |
| 125 | Dynamic evaluation of renal resistive index in normoalbuminuric patients with newly diagnosed hypertension or type 2 diabetes. <i>Diabetologia</i> , 2011, 54, 2430-2439. | 6.3 | 48 |
| 126 | Effect of a fatty meal on inflammatory markers in healthy volunteers with a family history of type 2 diabetes. <i>British Journal of Nutrition</i> , 2011, 106, 364-368. | 2.3 | 18 |

| # | ARTICLE | IF | CITATIONS |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 127 | RAS Blockade for Every Diabetic Patient: Pro and Con. <i>Diabetes Care</i> , 2011, 34, S320-S324. | 8.6 | 10 |
| 128 | Pattern of expression of inflammatory markers in adipose tissue of untreated hypertensive patients. <i>Journal of Hypertension</i> , 2010, 28, 1459-1465. | 0.5 | 16 |
| 129 | Hypothyroidism and Intermediate Metabolism: A Complex Relationship. <i>Thyroid</i> , 2010, 20, 837-839. | 4.5 | 7 |
| 130 | Soluble Human Leukocyte Antigen-G Expression and Glucose Tolerance in Subjects with Different Degrees of Adiposity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010, 95, 3342-3346. | 3.6 | 25 |
| 131 | Retinol-Binding Protein-4 in Women With Untreated Essential Hypertension. <i>American Journal of Hypertension</i> , 2009, 22, 1001-1006. | 2.0 | 61 |
| 132 | Soluble CD40 Ligand Levels in Essential Hypertensive Men: Evidence of a Possible Role of Insulin Resistance. <i>American Journal of Hypertension</i> , 2009, 22, 1007-1013. | 2.0 | 14 |
| 133 | Extracellular Adenosine 5â€™-Triphosphate Modulates Insulin Secretion via Functionally Active Purinergic Receptors of X and Y Subtype. <i>Endocrinology</i> , 2009, 150, 2596-2602. | 2.8 | 17 |
| 134 | Short-term Acute Hyperinsulinemia and Prothrombotic Factors in Subjects with Normal Glucose Tolerance. <i>Hormone and Metabolic Research</i> , 2009, 41, 568-572. | 1.5 | 2 |
| 135 | 1513A>C Polymorphism in the P2X7 Receptor Gene in Patients with Papillary Thyroid Cancer: Correlation with Histological Variants and Clinical Parameters. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009, 94, 695-698. | 3.6 | 43 |
| 136 | Family history of hypertension, anthropometric parameters and markers of early atherosclerosis in young healthy individuals. <i>Journal of Human Hypertension</i> , 2009, 23, 801-807. | 2.2 | 17 |
| 137 | TIMP3 Is Reduced in Atherosclerotic Plaques From Subjects With Type 2 Diabetes and Increased by SirT1. <i>Diabetes</i> , 2009, 58, 2396-2401. | 0.6 | 132 |
| 138 | Effects of different LDL particles on inflammatory molecules in human mesangial cells. <i>Diabetologia</i> , 2008, 51, 2117-2125. | 6.3 | 35 |
| 139 | Rosiglitazone increases matrix production and quenches inflammation: studies in human cells. <i>Diabetes/Metabolism Research and Reviews</i> , 2008, 24, 197-204. | 4.0 | 4 |
| 140 | Hyperinsulinemia and insulin resistance are independently associated with plasma lipids, uric acid and blood pressure in non-diabetic subjects. The GISIR database. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2008, 18, 624-631. | 2.6 | 67 |
| 141 | Effect of statins on soluble CD40 ligand in hypercholesterolemic Type 2 diabetic patients. <i>Journal of Endocrinological Investigation</i> , 2008, 31, 660-665. | 3.3 | 18 |
| 142 | Increased P2X7 Receptor Expression and Function in Thyroid Papillary Cancer: A New Potential Marker of the Disease?. <i>Endocrinology</i> , 2008, 149, 389-396. | 2.8 | 123 |
| 143 | Multiple P2X receptors are involved in the modulation of apoptosis in human mesangial cells: evidence for a role of P2X4. <i>American Journal of Physiology - Renal Physiology</i> , 2007, 292, F1537-F1547. | 2.7 | 30 |
| 144 | Effects of endothelin-1 on fibroblasts from type 2 diabetic patients: Possible role in wound healing and tissue repair. <i>Growth Factors</i> , 2007, 25, 392-399. | 1.7 | 13 |

| # | ARTICLE | IF | CITATIONS |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 145 | Increased sensitivity to extracellular ATP of fibroblasts from patients affected by systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2007, 66, 1124-1125. | 0.9 | 9 |
| 146 | Correspondence Between the International Diabetes Federation Criteria for Metabolic Syndrome and Insulin Resistance in a Cohort of Italian Nondiabetic Caucasians: The GISIR database. <i>Diabetes Care</i> , 2007, 30, e33-e33. | 8.6 | 4 |
| 147 | Acute retinal ganglion cell injury caused by intraocular pressure spikes is mediated by endogenous extracellular ATP. <i>European Journal of Neuroscience</i> , 2007, 25, 2741-2754. | 2.6 | 128 |
| 148 | High glucose and homocysteine synergistically affect the metalloproteinasesâ€tissue inhibitors of metalloproteinases pattern, but not TGFB expression, in human fibroblasts. <i>Diabetologia</i> , 2006, 49, 2499-2506. | 6.3 | 26 |
| 149 | Acute Vascular Events and Electrolytes Variations in Elderly Patients. <i>Hormone and Metabolic Research</i> , 2006, 38, 197-202. | 1.5 | 13 |
| 150 | Effect of short-term folic acid supplementation on insulin sensitivity and inflammatory markers in overweight subjects. <i>International Journal of Obesity</i> , 2006, 30, 1197-1202. | 3.4 | 105 |
| 151 | Enhanced angiotensin II-mediated effects in fibroblasts of patients with familial hypercholesterolemia. <i>Journal of Hypertension</i> , 2005, 23, 367-374. | 0.5 | 10 |
| 152 | Purinergic modulation of mesangial extracellular matrix production: Role in diabetic and other glomerular diseases. <i>Kidney International</i> , 2005, 67, 875-885. | 5.2 | 63 |
| 153 | Selective Insulin Resistance Affecting Nitric Oxide Release But Not Plasminogen Activator Inhibitor-1 Synthesis in Fibroblasts From Insulin-Resistant Individuals. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005, 25, 2392-2397. | 2.4 | 18 |
| 154 | Antihypertensive Treatment and Multifactorial Approach for Renal Protection in Diabetes. <i>Journal of the American Society of Nephrology: JASN</i> , 2005, 16, S18-S21. | 6.1 | 27 |
| 155 | Extracellular Adenosine 5â€-Triphosphate Modulates Interleukin-6 Production by Human Thyrocytes through Functional Purinergic P2 Receptors. <i>Endocrinology</i> , 2005, 146, 3172-3178. | 2.8 | 21 |
| 156 | Enhanced P2X 7 Activity in Human Fibroblasts From Diabetic Patients. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2004, 24, 1240-1245. | 2.4 | 50 |
| 157 | Lipoprotein lipase gene variants and progression of nephropathy in hypercholesterolaemic patients with type 2 diabetes. <i>Journal of Internal Medicine</i> , 2004, 256, 30-36. | 6.0 | 11 |
| 158 | Defective P2Y purinergic receptor function: A possible novel mechanism for impaired glucose transport. <i>Journal of Cellular Physiology</i> , 2003, 197, 435-444. | 4.1 | 26 |
| 159 | Effect of metabolic control on homocysteine levels in type 2 diabetic patients: a 3-year follow-up. <i>Journal of Internal Medicine</i> , 2003, 254, 264-271. | 6.0 | 23 |
| 160 | Atorvastatin improves metabolic control and endothelial function in Type 2 diabetic patients: A placebo-controlled study. <i>Journal of Endocrinological Investigation</i> , 2003, 26, 73-78. | 3.3 | 44 |
| 161 | PC-1 Amino Acid Variant Q121 Is Associated With a Lower Glomerular Filtration Rate in Type 2 Diabetic Patients With Abnormal Albumin Excretion Rates. <i>Diabetes Care</i> , 2003, 26, 2898-2902. | 8.6 | 17 |
| 162 | Metabolic Profile in Patients with Benign Prostate Hyperplasia or Prostate Cancer and Normal Glucose Tolerance. <i>Hormone and Metabolic Research</i> , 2003, 35, 296-300. | 1.5 | 13 |

| # | ARTICLE | IF | CITATIONS |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 163 | The Angiotensin-Converting Enzyme DD Genotype Is Associated With Glomerulopathy Lesions in Type 2 Diabetes. <i>Diabetes</i> , 2002, 51, 251-255. | 0.6 | 43 |
| 164 | P2 receptors: new potential players in atherosclerosis. <i>British Journal of Pharmacology</i> , 2002, 135, 831-842. | 5.4 | 113 |
| 165 | Plasma homocysteine, methylenetetrahydrofolate reductase mutation and carotid damage in elderly healthy women. <i>Atherosclerosis</i> , 2001, 157, 175-180. | 0.8 | 41 |
| 166 | Role of diabetes in influencing leptin concentration in elderly overweight patients. <i>European Journal of Endocrinology</i> , 2001, 145, 173-179. | 3.7 | 8 |
| 167 | ACE Genotype and Endothelium-Dependent Vasodilation of Conduit Arteries and Forearm Microcirculation in Humans. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2001, 21, 1313-1319. | 2.4 | 20 |
| 168 | A Defect in Glycogen Synthesis Characterizes Insulin Resistance in Hypertensive Patients With Type 2 Diabetes. <i>Hypertension</i> , 2001, 37, 1492-1496. | 2.7 | 15 |
| 169 | Factors influencing plasma homocysteine levels in type 2 diabetes. <i>Diabetes Care</i> , 2000, 23, 420-421. | 8.6 | 17 |
| 170 | High glucose modulates P2X 7 receptor-mediated function in human primary fibroblasts. <i>Diabetologia</i> , 2000, 43, 1248-1256. | 6.3 | 51 |
| 171 | The relationship of plasma glucose and electrocardiographic parameters in elderly women with different degrees of glucose tolerance. <i>Aging Clinical and Experimental Research</i> , 2000, 12, 249-255. | 2.9 | 8 |
| 172 | Polymorphisms of angiotensin-converting enzyme and angiotensinogen genes in type 2 diabetic sibships in relation to albumin excretion rate. <i>American Journal of Kidney Diseases</i> , 1999, 34, 1002-1009. | 1.9 | 24 |
| 173 | Human primary fibroblasts in vitro express a purinergic P2X7 receptor coupled to ion fluxes, microvesicle formation and IL-6 release. <i>Journal of Cell Science</i> , 1999, 112, 297-305. | 2.0 | 134 |
| 174 | Human primary fibroblasts in vitro express a purinergic P2X7 receptor coupled to ion fluxes, microvesicle formation and IL-6 release. <i>Journal of Cell Science</i> , 1999, 112 (Pt 3), 297-305. | 2.0 | 52 |
| 175 | Elevated PC-1 content in cultured skin fibroblasts correlates with decreased in vivo and in vitro insulin action in nondiabetic subjects: evidence that PC-1 may be an intrinsic factor in impaired insulin receptor signaling. <i>Diabetes</i> , 1998, 47, 1095-1100. | 0.6 | 66 |
| 176 | Derangements in Protein Metabolism Induced by Type I Diabetes mellitus. <i>Mineral and Electrolyte Metabolism</i> , 1998, 24, 41-46. | 1.1 | 6 |
| 177 | Protein Metabolism in Human Obesity: Relationship with Glucose and Lipid Metabolism and with Visceral Adipose Tissue. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997, 82, 2552-2558. | 3.6 | 46 |
| 178 | Glycosaminoglycans Delay the Progression of Nephropathy in NIDDM. <i>Diabetes Care</i> , 1997, 20, 819-823. | 8.6 | 74 |
| 179 | and countertransport activity in hypertensive non-insulin-dependent diabetic patients: Role in insulin resistance and antihypertensive treatment. <i>Metabolism: Clinical and Experimental</i> , 1997, 46, 1316-1323. | 3.4 | 20 |
| 180 | Protein Metabolism in Human Obesity: Relationship with Glucose and Lipid Metabolism and with Visceral Adipose Tissue. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997, 82, 2552-2558. | 3.6 | 43 |

| # | ARTICLE | IF | CITATIONS |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 181 | Intracellular calcium handling by fibroblasts from non-insulin dependent diabetic patients with and without hypertension and microalbuminuria. <i>Kidney International</i> , 1996, 50, 618-626. | 5.2 | 6 |
| 182 | Hypertension, Cardiovascular Disease, Diabetes Mellitus, and Diabetic Nephropathy: Role of Insulin Resistance. , 1996, , 61-74. | | 2 |
| 183 | Insulin Sensitivity Is Not Impaired In Mexican-American Women Without a Family History Of Diabetes. <i>Diabetes Care</i> , 1995, 18, 825-833. | 8.6 | 8 |
| 184 | Lipoprotein abnormalities in non-insulin-dependent diabetic patients with impaired extrahepatic insulin sensitivity, hypertension, and microalbuminuria.. <i>Arteriosclerosis and Thrombosis: A Journal of Vascular Biology</i> , 1994, 14, 911-916. | 3.9 | 16 |
| 185 | Impaired Insulin-Induced Glucose Uptake by Extrahepatic Tissue Is Hallmark of NIDDM Patients Who Have or Will Develop Hypertension and Microalbuminuria. <i>Diabetes</i> , 1994, 43, 491-499. | 0.6 | 62 |
| 186 | Peripheral, rather than hepatic, insulin resistance and atherogenic lipoprotein phenotype predict cardiovascular complications in NIDDM. <i>European Journal of Clinical Investigation</i> , 1994, 24, 258-266. | 3.4 | 21 |
| 187 | Impaired insulin-induced glucose uptake by extrahepatic tissue is hallmark of NIDDM patients who have or will develop hypertension and microalbuminuria. <i>Diabetes</i> , 1994, 43, 491-499. | 0.6 | 27 |
| 188 | Role of hyperglycemia and insulin resistance in determining sodium retention in non-insulin-dependent diabetes. <i>Kidney International</i> , 1993, 44, 139-146. | 5.2 | 58 |
| 189 | In vivo Glucose Metabolism in Obese and Type II Diabetic Subjects With or Without Hypertension. <i>Diabetes</i> , 1993, 42, 764-772. | 0.6 | 37 |
| 190 | Effects of physiological hyperinsulinemia on the intracellular metabolic partition of plasma glucose. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 1993, 265, E943-E953. | 3.5 | 11 |
| 191 | Characterization of cellular defects of insulin action in type 2 (non-insulin-dependent) diabetes mellitus.. <i>Journal of Clinical Investigation</i> , 1993, 91, 484-494. | 8.2 | 152 |
| 192 | In vivo glucose metabolism in obese and type II diabetic subjects with or without hypertension. <i>Diabetes</i> , 1993, 42, 764-772. | 0.6 | 8 |
| 193 | Total Body Fat Content and Fat Topography Are Associated Differently With In Vivo Glucose Metabolism in Nonobese and Obese Nondiabetic Women. <i>Diabetes</i> , 1992, 41, 1151-1159. | 0.6 | 145 |
| 194 | Lower limb arterio-venous shunts, autonomic neuropathy and diabetic foot. <i>Diabetes Research and Clinical Practice</i> , 1992, 16, 123-130. | 2.8 | 52 |
| 195 | Glucose and amino acid metabolism in chronic renal failure: effect of insulin and amino acids. <i>American Journal of Physiology - Renal Physiology</i> , 1992, 262, F168-F176. | 2.7 | 32 |
| 196 | Poor metabolic control and predisposition to hypertension, rather than hypertension itself, are risk factors for nephropathy in type 2 diabetes. <i>Acta Diabetologica</i> , 1992, 29, 123-129. | 2.5 | 7 |
| 197 | Insulin resistance, hypertension and cellular ion transport systems. <i>Acta Diabetologica</i> , 1992, 29, 196-200. | 2.5 | 5 |
| 198 | Close relationship between microalbuminuria and insulin resistance in essential hypertension and non-insulin dependent diabetes mellitus.. <i>Journal of the American Society of Nephrology: JASN</i> , 1992, 3, S56. | 6.1 | 47 |

| # | ARTICLE | IF | CITATIONS |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 199 | Total body fat content and fat topography are associated differently with in vivo glucose metabolism in nonobese and obese nondiabetic women. <i>Diabetes</i> , 1992, 41, 1151-1159. | 0.6 | 33 |
| 200 | Somatostatin response to a mixed meal in normals and in type I diabetics. <i>Peptides</i> , 1986, 7, 287-291. | 2.4 | 5 |
| 201 | Somatostatin plasma levels and biological effects following subcutaneous administration of somatostatin in man. <i>European Journal of Endocrinology</i> , 1986, 113, 465-470. | 3.7 | 1 |