
## **Charles J Glueck**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6191732/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Metabolic Syndrome in Childhood Predicts Adult Metabolic Syndrome and Type 2 Diabetes Mellitus 25 to 30 Years Later. Journal of Pediatrics, 2008, 152, 201-206.	1.8	532
2	Pregnancy outcomes among women with polycystic ovary syndrome treated with metformin. Human Reproduction, 2002, 17, 2858-2864.	0.9	338
3	Metformin therapy is associated with a decrease in plasma plasminogen activator inhibitor-1, lipoprotein(a), and immunoreactive insulin levels in patients with the polycystic ovary syndrome. Metabolism: Clinical and Experimental, 1997, 46, 454-457.	3.4	258
4	Characteristics of obesity in polycystic ovary syndrome: Etiology, treatment, and genetics. Metabolism: Clinical and Experimental, 2019, 92, 108-120.	3.4	215
5	Apolipoprotein A-I Gene Polymorphism Associated with Premature Coronary Artery Disease and Familial Hypoalphalipoproteinemia. New England Journal of Medicine, 1986, 314, 671-677.	27.0	213
6	Hypofibrinolysis, Thrombophilia, Osteonecrosis. Clinical Orthopaedics and Related Research, 2001, 386, 19-33.	1.5	194
7	Relationships of serum plant sterols (phytosterols) and cholesterol in 595 hypercholesterolemic subjects, and familial aggregation of phytosterols, cholesterol, and premature coronary heart disease in hyperphytosterolemic probands and their first-degree relatives. Metabolism: Clinical and Experimental. 1991. 40. 842-848.	3.4	184
8	Low serum 25 (OH) vitamin D levels (<32 ng/mL) are associated with reversible myositis-myalgia in statin-treated patients. Translational Research, 2009, 153, 11-16.	5.0	167
9	Enoxaparin Prevents Progression of Stages I and II Osteonecrosis of the Hip. Clinical Orthopaedics and Related Research, 2005, &NA, 164-170.	1.5	163
10	Neonatal familial type II hyperlipoproteinemia: Cord blood cholesterol in 1800 births. Metabolism: Clinical and Experimental, 1971, 20, 597-608.	3.4	154
11	Thrombophilia and Hypofibrinolysis; Pathophysiologies of Osteonecrosis. Clinical Orthopaedics and Related Research, 1997, 334, 43???56.	1.5	146
12	Evidence that homocysteine is an independent risk factor for atherosclerosis in hyperlipidemic patients. American Journal of Cardiology, 1995, 75, 132-136.	1.6	142
13	Lipids, lipoproteins, and sexual maturation during adolescence: The princeton maturation study. Metabolism: Clinical and Experimental, 1979, 28, 641-649.	3.4	131
14	Severe depression of high-density lipoprotein cholesterol levels in weight lifters and body builders by self-administered exogenous testosterone and anabolic-androgenic steroids. Metabolism: Clinical and Experimental, 1984, 33, 971-975.	3.4	125
15	The Relationship of Mutations in the MTHFR, Prothrombin, and PAI-1 Genes to Plasma Levels of Homocysteine, Prothrombin, and PAI-1 in Children and Adults. Thrombosis and Haemostasis, 1999, 81, 739-744.	3.4	125
16	Procoagulants and osteonecrosis. Journal of Rheumatology, 2003, 30, 783-91.	2.0	124
17	The Plasminogen Activator Inhibitor-1 Gene, Hypofibrinolysis, and Osteonecrosis. Clinical Orthopaedics and Related Research, 1999, 366, 133-146.	1.5	118
18	Blood Lead Levels and Dietary Calcium Intake in 1- to 11-Year-Old Children: The Second National Health and Nutrition Examination Survey, 1976 to 1980. Pediatrics, 1986, 78, 257-262.	2.1	118

#	Article	IF	CITATIONS
19	Plasma cholesterol and triglyceride levels in 6775 school children, ages 6–17. Metabolism: Clinical and Experimental, 1977, 26, 1199-1211.	3.4	117
20	Plasma Cholesterol and Triglyceride Distributions in 13,665 Children and Adolescents: the Prevalence Study of the Lipid Research Clinics Program. Pediatric Research, 1980, 14, 194-202.	2.3	115
21	Hypofibrinolysis: A common, major cause of osteonecrosis. American Journal of Hematology, 1994, 45, 156-166.	4.1	115
22	METABOLIC CONSEQUENCES OF EXPOSURE TO POLYCHLORINATED BIPHENYLS (PCB) IN SEWAGE SLUDGE. American Journal of Epidemiology, 1980, 112, 553-563.	3.4	113
23	Estrogen-induced pancreatitis in patients with previously covert familial Type V hyperlipoproteinemia. Metabolism: Clinical and Experimental, 1972, 21, 657-666.	3.4	110
24	FATNESS AND FAT PATTERNS: ASSOCIATIONS WITH PLASMA LIPIDS AND BLOOD PRESSURES IN ADULTS, 18 TO 57 YEARS OF AGE1. American Journal of Epidemiology, 1987, 126, 614-628.	3.4	105
25	RELATIONSHIPS OF MEASUREMENTS OF BODY MASS TO PLASMA LIPOPROTEINS IN SCHOOLCHILDREN AND ADULTS. American Journal of Epidemiology, 1980, 111, 395-406.	3.4	104
26	Idiopathic intracranial hypertension, polycystic-ovary syndrome, and thrombophilia. Translational Research, 2005, 145, 72-82.	2.3	104
27	Protein C and S Deficiency, Thrombophilia, and Hypofibrinolysis: Pathophysiologic Causes of Legg-Perthes Disease. Pediatric Research, 1994, 35, 383-388.	2.3	102
28	Pioglitazone and metformin in obese women with polycystic ovary syndrome not optimally responsive to metformin. Human Reproduction, 2003, 18, 1618-1625.	0.9	99
29	LEGG-CALVÉ-PERTHES DISEASE AND THROMBOPHILIA. Journal of Bone and Joint Surgery - Series A, 2004, 86, 2642-2647.	3.0	96
30	Heritable Thrombophilia and Hypofibrinolysis. JAMA Ophthalmology, 1999, 117, 43.	2.4	93
31	Risk of Venous Thromboembolism in Men Receiving Testosterone Therapy. Mayo Clinic Proceedings, 2015, 90, 1038-1045.	3.0	90
32	Effects of probucol on plasma cholesterol, high and low density lipoprotein cholesterol, and apolipoproteins A1 and A2 in adults with primary familial hypercholesterolemia. Metabolism: Clinical and Experimental, 1980, 29, 956-964.	3.4	89
33	Relationships between lipoprotein(a), lipids, apolipoproteins, basal and stimulated fibrinolytic regulators, and d-dimer. Metabolism: Clinical and Experimental, 1993, 42, 236-246.	3.4	88
34	Obesity and extreme obesity, manifest by ages 20–24 years, continuing through 32–41 years in women, should alert physicians to the diagnostic likelihood of polycystic ovary syndrome as a reversible underlying endocrinopathy. European Journal of Obstetrics, Gynecology and Reproductive Biology, 2005, 122, 206-212.	1.1	88
35	Hyperlipemia in Progeny of Parents. American Journal of Diseases of Children, 1974, 127, 70.	0.5	87
36	Idiopathic intracranial hypertension: associations with coagulation disorders and polycystic-ovary syndrome. Translational Research, 2003, 142, 35-45.	2.3	87

#	Article	IF	CITATIONS
37	Low or lowered cholesterol and risk of death from suicide and trauma. Metabolism: Clinical and Experimental, 1993, 42, 45-56.	3.4	86
38	CLUSTERING AND INTERRELATIONSHIPS OF CORONARY HEART DISEASE RISK FACTORS IN SCHOOLCHILDREN, AGES 6–19. American Journal of Epidemiology, 1980, 112, 524-538.	3.4	84
39	Childhood risk factors predict cardiovascular disease, impaired fasting glucose plus type 2 diabetes mellitus, and high blood pressure 26 years later at a mean age of 38 years: the Princeton–lipid research clinics follow-up study. Metabolism: Clinical and Experimental, 2012, 61, 531-541.	3.4	84
40	Statin intolerance because of myalgia, myositis, myopathy, or myonecrosis can in most cases be safely resolved by vitamin d supplementation. North American Journal of Medical Sciences, 2015, 7, 86.	1.7	83
41	Idiopathic osteonecrosis, hypofibrinolysis, high plasminogen activator inhibitor, high lipoprotein(a), and therapy with stanozolol. American Journal of Hematology, 1995, 48, 213-220.	4.1	81
42	Heritable Thrombophilia-Hypofibrinolysis and Osteonecrosis of the Femoral Head. Clinical Orthopaedics and Related Research, 2008, 466, 1034-1040.	1.5	81
43	Resistance to Activated Protein C and Legg-Perthes Disease. Clinical Orthopaedics and Related Research, 1997, 338, 139-152.	1.5	80
44	Prevention of gestational diabetes by metformin plus diet in patients with polycystic ovary syndrome. Fertility and Sterility, 2008, 89, 625-634.	1.0	79
45	Pre-teen insulin resistance predicts weight gain, impaired fasting glucose, and type 2 diabetes at age 18–19 y: a 10-y prospective study of black and white girls. American Journal of Clinical Nutrition, 2008, 88, 778-788.	4.7	79
46	Vitamin D deficiency, myositis–myalgia, and reversible statin intolerance. Current Medical Research and Opinion, 2011, 27, 1683-1690.	1.9	77
47	Childhood Predictors of Adult Type 2 Diabetes at 9- and 26-Year Follow-ups. JAMA Pediatrics, 2010, 164, 53-60.	3.0	74
48	Determinants of high-density lipoprotein cholesterol in blacks and whites: The second National Health and Nutrition Examination Survey. American Heart Journal, 1984, 108, 641-653.	2.7	73
49	Hypocholesterolemia and Affective Disorders. American Journal of the Medical Sciences, 1994, 308, 218-225.	1.1	71
50	Genetic hypofibrinolysis in complicated pregnancies. Obstetrics and Gynecology, 2001, 97, 44-48.	2.4	70
51	Benign symmetric lipomatosis (launois-bensaude adenolipomatosis) with gout and hyperlipoproteinemia. American Journal of Medicine, 1970, 48, 239-246.	1.5	69
52	The pathophysiology of alveolar osteonecrosis of the jaw: Anticardiolipin antibodies, thrombophilia, and hypofibrinolysis. Translational Research, 1996, 127, 481-488.	2.3	66
53	Thrombophilia, Hypofibrinolysis, the eNOS T-786C Polymorphism, and Multifocal Osteonecrosis. Journal of Bone and Joint Surgery - Series A, 2008, 90, 2220-2229.	3.0	64
54	Risk factors for cardiovascular disease and type 2 diabetes retained from childhood to adulthood predict adult outcomes: the Princeton LRC Follow-up Study. International Journal of Pediatric Endocrinology (Springer), 2012, 2012, 6.	1.6	64

#	Article	IF	CITATIONS
55	Pregnancy Loss, Polycystic Ovary Syndrome, Thrombophilia, Hypofibrinolysis, Enoxaparin, Metformin. Clinical and Applied Thrombosis/Hemostasis, 2004, 10, 323-334.	1.7	62
56	Rosuvastatin 5 and 10 mg/d: A pilot study of the effects in hypercholesterolemic adults unable to tolerate other statins and reach Idl cholesterol goals with nonstatin lipid-lowering therapies. Clinical Therapeutics, 2006, 28, 933-942.	2.5	62
57	Nonarteritic anterior ischemic optic neuropathy: associations with homozygosity for the C677T methylenetetrahydrofolate reductase mutation. Translational Research, 2004, 143, 184-192.	2.3	61
58	Testosterone therapy, thrombosis, thrombophilia, cardiovascular events. Metabolism: Clinical and Experimental, 2014, 63, 989-994.	3.4	60
59	Plasminogen activator inhibitor activity, 4G5G polymorphism of the plasminogen activator inhibitor 1 gene, and first-trimester miscarriage in women with polycystic ovary syndrome. Metabolism: Clinical and Experimental, 2006, 55, 345-352.	3.4	59
60	CORRELATION OF THROMBOPHILIA AND HYPOFIBRINOLYSIS WITH PULMONARY EMBOLISM FOLLOWING TOTAL HIP ARTHROPLASTY. Journal of Bone and Joint Surgery - Series A, 2002, 84, 2161-2167.	3.0	57
61	Obesity, free testosterone, and cardiovascular risk factors in adolescents with polycystic ovary syndrome and regularly cycling adolescents. Metabolism: Clinical and Experimental, 2006, 55, 508-514.	3.4	56
62	Prospective Assessment of Coronary Heart Disease Risk Factors: The NHANES I Epidemiologic Follow-up Study (NHEFS) 16-Year Follow-up. Journal of the American College of Nutrition, 1998, 17, 263-269.	1.8	55
63	Thrombophilia, hypofibrinolysis, and alveolar osteonecrosis of the jaws. Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics, 1996, 81, 557-566.	1.4	54
64	Pediatric triglycerides predict cardiovascular disease events in the fourth to fifth decade of life. Metabolism: Clinical and Experimental, 2009, 58, 1277-1284.	3.4	52
65	Polycystic ovary syndrome, the G1691A factor V Leiden mutation, and plasminogen activator inhibitor activity: associations with recurrent pregnancy loss. Metabolism: Clinical and Experimental, 2003, 52, 1627-1632.	3.4	50
66	An observational study of severe hypertriglyceridemia, hypertriglyceridemic acute pancreatitis, and failure of triglyceride-lowering therapy when estrogens are given to women with and without familial hypertriglyceridemia. Clinica Chimica Acta, 2003, 332, 11-19.	1.1	50
67	Familial Heterozygous Protein-S Deficiency in a Patient Who Had Multifocal Osteonecrosis. A Case Report*. Journal of Bone and Joint Surgery - Series A, 1997, 79, 1079-84.	3.0	50
68	Heterozygosity for the leiden mutation of the factor v gene, a common pathoetiology for osteonecrosis of the jaw, with thrombophilia augmented by exogenous estrogens. Translational Research, 1997, 130, 540-543.	2.3	49
69	Thrombotic events after starting exogenous testosterone in men with previously undiagnosed familial thrombophilia. Translational Research, 2011, 158, 225-234.	5.0	49
70	Pediatric familial type II hyperlipoproteinemia: effects of diet on plasma cholesterol in the first year of life. American Journal of Clinical Nutrition, 1972, 25, 224-230.	4.7	48
71	Legg-Perthes disease in three siblings, two heterozygous and one homozygous for the factor V Leiden mutation. Journal of Pediatrics, 1998, 132, 885-888.	1.8	48
72	Treatment of polycystic ovary syndrome with insulin-lowering agents. Expert Opinion on Pharmacotherapy, 2002, 3, 1177-1189.	1.8	48

#	Article	IF	CITATIONS
73	Ocular Vascular Thrombotic Events: Central Retinal Vein and Central Retinal Artery Occlusions. Clinical and Applied Thrombosis/Hemostasis, 2008, 14, 286-294.	1.7	48
74	Early and late menarche are associated with oligomenorrhea and predict metabolic syndrome 26years later. Metabolism: Clinical and Experimental, 2013, 62, 1597-1606.	3.4	48
75	Genetic Hypofibrinolysis in Complicated Pregnancies. Obstetrics and Gynecology, 2001, 97, 44-48.	2.4	47
76	Growth, motor, and social development in breast- and formula-fed infants of metformin-treated women with polycystic ovary syndrome. Journal of Pediatrics, 2006, 148, 628-632.e2.	1.8	47
77	Sex Hormone-Binding Globulin, Oligomenorrhea, Polycystic Ovary Syndrome, and Childhood Insulin at Age 14 Years Predict Metabolic Syndrome and Class III Obesity at Age 24 Years. Journal of Pediatrics, 2011, 159, 308-313.e2.	1.8	47
78	THE JOHN CHARNLEY AWARD: Heritable Thrombophilia and Development of Thromboembolic Disease after Total Hip Arthroplasty. Clinical Orthopaedics and Related Research, 2005, 441, 40-55.	1.5	45
79	Associations of Thrombophilia, Hypofibrinolysis, and Retinal Vein Occlusion. Clinical and Applied Thrombosis/Hemostasis, 2005, 11, 375-389.	1.7	44
80	Coronary heart disease risk factors in adult premenopausal white women with polycystic ovary syndrome compared with a healthy female population. Metabolism: Clinical and Experimental, 2009, 58, 714-721.	3.4	39
81	Thrombophilia and retinal vascular occlusion. Clinical Ophthalmology, 2012, 6, 1377.	1.8	38
82	Metformin and gestational diabetes. Current Diabetes Reports, 2003, 3, 303-312.	4.2	37
83	The contentious nature of gestational diabetes: diet, insulin, glyburide and metformin. Expert Opinion on Pharmacotherapy, 2002, 3, 1557-1568.	1.8	36
84	Risk Factors for Pulmonary Emboli after Total Hip or Knee Arthroplasty. Clinical Orthopaedics and Related Research, 2004, 422, 154-163.	1.5	36
85	Sustainability of 8% weight loss, reduction of insulin resistance, and amelioration of atherogenic-metabolic risk factors over 4 years by metformin-diet in women with polycystic ovary syndrome. Metabolism: Clinical and Experimental, 2006, 55, 1582-1589.	3.4	36
86	Association Between the T-786C eNOS Polymorphism and Idiopathic Osteonecrosis of the Head of the Femur. Journal of Bone and Joint Surgery - Series A, 2007, 89, 2460-2468.	3.0	35
87	Testosterone Therapy, Thrombophilia–Hypofibrinolysis, and Hospitalization for Deep Venous Thrombosis-Pulmonary Embolus. Clinical and Applied Thrombosis/Hemostasis, 2014, 20, 244-249.	1.7	35
88	Metformin before and during pregnancy and lactation in polycystic ovary syndrome. Expert Opinion on Drug Safety, 2007, 6, 191-198.	2.4	34
89	Effects of metformin-diet intervention before and throughout pregnancy on obstetric and neonatal outcomes in patients with polycystic ovary syndrome. Current Medical Research and Opinion, 2013, 29, 55-62.	1.9	34
90	Thrombophilia in 67 Patients With Thrombotic Events After Starting Testosterone Therapy. Clinical and Applied Thrombosis/Hemostasis, 2016, 22, 548-553.	1.7	34

#	Article	IF	CITATIONS
91	Testosterone, thrombophilia, thrombosis. Translational Research, 2015, 165, 537-548.	5.0	33
92	Interaction of Duloxetine and Warfarin Causing Severe Elevation of International Normalized Ratio. JAMA - Journal of the American Medical Association, 2006, 295, 1513.	7.4	32
93	Metformin-diet benefits in women with polycystic ovary syndrome in the bottom and top quintiles for insulin resistance. Metabolism: Clinical and Experimental, 2005, 54, 113-121.	3.4	31
94	Symptomatic myositis-myalgia in hypercholesterolemic statin-treated patients with concurrent vitamin D deficiency leading to statin intolerance may reflect a reversible interaction between vitamin D deficiency and statins on skeletal muscle. Medical Hypotheses, 2011, 77, 658-661.	1.5	31
95	Hyperinsulinemia and metabolic syndrome at mean age of 10 years in black and white schoolgirls and development of impaired fasting glucose and type 2 diabetes mellitus by mean age of 24 years. Metabolism: Clinical and Experimental, 2011, 60, 24-31.	3.4	31
96	The Role of the Factor V Leiden Mutation in Osteonecrosis of the Hip. Clinical and Applied Thrombosis/Hemostasis, 2013, 19, 499-503.	1.7	31
97	Evidence that anticardiolipin antibodies are independent risk factors for atherosclerotic vascular disease. American Journal of Cardiology, 1999, 83, 1490-1494.	1.6	29
98	Anticoagulant therapy for osteonecrosis associated with heritable hypofibrinolysis and thrombophilia. Expert Opinion on Investigational Drugs, 2001, 10, 1309-1316.	4.1	28
99	An observational study of reduction of insulin resistance and prevention of development of type 2 diabetes mellitus in women with polycystic ovary syndrome treated with metformin and diet. Metabolism: Clinical and Experimental, 2008, 57, 954-960.	3.4	28
100	Factor V Leiden mutation: a treatable etiology for sporadic and recurrent pregnancy loss. Fertility and Sterility, 2008, 89, 410-416.	1.0	28
101	Osteonecrosis: a multifactorial etiology. Journal of Oral and Maxillofacial Surgery, 2004, 62, 904-905.	1.2	27
102	Population-specific alleles: the polymorphism (k121q) of the human glycoprotein PC-1 gene is strongly associated with race but not with insulin resistance in black and white children. Metabolism: Clinical and Experimental, 2004, 53, 465-468.	3.4	27
103	Testosterone, Thrombophilia, and Thrombosis. Clinical and Applied Thrombosis/Hemostasis, 2014, 20, 22-30.	1.7	27
104	Adolescent Oligomenorrhea (Age 14–19) Tracks Into the Third Decade of Life (Age 20–28) and Predicts Increased Cardiovascular Risk Factors and Metabolic Syndrome. Metabolism: Clinical and Experimental, 2015, 64, 539-553.	3.4	27
105	Testosterone Therapy, Thrombophilia, Venous Thromboembolism, and Thrombotic Events. Journal of Clinical Medicine, 2019, 8, 11.	2.4	27
106	The factor V Leiden mutation, high factor VIII, and high plasminogen activator inhibitor activity: etiologies for sporadic miscarriage. Metabolism: Clinical and Experimental, 2005, 54, 1345-1349.	3.4	25
107	Stromelysin-1 5A/6A and eNOS T-786C Polymorphisms, MTHFR C677T and A1298C Mutations, and Cigarette-Cannabis Smoking: A Pilot, Hypothesis-Generating Study of Gene-Environment Pathophysiological Associations With Buerger's Disease. Clinical and Applied Thrombosis/Hemostasis, 2006, 12, 427-439.	1.7	24
108	Changes in weight, papilledema, headache, visual field, and life status in response to diet and metformin in women with idiopathic intracranial hypertension with and without concurrent polycystic ovary syndrome or hyperinsulinemia. Translational Research, 2006, 148, 215-222.	5.0	23

#	Article	IF	CITATIONS
109	Long Term Anticoagulation (4–16 Years) Stops Progression of Idiopathic Hip Osteonecrosis Associated with Familial Thrombophilia. Advances in Orthopedics, 2015, 2015, 1-7.	1.0	23
110	Treatment of severe familial hypertriglyceridemia during pregnancy with very-low-fat diet and n-3 fatty acids. Nutrition, 1996, 12, 202-205.	2.4	22
111	Dietary trans fatty acid intake is associated with increased fetal loss. Fertility and Sterility, 2008, 90, 385-390.	1.0	22
112	Safety of 50,000-100,000 units of vitamin D3/week in vitamin D-deficient, hypercholesterolemic patients with reversible statin intolerance. North American Journal of Medical Sciences, 2016, 8, 156.	1.7	22
113	Interaction of Heritable and Estrogen-induced Thrombophilia: Possible Etiologies for Ischemic Optic Neuropathy and Ischemic Stroke. Thrombosis and Haemostasis, 2001, 85, 256-259.	3.4	21
114	Staging Bisphosphonate-Related Osteonecrosis of the Jaw Should Include Early Stages of Disease. Journal of Oral and Maxillofacial Surgery, 2007, 65, 1899-1900.	1.2	21
115	Should high creatine kinase discourage the initiation or continuance of statins for the treatment of hypercholesterolemia?. Metabolism: Clinical and Experimental, 2009, 58, 233-238.	3.4	21
116	Thrombosis in Three Postmenopausal Women Receiving Testosterone Therapy for Low Libido. Women's Health, 2013, 9, 405-410.	1.5	21
117	Testosterone, thrombophilia, thrombosis. Blood Coagulation and Fibrinolysis, 2014, 25, 683-687.	1.0	21
118	Association Between the T-786C eNOS Polymorphism and Idiopathic Osteonecrosis of the Head of the Femur. Journal of Bone and Joint Surgery - Series A, 2007, 89, 2460-2468.	3.0	21
119	Selected Nutrient Intakes of Free-Living White Children Ages 6–19 Years. The Lipid Research Clinics Program Prevalence Study(40, 41). Pediatric Research, 1983, 17, 124-130.	2.3	20
120	Contribution of fasting hyperinsulinemia to prediction of atherosclerotic cardiovascular disease status in 293 hyperlipidemic patients. Metabolism: Clinical and Experimental, 1999, 48, 1437-1444.	3.4	20
121	Diagnostic ramifications of ocular vascular occlusion as a first thrombotic event associated with factor V Leiden and prothrombin gene heterozygosity. Clinical Ophthalmology, 2015, 9, 591.	1.8	20
122	High and low density lipoprotein cholesterol levels in hypercholesterolemic school children. Lipids, 1979, 14, 99-104.	1.7	19
123	Enoxaparin-metformin and enoxaparin alone may safely reduce pregnancy loss. Translational Research, 2009, 153, 33-43.	5.0	19
124	Retinal artery and vein thrombotic occlusion during pregnancy: markers for familial thrombophilia and adverse pregnancy outcomes. Clinical Ophthalmology, 2016, 10, 935.	1.8	19
125	Thrombophilia in Klinefelter Syndrome With Deep Venous Thrombosis, Pulmonary Embolism, and Mesenteric Artery Thrombosis on Testosterone Therapy: A Pilot Study. Clinical and Applied Thrombosis/Hemostasis, 2017, 23, 973-979.	1.7	19
126	Endothelial Nitric Oxide Synthase T-786C Mutation, A Reversible Etiology of Prinzmetal's Angina Pectoris. American Journal of Cardiology, 2010, 105, 792-796.	1.6	18

#	Article	IF	CITATIONS
127	Plasma 25-Hydroxy-vitamin D in Familial Hypercholesterolemic Children Receiving Colestipol Resin. Pediatric Research, 1978, 12, 980-982.	2.3	17
128	Idiopathic Intracranial Hypertension: Associations with Thrombophilia and Hypofibrinolysis in Men. Clinical and Applied Thrombosis/Hemostasis, 2005, 11, 441-448.	1.7	17
129	Paradoxically High Adiponectin in Obese 16-Year-Old Girls Protects Against Appearance of the Metabolic Syndrome and its Components Seven Years Later. Journal of Pediatrics, 2011, 158, 208-214.e1.	1.8	17
130	Effects of diet and high density lipoprotein subfractions on the removal of cellular cholesterol. Lipids, 1980, 15, 230-235.	1.7	16
131	Homeostasis model assessment of insulin resistance*body mass index interactions at ages 9 to 10 years predict metabolic syndrome risk factor aggregate score at ages 18 to 19 years: a 10-year prospective study of black and white girls. Metabolism: Clinical and Experimental, 2009, 58, 290-295.	3.4	16
132	Ocular Vascular Thrombotic Events: A Diagnostic Window to Familial Thrombophilia (Compound) Tj ETQq0 0 0 rg Thrombosis/Hemostasis, 2009, 15, 12-18.	gBT /Overl 1.7	ock 10 Tf 50 5 16
133	Preteen insulin levels interact with caloric intake to predict increases in obesity at ages 18 to 19 years: a 10-year prospective study of black and white girls. Metabolism: Clinical and Experimental, 2010, 59, 718-727.	3.4	16
134	Estrogen replacement therapy, thrombophilia, and atherothrombosis. Metabolism: Clinical and Experimental, 2002, 51, 724-732.	3.4	15
135	The endothelial nitric oxide synthase T-786c mutation, a treatable etiology of Prinzmetal's angina. Translational Research, 2013, 162, 64-66.	5.0	15
136	Medical Treatment of Osteonecrosis of the Knee Associated With Thrombophilia-Hypofibrinolysis. Orthopedics, 2014, 37, e911-6.	1.1	15
137	Eligibility for PCSK9 treatment in 734 Hypercholesterolemic patients referred to a regional cholesterol treatment center with LDL cholesterol ≥70Âmg/dl despite maximal tolerated cholesterol lowering therapy. Lipids in Health and Disease, 2016, 15, 55.	3.0	15
138	Low serum vitamin D, statin associated muscle symptoms, vitamin D supplementation. Atherosclerosis, 2017, 256, 125-127.	0.8	15
139	Thrombophilia–hypofibrinolysis and atherothrombotic cardiovascular disease ≤ge 45 years. Translational Research, 2007, 150, 93-100.	5.0	14
140	Adolescent oligomenorrhea in a biracial schoolgirl cohort: a simple clinical parameter predicting impaired fasting glucose plus type 2 diabetes mellitus, insulin, glucose, insulin resistance, and centripetal obesity from age 19 to 25 years. Metabolism: Clinical and Experimental, 2011, 60, 1285-1293.	3.4	14
141	Titrating lovaza from 4 to 8 to 12 grams/day in patients with primary hypertriglyceridemia who had triglyceride levels >500 mg/dl despite conventional triglyceride lowering therapy. Lipids in Health and Disease, 2012, 11, 143.	3.0	14
142	Testosterone, anastrozole, factor V Leiden heterozygosity and osteonecrosis of the jaws. Blood Coagulation and Fibrinolysis, 2014, 25, 286-288.	1.0	14
143	Pharmacoeconomics of PCSK9 inhibitors in 103 hypercholesterolemic patients referred for diagnosis and treatment to a cholesterol treatment center. Lipids in Health and Disease, 2016, 15, 132.	3.0	14
144	Exogenous Estrogen May Exacerbate Thrombophilia, Impair Bone Healing and Contribute to Development of Chronic Facial Pain. Cranio - Journal of Craniomandibular Practice, 1998, 16, 143-153.	1.4	13

9

#	Article	IF	CITATIONS
145	Intrafamilial associations of cholesterol and triglyceride among related and unrelated household members. Clinical Genetics, 1980, 18, 321-328.	2.0	13
146	Insulin Resistance, Obesity, Hypofibrinolysis, Hyperandrogenism, and Coronary Heart Disease Risk Factors in 25 Pre-Perimenarchal Girls Age <14 Years, 13 with Precocious Puberty, 23 with a First-degree Relative with Polycystic Ovary Syndrome. Journal of Pediatric Endocrinology and Metabolism, 2008, 21, 973-84.	0.9	12
147	Is pharmacogenomics our future? Metformin, ovulation and polymorphism of theSTK11gene in polycystic ovary syndrome. Pharmacogenomics, 2008, 9, 1163-1165.	1.3	12
148	Alirocumab in high-risk patients: Observations from the open-label expanded use program. Journal of Clinical Lipidology, 2018, 12, 662-668.	1.5	12
149	PARENT-CHILD CORONARY HEART DISEASE RISK FACTOR ASSOCIATIONS. American Journal of Epidemiology, 1981, 114, 827-835.	3.4	11
150	Legg-Calve-Perthes Disease, Venous and Arterial Thrombi, and the Factor V Leiden Mutation in a Four-Generation Kindred. Journal of Pediatric Orthopaedics, 2007, 27, 834-837.	1.2	11
151	Case report: primary osteonecrosis associated with thrombophilia-hypofibrinolysis and worsened by testosterone therapy. BMC Hematology, 2017, 17, 5.	2.6	11
152	Determinants of Persistent Obesity and Hyperinsulinemia in a Biracial Cohort: A 15-Year Prospective Study of Schoolgirls. Journal of Pediatrics, 2010, 157, 559-565.	1.8	10
153	Paradoxically high adiponectin and the healthy obese phenotype in obese black and white 16-year-old girls. Translational Research, 2010, 156, 302-308.	5.0	10
154	Determinants of ApoB, ApoA1, and the ApoB/ApoA1 ratio in healthy schoolgirls, prospectively studied from mean ages 10 to 19 years: the Cincinnati National Growth and Health Study. Metabolism: Clinical and Experimental, 2012, 61, 1377-1387.	3.4	10
155	Hospitalization for pulmonary embolism associated with antecedent testosterone or estrogen therapy in patients found to have familial and acquired thrombophilia. BMC Hematology, 2016, 16, 6.	2.6	10
156	Testosterone Therapy Can Interact With Thrombophilia, Leading to Osteonecrosis. Orthopedics, 2015, 38, e1073-8.	1.1	10
157	LIPID AND LIPOPROTEIN TRACKING IN CHILDREN. Pediatric Research, 1979, 13, 1082-1084.	2.3	9
158	Detecting Thrombophilia, Hypofibrinolysis and Reduced Nitric Oxide Production in Osteonecrosis. Seminars in Arthroplasty, 2007, 18, 184-191.	0.7	9
159	Tâ~786C polymorphism of the endothelial nitric oxide synthase gene and neuralgia-inducing cavitational osteonecrosis of the jaws. Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics, 2010, 109, 548-553.	1.4	9
160	Efficacy, safety, Low density lipoprotein cholesterol lowering, and calculated 10-year cardiovascular risk reduction of alirocumab and evolocumab in addition to maximal tolerated cholesterol lowering therapy: a post-commercialization study. Lipids in Health and Disease, 2017, 16, 19.	3.0	9
161	Thrombophilia in 153 Patients With Premature Cardiovascular Disease â‰ <b>¤</b> ge 45. Clinical and Applied Thrombosis/Hemostasis, 2018, 24, 295-302.	1.7	9
162	Long-term Anticoagulation Prevents Progression of Stages I and II Primary Osteonecrosis of the Hip in Patients With Familial Thrombophilia. Orthopedics, 2020, 43, e208-e214.	1.1	9

#	Article	IF	CITATIONS
163	Severe vitamin D deficiency, myopathy, and rhabdomyolysis. North American Journal of Medical Sciences, 2013, 5, 494.	1.7	9
164	VKORC1 Variant Genotypes Influence Warfarin Response in Patients Undergoing Total Joint Arthroplasty: A Pilot Study. Clinical Orthopaedics and Related Research, 2009, 467, 1773-1780.	1.5	8
165	Ramifications of adolescent menstrual cycles ≥42 days in young adults. Fertility and Sterility, 2011, 96, 236-240.e1.	1.0	8
166	The Child as Proband for Future Parental Cardiometabolic Disease: The 26-Year Prospective Princeton Lipid Research Clinics Follow-Up Study. Journal of Pediatrics, 2012, 160, 590-597.e3.	1.8	8
167	Adolescent and Young Adult Female Determinants of Visceral Adipose Tissue at Ages 26-28 Years. Journal of Pediatrics, 2015, 166, 936-946.e3.	1.8	8
168	Testosterone therapy, thrombophilia, and hospitalization for deep venous thrombosis–pulmonary embolus, an exploratory, hypothesis-generating study. Medical Hypotheses, 2015, 84, 341-343.	1.5	8
169	Familial Hypertriglyceridemia in Children: Dietary Management. Pediatric Research, 1977, 11, 953-957.	2.3	7
170	Low and High Density Lipoprotein Cholesterol Interrelationships in Neonates with Low Density Lipoprotein Cholesterol ≤he 10th Percentile and in Neonates with High Density Lipoprotein Cholesterol ≥ the 90th Percentile. Pediatric Research, 1977, 11, 957-959.	2.3	7
171	High factor XI, recurrent pregnancy loss, enoxaparin. Fertility and Sterility, 2010, 94, 2828-2831.	1.0	7
172	Treatable high homocysteine alone or in concert with five other thrombophilias in 1014 patients with thrombotic events. Blood Coagulation and Fibrinolysis, 2015, 26, 736-742.	1.0	7
173	Beyond bisphosphonates: Thrombophilia, hypofibrinolysis, and jaw osteonecrosis. Journal of Oral and Maxillofacial Surgery, 2006, 64, 1704-1705.	1.2	6
174	Amaurosis Fugax Caused by Heritable Thrombophilia-Hypofibrinolysis in Cases Without Carotid Atherosclerosis. Clinical and Applied Thrombosis/Hemostasis, 2007, 13, 124-129.	1.7	6
175	Bone Marrow Edema: Mild or Nascent Variant of Ischemic Bone Disease. Journal of Oral and Maxillofacial Surgery, 2008, 66, 205-208.	1.2	6
176	Metformin-Diet Ameliorates Coronary Heart Disease Risk Factors and Facilitates Resumption of Regular Menses in Adolescents with Polycystic Ovary Syndrome. Journal of Pediatric Endocrinology and Metabolism, 2009, 22, 815-26.	0.9	6
177	Retinal vascular occlusion: a window to diagnosis of familial and acquired thrombophilia and hypofibrinolysis, with important ramifications for pregnancy outcomes. Clinical Ophthalmology, 2016, Volume 10, 1479-1486.	1.8	6
178	Thromboembolism Peaking 3 Months after Starting Testosterone Therapy: Testosterone–Thrombophilia Interactions. Journal of Investigative Medicine, 2018, 66, 733-738.	1.6	6
179	A Likelihood Model That Accounts for Censoring Due to Fetal Loss Can Accurately Test the Effects of Maternal and Fetal Genotype on the Probability of Miscarriage. Human Heredity, 2009, 67, 57-65.	0.8	5
180	Pediatric Non–High-Density Lipoprotein Cholesterol, the Metabolic Syndrome, and Insulin Resistance. Journal of Pediatrics, 2011, 158, 179-181.	1.8	5

#	Article	IF	CITATIONS
181	Four Thrombotic Events Over 5 Years, Two Pulmonary Emboli and Two Deep Venous Thrombosis, When Testosterone-HCG Therapy Was Continued Despite Concurrent Anticoagulation in a 55-Year-Old Man With Lupus Anticoagulant. Journal of Investigative Medicine High Impact Case Reports, 2016, 4, 232470961666183.	0.6	5
182	Treatment of Osteonecrosis of the Hip and Knee with Enoxaparin. , 2014, , 241-247.		5
183	Venous Thromboembolism After Knee Arthroscopy in Undiagnosed Familial Thrombophilia. Orthopedics, 2016, 39, e1052-e1057.	1.1	5
184	A multivariate analysis of familial associations of lipoprotein levels in the Lipid Research Clinics Collaborative Family Study: I. Familial correlation and regression analyses. Genetic Epidemiology, 1985, 2, 283-300.	1.3	4
185	Ramifications of Four Concurrent Thrombophilic Mutations and One Hypofibrinolytic Mutation. Clinical and Applied Thrombosis/Hemostasis, 2004, 10, 365-371.	1.7	4
186	Race, Childhood Insulin, Childhood Caloric Intake, and Class 3 Obesity at Age 24: 14‥ear Prospective Study of Schoolgirls. Obesity, 2012, 20, 597-604.	3.0	4
187	Efficacy and safety of proprotein convertase subtilisin-kexin type 9 (PCSK9) inhibitors, alirocumab and evolocumab, a post-commercialization study. Lipids in Health and Disease, 2017, 16, 141.	3.0	4
188	High-Sensitivity C Reactive Protein: Associations with Cardiovascular Risk Factors and Tracking in Female Adolescents and Young Adults. ISRN Pediatrics, 2011, 2011, 1-9.	1.2	3
189	T786C Mutation in the Endothelial Nitric Oxide Synthase Gene in Patients With Primary Osteonecrosis. Orthopedics, 2017, 40, e898-e903.	1.1	3
190	Age of Menarche and Oligomenorrhea in Adolescence: Cardiometabolic Ramifications in Adulthood. Journal of Pediatrics, 2015, 167, 225-227.	1.8	2
191	Eligibility for alirocumab or evolocumab treatment in 1090 hypercholesterolemic patients referred to a regional cholesterol treatment center with LDL cholesterol ≥70 mg/dL despite maximal-tolerated LDL-cholesterol-lowering therapy. Vascular Health and Risk Management, 2017, Volume 13, 247-253.	2.3	2
192	Statin Intolerance and Vitamin D Supplementation. North American Journal of Medical Sciences, 2015, 7, 339-40.	1.7	2
193	Successful anticoagulation for bilateral central retinal vein occlusion. Clinica Chimica Acta, 2011, 412, 1165-1166.	1.1	1
194	The eNOS T786C mutation, Prinzmetal's Variant Angina, and amelioration of angina by l-arginine in 59 patients with intractable angina despite calcium channel blocker–nitrate therapy. IJC Metabolic & Endocrine, 2015, 8, 13-19.	0.5	1
195	Osteonecrosis and Thrombophilia: Pathophysiology, Diagnosis, and Treatment. , 2017, , 3-17.		1
196	Does low serum 25 OH vitamin D interact with very strenuous physical activity, facilitating development of rhabdomyolysis?. Medical Hypotheses, 2013, 81, 551-552.	1.5	0
197	Progressively Worsening Premature Coronary Artery Disease: Adding Anticoagulation Stabilizes–Reverses Clinical Symptomatic Disease Progression in Thrombophilic–Atherothrombotic Patients: A Pilot Study. Clinical and Applied Thrombosis/Hemostasis, 2017, 23, 1060-1070.	1.7	0

198 The Long-Term Metabolic Complications of Childhood Obesity. , 2010, , 253-263.

#	Article	IF	CITATIONS
199	Response to diet and metformin in women with idiopathic intracranial hypertension with and without concurrent polycystic ovary syndrome or hyperinsulinemia. MedGenMed: Medscape General Medicine, 2005, 7, 41.	0.2	0