Charlotte Hanlon

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6193841/publications.pdf

Version: 2024-02-01

237 papers

9,403 citations

44069 48 h-index 60623 81 g-index

253 all docs

253 docs citations

times ranked

253

7252 citing authors

#	Article	IF	CITATIONS
1	COVID-19 mental health impact and responses in low-income and middle-income countries: reimagining global mental health. Lancet Psychiatry,the, 2021, 8, 535-550.	7.4	455
2	PRIME: A Programme to Reduce the Treatment Gap for Mental Disorders in Five Low- and Middle-Income Countries. PLoS Medicine, 2012, 9, e1001359.	8.4	436
3	Acceptability and feasibility of using non-specialist health workers to deliver mental health care: Stakeholder perceptions from the PRIME district sites in Ethiopia, India, Nepal, South Africa, and Uganda. Social Science and Medicine, 2014, 118, 33-42.	3.8	259
4	Challenges and Opportunities for Implementing Integrated Mental Health Care: A District Level Situation Analysis from Five Low- and Middle-Income Countries. PLoS ONE, 2014, 9, e88437.	2.5	255
5	WPA guidance on steps, obstacles and mistakes to avoid in the implementation of community mental health care. World Psychiatry, 2010, 9, 67-77.	10.4	224
6	Detecting perinatal common mental disorders in Ethiopia: Validation of the self-reporting questionnaire and Edinburgh Postnatal Depression Scale. Journal of Affective Disorders, 2008, 108, 251-262.	4.1	222
7	Strengthening mental health systems in low- and middle-income countries: the Emerald programme. BMC Medicine, 2015, 13, 79.	5.5	190
8	Service user and caregiver involvement in mental health system strengthening in low- and middle-income countries: systematic review. BMC Health Services Research, 2016, 16, 79.	2.2	175
9	Detecting postnatal common mental disorders in Addis Ababa, Ethiopia: Validation of the Edinburgh Postnatal Depression Scale and Kessler Scales. Journal of Affective Disorders, 2010, 122, 102-108.	4.1	165
10	Efficacy of psychosocial interventions for mental health outcomes in low-income and middle-income countries: an umbrella review. Lancet Psychiatry,the, 2020, 7, 162-172.	7.4	163
11	Impact of antenatal common mental disorders upon perinatal outcomes in Ethiopia: the Pâ€MaMiE populationâ€based cohort study. Tropical Medicine and International Health, 2009, 14, 156-166.	2.3	132
12	Validity of brief screening questionnaires to detect depression in primary care in Ethiopia. Journal of Affective Disorders, 2015, 186, 32-39.	4.1	131
13	Excess mortality in severe mental illness: 10-year population-based cohort study in rural Ethiopia. British Journal of Psychiatry, 2015, 206, 289-296.	2.8	130
14	Strengthening mental health system governance in six low- and middle-income countries in Africa and South Asia: challenges, needs and potential strategies. Health Policy and Planning, 2017, 32, 699-709.	2.7	127
15	Stigma, explanatory models and unmet needs of caregivers of children with developmental disorders in a low-income African country: a cross-sectional facility-based survey. BMC Health Services Research, 2016, 16, 152.	2.2	119
16	Maternal mental health in primary care in five low- and middle-income countries: a situational analysis. BMC Health Services Research, 2016, 16, 53.	2.2	110
17	Mental health care in Nepal: current situation and challenges for development of a district mental health care plan. Conflict and Health, 2015, 9, 3.	2.7	108
18	Arginine residues at codons 112 and 158 in the apolipoprotein E gene correspond to the ancestral state in humans. Atherosclerosis, 1995, 112, 85-90.	0.8	107

#	Article	IF	CITATIONS
19	Prevalence and predictors of undernutrition among infants aged six and twelve months in Butajira, Ethiopia: The P-MaMiE Birth Cohort. BMC Public Health, 2010, 10, 27.	2.9	106
20	Population level mental distress in rural Ethiopia. BMC Psychiatry, 2014, 14, 194.	2.6	98
21	Development of a scalable mental healthcare plan for a rural district in Ethiopia. British Journal of Psychiatry, 2016, 208, s4-s12.	2.8	93
22	Evaluation of district mental healthcare plans: The PRIME consortium methodology. British Journal of Psychiatry, 2016, 208, s63-s70.	2.8	92
23	Lessons learned in developing community mental health care in Africa. World Psychiatry, 2010, 9, 185-189.	10.4	85
24	Service user involvement in mental health system strengthening in a rural African setting: qualitative study. BMC Psychiatry, 2017, 17, 187.	2.6	83
25	Service user and caregiver involvement in mental health system strengthening in low- and middle-income countries: a cross-country qualitative study. Epidemiology and Psychiatric Sciences, 2018, 27, 29-39.	3.9	81
26	"l cry every day and night, I have my son tied in chains― physical restraint of people with schizophrenia in community settings in Ethiopia. Globalization and Health, 2017, 13, 47.	4.9	78
27	Generating evidence to narrow the treatment gap for mental disorders in sub-Saharan Africa: rationale, overview and methods of AFFIRM. Epidemiology and Psychiatric Sciences, 2015, 24, 233-240.	3.9	77
28	District mental healthcare plans for five low-and middle-income countries: Commonalities, variations and evidence gaps. British Journal of Psychiatry, 2016, 208, s47-s54.	2.8	73
29	Efficacy of interventions to reduce coercive treatment in mental health services: umbrella review of randomised evidence. British Journal of Psychiatry, 2021, 218, 185-195.	2.8	73
30	Prevalence and risk factors for soil-transmitted helminth infection in mothers and their infants in Butajira, Ethiopia: a population based study. BMC Public Health, 2010, 10, 21.	2.9	71
31	Apo E genotypes in multiple sclerosis, Parkinson's disease, schwannomas and late-onset Alzheimer's disease. Molecular and Cellular Probes, 1994, 8, 519-525.	2.1	69
32	Validation of the World Health Organization Disability Assessment Schedule in people with severe mental disorders in rural Ethiopia. Health and Quality of Life Outcomes, 2017, 15, 64.	2.4	64
33	Communityâ€based mental health care in Africa: mental health workers' views. World Psychiatry, 2008, 7, 54-57.	10.4	63
34	Community-based rehabilitation intervention for people with schizophrenia in Ethiopia (RISE): a 12 month mixed methods pilot study. BMC Psychiatry, 2018, 18, 250.	2.6	63
35	Impact of perinatal somatic and common mental disorder symptoms on functioning in Ethiopian women: The P-MaMiE population-based cohort study. Journal of Affective Disorders, 2012, 136, 340-349.	4.1	62
36	Maternal common mental disorders and infant development in Ethiopia: the P-MaMiE Birth Cohort. BMC Public Health, 2010, 10, 693.	2.9	61

#	Article	IF	CITATIONS
37	Treatment gap and help-seeking for postpartum depression in a rural African setting. BMC Psychiatry, 2016, 16, 196.	2.6	61
38	The effect of maternal common mental disorders on infant undernutrition in Butajira, Ethiopia: The P-MaMiE study. BMC Psychiatry, 2010, 10, 32.	2.6	60
39	Perspectives on reasons for non-adherence to medication in persons with schizophrenia in Ethiopia: a qualitative study of patients, caregivers and health workers. BMC Psychiatry, 2013, 13, 168.	2.6	60
40	Postnatal mental distress in relation to the sociocultural practices of childbirth: An exploratory qualitative study from Ethiopia. Social Science and Medicine, 2009, 69, 1211-1219.	3.8	58
41	Treatment gap, help-seeking, stigma and magnitude of alcohol use disorder in rural Ethiopia. Substance Abuse Treatment, Prevention, and Policy, 2019, 14, 4.	2.2	58
42	Perinatal mental distress and infant morbidity in Ethiopia: a cohort study. Archives of Disease in Childhood: Fetal and Neonatal Edition, 2011, 96, F59-F64.	2.8	56
43	Psychological interventions for common mental disorders in women experiencing intimate partner violence in low-income and middle-income countries: a systematic review and meta-analysis. Lancet Psychiatry,the, 2020, 7, 173-190.	7.4	56
44	Participatory planning of a primary care service for people with severe mental disorders in rural Ethiopia. Health Policy and Planning, 2016, 31, 367-376.	2.7	55
45	Health systems context(s) for integrating mental health into primary health care in six Emerald countries: a situation analysis. International Journal of Mental Health Systems, 2017, 11, 7.	2.7	55
46	Ethical and professional challenges in mental health care in low- and middle-income countries. International Review of Psychiatry, 2010, 22, 245-251.	2.8	54
47	Alcohol use disorders and associated factors among people living with HIV who are attending services in south west Ethiopia. BMC Research Notes, 2014, 7, 828.	1.4	54
48	Migration and Mental Health: a Study of Low-Income Ethiopian Women Working in Middle Eastern Countries. International Journal of Social Psychiatry, 2009, 55, 557-568.	3.1	53
49	Untreated depression and tuberculosis treatment outcomes, quality of life and disability, Ethiopia. Bulletin of the World Health Organization, 2018, 96, 243-255.	3.3	53
50	Perceived challenges and opportunities arising from integration of mental health into primary care: a cross-sectional survey of primary health care workers in south-west Ethiopia. BMC Health Services Research, 2014, 14, 113.	2.2	52
51	Suicide and suicide attempts in people with severe mental disorders in Butajira, Ethiopia: 10Âyear follow-up of a population-based cohort. BMC Psychiatry, 2014, 14, 150.	2.6	51
52	Scaling up integrated primary mental health in six low- and middle-income countries: obstacles, synergies and implications for systems reform. BJPsych Open, 2019, 5, e69.	0.7	51
53	Developing capacity-building activities for mental health system strengthening in low- and middle-income countries for service users and caregivers, service planners, and researchers. Epidemiology and Psychiatric Sciences, 2018, 27, 11-21.	3.9	50
54	Challenges and opportunities to improve autism services in low-income countries: lessons from a situational analysis in Ethiopia. Global Mental Health (Cambridge, England), 2016, 3, e21.	2.5	48

#	Article	IF	CITATIONS
55	Adapting and pre-testing the World Health Organization's Caregiver Skills Training programme for autism and other developmental disorders in a very low-resource setting: Findings from Ethiopia. Autism, 2020, 24, 51-63.	4.1	48
56	Burden of mental disorders and unmet needs among street homeless people in Addis Ababa, Ethiopia. BMC Medicine, 2014, 12, 138.	5.5	47
57	Depression and disability in people with podoconiosis: a comparative cross-sectional study in rural Northern Ethiopia. International Health, 2016, 8, 124-131.	2.0	47
58	Scaling-up services for psychosis, depression and epilepsy in sub-Saharan Africa and South Asia: development and application of a mental health systems planning tool (OneHealth). Epidemiology and Psychiatric Sciences, 2017, 26, 234-244.	3.9	47
59	Depression among patients with tuberculosis: determinants, course and impact on pathways to care and treatment outcomes in a primary care setting in southern Ethiopia—a study protocol. BMJ Open, 2015, 5, e007653.	1.9	46
60	Perinatal mental health care in a rural African district, Uganda: a qualitative study of barriers, facilitators and needs. BMC Health Services Research, 2016, 16, 295.	2.2	46
61	Antenatal depressive symptoms and maternal health care utilisation: a population-based study of pregnant women in Ethiopia. BMC Pregnancy and Childbirth, 2016, 16, 301.	2.4	46
62	Hazardous alcohol use and associated factors in a rural Ethiopian district: a cross-sectional community survey. BMC Public Health, 2016, 16, 218.	2.9	46
63	Conceptualisation of job-related wellbeing, stress and burnout among healthcare workers in rural Ethiopia: a qualitative study. BMC Health Services Research, 2017, 17, 412.	2.2	46
64	Developing a Theory of Change model of service user and caregiver involvement in mental health system strengthening in primary health care in rural Ethiopia. International Journal of Mental Health Systems, 2020, 14, 51.	2.7	46
65	Development of a Community-Based Rehabilitation Intervention for People with Schizophrenia in Ethiopia. PLoS ONE, 2015, 10, e0143572.	2.5	46
66	Between life and death: exploring the sociocultural context of antenatal mental distress in rural Ethiopia. Archives of Women's Mental Health, 2010, 13, 385-393.	2.6	45
67	Involvement of people with schizophrenia in decision-making in rural Ethiopia: a qualitative study. Globalization and Health, 2018, 14, 85.	4.9	45
68	Catastrophic health expenditure and impoverishment in households of persons with depression: a cross-sectional, comparative study in rural Ethiopia. BMC Public Health, 2019, 19, 930.	2.9	44
69	Coping strategies of women with postpartum depression symptoms in rural Ethiopia: a cross-sectional community study. BMC Psychiatry, 2018, 18, 41.	2.6	43
70	Task-sharing of psychological treatment for antenatal depression in Khayelitsha, South Africa: Effects on antenatal and postnatal outcomes in an individual randomised controlled trial. Behaviour Research and Therapy, 2020, 130, 103466.	3.1	43
71	"Like a doctor, like a brotherâ€. Achieving competence amongst lay health workers delivering community-based rehabilitation for people with schizophrenia in Ethiopia. PLoS ONE, 2021, 16, e0246158.	2.5	43
72	Exposure to <i><scp>H</scp>elicobacter pylori</i> infection in early childhood and the risk of allergic disease and atopic sensitization: a longitudinal birth cohort study. Clinical and Experimental Allergy, 2014, 44, 563-571.	2.9	42

#	Article	IF	CITATIONS
73	Burden and presentation of depression among newly diagnosed individuals with TB in primary care settings in Ethiopia. BMC Psychiatry, 2017, 17, 57.	2.6	42
74	Maternal depression in low- and middle-income countries. International Health, 2013, 5, 4-5.	2.0	41
75	Mental health financing challenges, opportunities and strategies in low- and middle-income countries: findings from the Emerald project. BJPsych Open, 2019, 5, e68.	0.7	41
76	Participatory action research to pilot a model of mental health service user involvement in an Ethiopian rural primary healthcare setting: study protocol. Research Involvement and Engagement, 2020, 6, 2.	2.9	41
77	Antenatal depressive symptoms and perinatal complications: a prospective study in rural Ethiopia. BMC Psychiatry, 2017, 17, 301.	2.6	40
78	Explanatory models of depression in sub-Saharan Africa: Synthesis of qualitative evidence. Social Science and Medicine, 2020, 246, 112760.	3.8	40
79	Acute psychiatric care: approaches to increasing the range of services and improving access and quality of care. World Psychiatry, 2022, 21, 220-236.	10.4	40
80	Violence and violent victimization in people with severe mental illness in a rural low-income country setting: A comparative cross-sectional community study. Schizophrenia Research, 2014, 152, 275-282.	2.0	39
81	Task sharing for the care of severe mental disorders in a low-income country (TaSCS): study protocol for a randomised, controlled, non-inferiority trial. Trials, 2016, 17, 76.	1.6	39
82	Homelessness and severe mental illness in low- and middle-income countries: scoping review. BJPsych Open, 2019, 5, e57.	0.7	39
83	Beyond the Biomedical: Community Resources for Mental Health Care in Rural Ethiopia. PLoS ONE, 2015, 10, e0126666.	2.5	39
84	Mental distress and podoconiosis in Northern Ethiopia: a comparative cross-sectional study. International Health, 2015, 7, 16-25.	2.0	37
85	Information systems for mental health in six low and middle income countries: cross country situation analysis. International Journal of Mental Health Systems, 2016, 10, 60.	2.7	37
86	Indicators for routine monitoring of effective mental healthcare coverage in low- and middle-income settings: a Delphi study. Health Policy and Planning, 2016, 31, 1100-1106.	2.7	36
87	Food insecurity, mental health and quality of life among people living with HIV commencing antiretroviral treatment in Ethiopia: a cross-sectional study. Health and Quality of Life Outcomes, 2016, 14, 37.	2.4	36
88	Service user and care giver involvement in mental health system strengthening in Nepal: a qualitative study on barriers and facilitating factors. International Journal of Mental Health Systems, 2017, 11, 30.	2.7	36
89	Recognition of depression by primary care clinicians in rural Ethiopia. BMC Family Practice, 2017, 18, 56.	2.9	36
90	Community-based Rehabilitation Intervention for people with Schizophrenia in Ethiopia (RISE): study protocol for a cluster randomised controlled trial. Trials, 2016, 17, 299.	1.6	34

#	Article	IF	CITATIONS
91	Health system governance to support scale up of mental health care in Ethiopia: a qualitative study. International Journal of Mental Health Systems, 2017, 11, 38.	2.7	34
92	National and regional prevalence of posttraumatic stress disorder in sub-Saharan Africa: A systematic review and meta-analysis. PLoS Medicine, 2020, 17, e1003090.	8.4	34
93	Engaging and staying engaged: a phenomenological study of barriers to equitable access to mental healthcare for people with severe mental disorders in a rural African setting. International Journal for Equity in Health, 2017, 16, 156.	3.5	33
94	Impact of district mental health care plans on symptom severity and functioning of patients with priority mental health conditions: the Programme for Improving Mental Health Care (PRIME) cohort protocol. BMC Psychiatry, 2018, 18, 61.	2.6	33
95	Under detection of depression in primary care settings in low and middle-income countries: a systematic review and meta-analysis. Systematic Reviews, 2022, 11, 21.	5.3	32
96	Non-fatal suicidal behaviour in rural Ethiopia: a cross-sectional facility- and population-based study. BMC Psychiatry, 2016, 16, 75.	2.6	31
97	Training needs and perspectives of community health workers in relation to integrating child mental health care into primary health care in a rural setting in sub-Saharan Africa: a mixed methods study. International Journal of Mental Health Systems, 2017, 11, 15.	2.7	31
98	Evaluating capacity-building for mental health system strengthening in low- and middle-income countries for service users and caregivers, service planners and researchers. Epidemiology and Psychiatric Sciences, 2018, 27, 3-10.	3.9	31
99	Improving collaboration between primary care and mental health services. World Journal of Biological Psychiatry, 2019, 20, 748-765.	2.6	31
100	Next steps for meeting the needs of people with severe mental illness in low- and middle-income countries. Epidemiology and Psychiatric Sciences, 2017, 26, 348-354.	3.9	30
101	Service user involvement for mental health system strengthening in India: a qualitative study. BMC Psychiatry, 2016, 16, 269.	2.6	29
102	Postpartum depressive symptoms in the context of high social adversity and reproductive health threats: a population-based study. International Journal of Mental Health Systems, 2018, 12, 42.	2.7	29
103	Food insecurity among people with severe mental disorder in a rural Ethiopian setting: a comparative, population-based study. Epidemiology and Psychiatric Sciences, 2019, 28, 397-407.	3.9	29
104	Restoring the person's life': a qualitative study to inform development of care for people with severe mental disorders in rural Ethiopia. Epidemiology and Psychiatric Sciences, 2017, 26, 43-52.	3.9	28
105	Long-term clinical course and outcome of schizophrenia in rural Ethiopia: 10-year follow-up of a population-based cohort. Schizophrenia Research, 2015, 161, 414-420.	2.0	27
106	Adapting the Bayley Scales of infant and toddler development in Ethiopia: evaluation of reliability and validity. Child: Care, Health and Development, 2016, 42, 699-708.	1.7	27
107	Global mental health and schizophrenia. Current Opinion in Psychiatry, 2018, 31, 193-199.	6.3	27
108	Using a mentorship model to localise the Practical Approach to Care Kit (PACK): from South Africa to Ethiopia. BMJ Global Health, 2019, 3, e001108.	4.7	27

#	Article	IF	CITATIONS
109	Ethiopian community health workers' beliefs and attitudes towards children with autism: Impact of a brief training intervention. Autism, 2019, 23, 39-49.	4.1	27
110	Impact of integrated district level mental health care on clinical and social outcomes of people with severe mental illness in rural Ethiopia: an intervention cohort study. Epidemiology and Psychiatric Sciences, 2020, 29, e45.	3.9	27
111	Perceptions and experiences of stigma among parents of children with developmental disorders in Ethiopia: A qualitative study. Social Science and Medicine, 2020, 256, 113034.	3.8	27
112	Measuring common mental disorders in women in Ethiopia. Social Psychiatry and Psychiatric Epidemiology, 2008, 43, 653-9.	3.1	26
113	Khat chewing in persons with severe mental illness in Ethiopia: A qualitative study exploring perspectives of patients and caregivers. Transcultural Psychiatry, 2011, 48, 455-472.	1.6	26
114	Conceptualizing and contextualizing functioning in people with severe mental disorders in rural Ethiopia: a qualitative study. BMC Psychiatry, 2015 , 15 , 34 .	2.6	26
115	Psychological interventions for bipolar disorder in low- and middle-income countries: systematic review. BJPsych Open, 2018, 4, 375-384.	0.7	26
116	Burnout among primary healthcare workers during implementation of integrated mental healthcare in rural Ethiopia: a cohort study. Human Resources for Health, 2019, 17, 58.	3.1	26
117	The D-score: a metric for interpreting the early development of infants and toddlers across global settings. BMJ Global Health, 2019, 4, e001724.	4.7	25
118	Strengthening mental health systems in low- and middle-income countries: recommendations from the Emerald programme. BJPsych Open, 2019, 5, e73.	0.7	25
119	Sociocultural practices in Ethiopia: association with onset and persistence of postnatal common mental disorders. British Journal of Psychiatry, 2010, 197, 468-475.	2.8	24
120	Prevalence and risk factors of wheeze and eczema in 1â€yearâ€old children: the Butajira birth cohort, Ethiopia. Clinical and Experimental Allergy, 2010, 40, 619-626.	2.9	24
121	Equitable access to integrated primary mental healthcare for people with severe mental disorders in Ethiopia: a formative study. International Journal for Equity in Health, 2016, 15, 121.	3 . 5	24
122	Adaptation and validation of the short version WHOQOL-HIV in Ethiopia. International Journal of Mental Health Systems, 2016, 10, 29.	2.7	24
123	Moving towards universal health coverage for mental disorders in Ethiopia. International Journal of Mental Health Systems, 2019, 13, 11.	2.7	24
124	Partnerships in a Global Mental Health Research Programmeâ€"the Example of PRIME. Global Social Welfare, 2019, 6, 159-175.	1.9	24
125	The psychosis treatment gap and its consequences in rural Ethiopia. BMC Psychiatry, 2019, 19, 325.	2.6	23
126	Catastrophic out-of-pocket payments for households of people with severe mental disorder: a comparative study in rural Ethiopia. International Journal of Mental Health Systems, 2019, 13, 39.	2.7	23

#	Article	IF	Citations
127	Effects of Early Life Paracetamol Use on the Incidence of Allergic Disease and Sensitization: 5 Year Follow-Up of an Ethiopian Birth Cohort. PLoS ONE, 2014, 9, e93869.	2.5	22
128	Building the capacity of policy-makers and planners to strengthen mental health systems in low- and middle-income countries: a systematic review. BMC Health Services Research, 2016, 16, 601.	2.2	22
129	Antenatal depressive symptoms and utilisation of delivery and postnatal care: a prospective study in rural Ethiopia. BMC Pregnancy and Childbirth, 2017, 17, 206.	2.4	22
130	Household economic costs associated with mental, neurological and substance use disorders: a cross-sectional survey in six low- and middle-income countries. BJPsych Open, 2019, 5, e34.	0.7	21
131	Community-based rehabilitation intervention for people with schizophrenia in Ethiopia (RISE): results of a 12-month cluster-randomised controlled trial. The Lancet Global Health, 2022, 10, e530-e542.	6.3	21
132	Khat use in people living with HIV: a facility-based cross-sectional survey from South West Ethiopia. BMC Psychiatry, 2015, 15, 69.	2.6	20
133	Development and validation of a contextual measure of functioning for people living with severe mental disorders in rural Africa. BMC Psychiatry, 2016, 16, 311.	2.6	20
134	Developing a measure of mental health service satisfaction for use in low income countries: a mixed methods study. BMC Health Services Research, 2017, 17, 183.	2.2	20
135	Ethics in global health research: the need for balance. The Lancet Global Health, 2015, 3, e516-e517.	6.3	19
136	Equitable access to mental healthcare integrated in primary care for people with severe mental disorders in rural Ethiopia: a community-based cross-sectional study. International Journal of Mental Health Systems, 2019, 13, 78.	2.7	19
137	Dental caries experience and associated factors in adults: a cross-sectional community survey within Ethiopia. BMC Public Health, 2021, 21, 180.	2.9	19
138	Help seeking and suicidality among people with epilepsy in a rural low income country setting: cross-sectional survey. International Journal of Mental Health Systems, 2017, 11, 44.	2.7	18
139	Maternal depression symptoms are highly prevalent among food-insecure households in Ethiopia. Public Health Nutrition, 2018, 21, 849-856.	2.2	18
140	Functional impairment among people with severe and enduring mental disorder in rural Ethiopia: a cross-sectional study. Social Psychiatry and Psychiatric Epidemiology, 2018, 53, 803-814.	3.1	18
141	Mental health problems and socioeconomic disadvantage: a controlled household study in rural Ethiopia. International Journal for Equity in Health, 2019, 18, 121.	3.5	18
142	Developing mental health research in sub-Saharan Africa: capacity building in the AFFIRM project. Global Mental Health (Cambridge, England), 2016, 3, e33.	2.5	17
143	Evaluation of capacity-building strategies for mental health system strengthening in low- and middle-income countries for service users and caregivers, policymakers and planners, and researchers. BJPsych Open, 2019, 5, e67.	0.7	17
144	Enhancing mental health research capacity: emerging voices from the National Institute of Mental Health (NIMH) global hubs. International Journal of Mental Health Systems, 2019, 13, 21.	2.7	17

#	Article	IF	Citations
145	Antenatal predictors of incident and persistent postnatal depressive symptoms in rural Ethiopia: a population-based prospective study. Reproductive Health, 2019, 16, 28.	3.1	17
146	Characteristics of perinatal depression in rural central, India: a cross-sectional study. International Journal of Mental Health Systems, 2018, 12, 68.	2.7	16
147	Mental health research capacity building in sub-Saharan Africa: the African Mental Health Research Initiative. Global Mental Health (Cambridge, England), 2020, 7, e8.	2.5	16
148	Missed opportunity for alcohol use disorder screening and management in primary health care facilities in northern rural Tanzania: a cross-sectional survey. Substance Abuse Treatment, Prevention, and Policy, 2022, 17, .	2.2	16
149	Adjuvant therapy with minocycline for schizophrenia (The MINOS Trial): study protocol for a double-blind randomized placebo-controlled trial. Trials, 2013, 14, 406.	1.6	15
150	Impact of maternal common mental disorders on child educational outcomes at 7 and 9 years: a population-based cohort study in Ethiopia. BMJ Open, 2018, 8, e018916.	1.9	15
151	A guide to systems-level, participatory, theory-informed implementation research in global health. BMJ Global Health, 2021, 6, e005365.	4.7	15
152	Proposed Training Areas for Global Mental Health Researchers. Academic Psychiatry, 2016, 40, 679-685.	0.9	14
153	Implementing integrated services for people with epilepsy in primary care in Ethiopia: a qualitative study. BMC Health Services Research, 2018, 18, 372.	2.2	14
154	"We need more big trees as well as the grass roots― going beyond research capacity building to develop sustainable careers in mental health research in African countries. International Journal of Mental Health Systems, 2020, 14, 66.	2.7	13
155	Psychotherapy Knowledge Translation and Interpersonal Psychotherapy: Using Best-Education Practices to Transform Mental Health Care in Canada and Ethiopia. American Journal of Psychotherapy, 2014, 68, 463-488.	1.2	12
156	Care for bipolar disorder in LMICs needs evidence from local settings. Lancet Psychiatry,the, 2015, 2, 772-773.	7.4	12
157	The management of adult psychiatric emergencies in low-income and middle-income countries: a systematic review. Lancet Psychiatry,the, 2015, 2, 540-547.	7.4	12
158	Development of programme theory for integration of service user and caregiver involvement in mental health system strengthening: protocol for realist systematic review. International Journal of Mental Health Systems, 2018, 12, 41.	2.7	12
159	Experience of implementing new mental health indicators within information systems in six low- and middle-income countries. BJPsych Open, 2019, 5, e71.	0.7	12
160	Stakeholder perspectives on antenatal depression and the potential for psychological intervention in rural Ethiopia: a qualitative study. BMC Pregnancy and Childbirth, 2020, 20, 371.	2.4	12
161	Integration of Alcohol Use Disorder Interventions in General Health Care Settings in Sub-Saharan Africa: A Scoping Review. Frontiers in Psychiatry, 2022, 13, 822791.	2.6	12
162	Comparison of liaison psychiatry service models for older patients. Psychiatric Bulletin, 2004, 28, 171-173.	0.3	11

#	Article	IF	CITATIONS
163	Fatigue as a manifestation of psychosocial distress in a lowâ€income country: a populationâ€based panel study. Tropical Medicine and International Health, 2016, 21, 365-372.	2.3	11
164	The experience of mental health service users in health system strengthening: lessons from Uganda. International Journal of Mental Health Systems, 2019, 13, 60.	2.7	11
165	Perinatal mental health around the world: priorities for research and service development in Africa. BJPsych International, 2020, 17, 56-59.	1.4	11
166	Towards racial equity in global mental health research. Lancet Psychiatry, the, 2021, 8, 553-555.	7.4	11
167	HeAlth System StrEngThening in four sub-Saharan African countries (ASSET) to achieve high-quality, evidence-informed surgical, maternal and newborn, and primary care: protocol for pre-implementation phase studies. Global Health Action, 2022, 15, 1987044.	1.9	11
168	Efficacy and cost-effectiveness of task-shared care for people with severe mental disorders in Ethiopia (TaSCS): a single-blind, randomised, controlled, phase 3 non-inferiority trial. Lancet Psychiatry,the, 2022, 9, 59-71.	7.4	11
169	Body composition during early infancy and developmental progression from 1 to 5 years of age: the Infant Anthropometry and Body Composition (iABC) cohort study among Ethiopian children. British Journal of Nutrition, 2018, 119, 1263-1273.	2.3	10
170	Effective methods for knowledge transfer to strengthen mental health systems in low- and middle-income countries. BJPsych Open, 2019, 5, e72.	0.7	10
171	Rural vs urban residence and experience of discrimination among people with severe mental illnesses in Ethiopia. BMC Psychiatry, 2019, 19, 340.	2.6	10
172	Frequent Use of Paracetamol and Risk of Allergic Disease Among Women in an Ethiopian Population. PLoS ONE, 2011, 6, e22551.	2.5	10
173	Relation between body composition at birth and child development at 2 years of age: a prospective cohort study among Ethiopian children. European Journal of Clinical Nutrition, 2017, 71, 1411-1417.	2.9	9
174	"He can send her to her parentsâ€. The interaction between marriageability, gender and serious mental illness in rural Ethiopia. BMC Psychiatry, 2019, 19, 315.	2.6	9
175	Impact of integrated mental health care on food insecurity of households of people with severe mental illness in a rural African district: a communityâ€based, controlled before–after study. Tropical Medicine and International Health, 2020, 25, 414-423.	2.3	9
176	Common Mental Disorder Symptoms among Patients with Malaria Attending Primary Care in Ethiopia: A Cross-Sectional Survey. PLoS ONE, 2014, 9, e108923.	2.5	9
177	Mental health stigma and discrimination in Ethiopia: evidence synthesis to inform stigma reduction interventions. International Journal of Mental Health Systems, 2022, 16, .	2.7	9
178	The effect of nutritional supplementation on quality of life in people living with <scp>HIV</scp> : a randomised controlled trial. Tropical Medicine and International Health, 2016, 21, 735-742.	2.3	8
179	The effect of gender on the long-term course and outcome of schizophrenia in rural Ethiopia: a population-based cohort. Social Psychiatry and Psychiatric Epidemiology, 2020, 55, 1581-1591.	3.1	8
180	Implementation science protocol for a participatory, theory-informed implementation research programme in the context of health system strengthening in sub-Saharan Africa (ASSET-ImplementER). BMJ Open, 2021, 11, e048742.	1.9	8

#	Article	IF	CITATIONS
181	Group problem solving therapy for perinatal depression in primary health care settings in rural Uganda: an intervention cohort study. BMC Pregnancy and Childbirth, 2021, 21, 584.	2.4	8
182	Body Composition during Early Infancy and Mental Health Outcomes at 5 Years of Age: A Prospective Cohort Study of Ethiopian Children. Journal of Pediatrics, 2018, 200, 225-231.	1.8	7
183	Functional impairment among people diagnosed with depression in primary healthcare in rural Ethiopia: a comparative cross-sectional study. International Journal of Mental Health Systems, 2019, 13, 50.	2.7	7
184	Food insecurity and work impairment in people with severe mental disorders in a rural district of Ethiopia: a cross-sectional survey. Social Psychiatry and Psychiatric Epidemiology, 2019, 54, 1055-1066.	3.1	7
185	Trauma exposure, depression, suicidal ideation, and alcohol use in people with severe mental disorder in Ethiopia. Social Psychiatry and Psychiatric Epidemiology, 2019, 54, 835-842.	3.1	7
186	Comorbid mental disorders and quality of life of people with epilepsy attending primary health care clinics in rural Ethiopia. PLoS ONE, 2021, 16, e0238137.	2.5	7
187	Understanding the key processes of excellence as a prerequisite to establishing academic centres of excellence in Africa. BMC Medical Education, 2021, 21, 36.	2.4	7
188	Contextual equipoise: a novel concept to inform ethical implications for implementation research in low-income and middle-income countries. BMJ Global Health, 2020, 5, e003456.	4.7	7
189	Impact of child emotional and behavioural difficulties on educational outcomes of primary school children in Ethiopia: a population-based cohort study. Child and Adolescent Psychiatry and Mental Health, 2020, 14, 22.	2.5	7
190	Perinatal mental distress in a rural Ethiopian community: a critical examination of psychiatric labels. BMC Psychiatry, 2020, 20, 223.	2.6	6
191	Problem solving therapy (PST) tailored for intimate partner violence (IPV) versus standard PST and enhanced usual care for pregnant women experiencing IPV in rural Ethiopia: protocol for a randomised controlled feasibility trial. Trials, 2020, 21, 454.	1.6	6
192	Health service costs and their association with functional impairment among adults receiving integrated mental health care in five low- and middle-income countries: the PRIME cohort study. Health Policy and Planning, 2020, 35, 567-576.	2.7	6
193	Brief problem-solving therapy for antenatal depressive symptoms in primary care in rural Ethiopia: protocol for a randomised, controlled feasibility trial. Pilot and Feasibility Studies, 2021, 7, 35.	1.2	6
194	"We improved our life because I cut my drinking― Qualitative analysis of a brief intervention for people with alcohol use disorder in Ethiopian primary health care. Journal of Substance Abuse Treatment, 2022, 132, 108636.	2.8	6
195	The path to global equity in mental health care in the context of COVID-19. Lancet, The, 2021, 398, 1670-1672.	13.7	6
196	The role of advocacy and empowerment in shaping service development for families raising children with developmental disabilities. Health Expectations, 2022, 25, 1882-1891.	2.6	6
197	The experiences of parents raising children with developmental disabilities in Ethiopia. Autism, 2023, 27, 539-551.	4.1	6
198	Mental health impact on the unmet need for family planning and fertility rate in rural Ethiopia: a population-based cohort study. Epidemiology and Psychiatric Sciences, 2020, 29, e160.	3.9	5

#	Article	IF	CITATIONS
199	Announcing the Lancet Psychiatry Commission on Psychoses in Global Context. Lancet Psychiatry, the, 2021, 8, 743-744.	7.4	5
200	Pills and prayers: a comparative qualitative study of community conceptualisations of pre-eclampsia and pluralistic care in Ethiopia, Haiti and Zimbabwe. BMC Pregnancy and Childbirth, 2021, 21, 716.	2.4	5
201	Teaching psychiatry in Ethiopia. International Psychiatry: Bulletin of the Board of International Affairs of the Royal College of Psychiatrists, 2006, 3, 43-46.	0.1	4
202	Perinatal common mental disorders and child survival in <scp>E</scp> thiopia. Journal of Paediatrics and Child Health, 2014, 50, 57-63.	0.8	4
203	Developing sustainable capacity-building in mental health research: implementation outcomes of training of trainers in systematic reviewing. Global Health Action, 2020, 13, 1715325.	1.9	4
204	Common perinatal mental disorders and postâ€infancy child development in rural Ethiopia: A populationâ€based cohort study. Tropical Medicine and International Health, 2022, 27, 251-261.	2.3	4
205	Understandings of depression among community members and primary healthcare attendees in rural Ethiopia: A qualitative study. Transcultural Psychiatry, 2023, 60, 412-427.	1.6	4
206	Empowerment training to support service user involvement in mental health system strengthening in rural Ethiopia: a mixed-methods pilot study. BMC Health Services Research, 2022, 22, .	2.2	4
207	Mapping the evidence on pharmacological interventions for non-affective psychosis in humanitarian non-specialised settings: a UNHCR clinical guidance. BMC Medicine, 2017, 15, 197.	5.5	3
208	Is depression associated with pathways to care and diagnosis delay in people with tuberculosis in Ethiopia?. Global Mental Health (Cambridge, England), 2019, 6, e20.	2.5	3
209	The 10-year functional outcome of schizophrenia in Butajira, Ethiopia. Heliyon, 2019, 5, e01272.	3.2	3
210	Incidence of depression in people with newly diagnosed tuberculosis in Ethiopia: a cohort study. Global Mental Health (Cambridge, England), 2020, 7, e1.	2.5	3
211	Psychotherapy in a resource-constrained setting: Understanding context for adapting and integrating a brief psychological intervention into primary care Psychotherapy, 2021, 58, 557-575.	1.2	3
212	Family planning for women with severe mental illness in rural Ethiopia: a qualitative study. Reproductive Health, 2021, 18, 191.	3.1	3
213	Development of a psychological intervention for people with bipolar disorder in rural Ethiopia. BJPsych Open, 2021, 7, .	0.7	3
214	Adaptation and piloting of an integrated intervention model for alcohol use disorders in primary healthcare in rural Tanzania: a study protocol. BMJ Open, 2020, 10, e038615.	1.9	3
215	Adaptation and validation of two autism-related measures of skills and quality of life in Ethiopia. Autism, 2021, , 136236132110507.	4.1	3
216	Implementation strategy in collaboration with people with lived experience of mental illness to reduce stigma among primary care providers in Nepal (RESHAPE): protocol for a type 3 hybrid implementation effectiveness cluster randomized controlled trial. Implementation Science, 2022, 17, .	6.9	3

#	Article	IF	Citations
217	Using participatory action research to pilot a model of service user and caregiver involvement in mental health system strengthening in Ethiopian primary healthcare: a case study. International Journal of Mental Health Systems, 2022, 16, .	2.7	3
218	Impact of perinatal and repeated maternal common mental disorders on educational outcomes of primary school children in rural Ethiopia: population-based cohort study. BJPsych Open, 2019, 5, e87.	0.7	2
219	The 11-year functional outcome of bipolar and major depressive disorders in Butajira, Ethiopia. Neurology Psychiatry and Brain Research, 2019, 32, 68-76.	2.0	2
220	Why doesn't God say "enough� Experiences of living with bipolar disorder in rural Ethiopia. Social Science and Medicine, 2021, 270, 113625.	3.8	2
221	Posttraumatic stress disorder intervention for people with severe mental illness in a low-income country primary care setting: a randomized feasibility trial protocol. Pilot and Feasibility Studies, 2021, 7, 149.	1.2	2
222	Performance of primary health care workers in detection of mental disorders comorbid with epilepsy in rural Ethiopia. BMC Family Practice, 2021, 22, 204.	2.9	2
223	Psychometric properties of screening questionnaires to detect depression in primary healthcare setting in rural Ethiopia. , 2022, 23, .		2
224	Detecting postnatal common mental disorders in Addis ababa, ethiopia: Validation of the Edinburgh postnatal depression scale and kessler scales. European Psychiatry, 2011, 26, 1689-1689.	0.2	1
225	Sustaining international careers: a peer group for psychiatrists working in global mental health. BJPsych International, 2015, 12, 5-7.	1.4	1
226	A leap of faith for more effective mental health care. Lancet, The, 2020, 396, 584-585.	13.7	1
227	Guided self-help to reduce psychological distress: what's the potential for scaling up?. The Lancet Global Health, 2020, 8, e165-e166.	6.3	1
228	Pathways into and out of homelessness among people with severe mental illness in rural Ethiopia: a qualitative study. BMC Public Health, 2021, 21, 568.	2.9	1
229	Involvement of People with Lived Experience of Mental Health Conditions in Decision-Making to Improve Care in Rural Ethiopia., 2021,, 396-412.		1
230	Adaptation and piloting of an integrated intervention model for alcohol use disorders in primary healthcare in rural Tanzania: a study protocol. BMJ Open, 2020, 10, e038615.	1.9	1
231	Validation of the Communication Profile-Adapted in Ethiopian children with neurodevelopmental disorders. Global Mental Health (Cambridge, England), 2021, 8, e45.	2.5	1
232	Stakeholder experiences, attitudes and perspectives on inclusive education for children with developmental disabilities in sub-Saharan Africa: A systematic review of qualitative studies. Autism, 2022, 26, 1606-1625.	4.1	1
233	Taking the path less trodden: UK psychiatrists working in low- and middle-income countries. Psychiatric Bulletin, 2008, 32, 431-435.	0.3	0
234	COVID-19 and global mental health – Authors' reply. Lancet Psychiatry,the, 2021, 8, 459-460.	7.4	0

#	Article	IF	CITATIONS
235	Quality not just quantity: how health system strengthening is essential for scale up of quality mental health care. Epidemiology and Psychiatric Sciences, 2020, 29, e186.	3.9	0
236	Teaching psychiatry in Ethiopia. International Psychiatry: Bulletin of the Board of International Affairs of the Royal College of Psychiatrists, 2006, 3, 43-46.	0.1	0
237	MENTAL HEALTH OF HEALTHCARE WORKERS DURING COVID-19 PANDEMIC IN UKRAINE. Proceedings of the Shevchenko Scientific Society Medical Sciences, 2022, 66, .	0.3	O