Fatima Ribeiro-Dias

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6200012/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Interleukin-32γ in the Control of Acute Experimental Chagas Disease. Journal of Immunology Research, 2022, 2022, 1-9.	2.2	4
2	Tollâ€like receptor 10 controls TLR2â€induced cytokine production in monocytes from patients with Parkinson's disease. Journal of Neuroscience Research, 2021, 99, 2511-2524.	2.9	5
3	IL-15 enhances the capacity of primary human macrophages to control Leishmania braziliensis infection by IL-32/vitamin D dependent and independent pathways. Parasitology International, 2020, 76, 102097.	1.3	11
4	β-Glucan-Induced Trained Immunity Protects against Leishmania braziliensis Infection: a Crucial Role for IL-32. Cell Reports, 2019, 28, 2659-2672.e6.	6.4	102
5	Alterations in monocyte subsets and cytokine production after TLR activation in American Cutaneous Leishmaniasis. Parasite Immunology, 2019, 41, e12623.	1.5	3
6	Identification and characterization of Paracoccidioides lutzii proteins interacting with macrophages. Microbes and Infection, 2019, 21, 401-411.	1.9	12
7	Metacyclogenesis ofLeishmania (Viannia) guyanensis: a comprehensive study of the main transformation features in axenic culture and purification of metacyclic promastigotes by negative selection withBauhinia purpurealectin. Parasitology, 2019, 146, 716-727.	1.5	3
8	Human Interleukin-32Î ³ Plays a Protective Role in an Experimental Model of Visceral Leishmaniasis in Mice. Infection and Immunity, 2018, 86, .	2.2	14
9	Leishmania (Viannia) guyanensis in tegumentary leishmaniasis. Pathogens and Disease, 2018, 76, .	2.0	12
10	TLR4 and TLR2 activation is differentially associated with age during Parkinson's disease. Immunological Investigations, 2018, 47, 71-88.	2.0	9
11	The NOD2 receptor is crucial for immune responses towards New World Leishmania species. Scientific Reports, 2017, 7, 15219.	3.3	17
12	Interleukin 32: a novel player in the control of infectious diseases. Journal of Leukocyte Biology, 2017, 101, 39-52.	3.3	65
13	Cytokines and microbicidal molecules regulated by IL-32 in THP-1-derived human macrophages infected with New World Leishmania species. PLoS Neglected Tropical Diseases, 2017, 11, e0005413.	3.0	38
14	IL-32Î ³ promotes the healing of murine cutaneous lesions caused by Leishmania braziliensis infection in contrast to Leishmania amazonensis. Parasites and Vectors, 2017, 10, 336.	2.5	18
15	Leishmania (Viannia) braziliensis amastigotes induces the expression of TNFα and IL-10 by human peripheral blood mononuclear cells in vitro in a TLR4-dependent manner. Cytokine, 2016, 88, 184-192.	3.2	27
16	Essential role of leukotriene B4 on Leishmania (Viannia) braziliensis killing by human macrophages. Microbes and Infection, 2014, 16, 945-953.	1.9	27
17	Interleukin 32Î ³ (IL-32Î ³) is highly expressed in cutaneous and mucosal lesions of American Tegumentary Leishmaniasis patients: association with tumor necrosis factor (TNF) and IL-10. BMC Infectious Diseases, 2014, 14, 249.	2.9	25
18	Energetic metabolism of axenic promastigotes of Leishmania (Viannia) braziliensis. Experimental Parasitology, 2011, 128, 438-443.	1.2	9

#	Article	IF	CITATIONS
19	Crucial cytokine interactions in nitric oxide production induced by Mycoplasma arthritidis superantigen. Microbes and Infection, 2008, 10, 1543-1551.	1.9	7