

Sang-Won Lee

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6242133/publications.pdf>

Version: 2024-02-01

213
papers

2,326
citations

257450

24
h-index

345221

36
g-index

224
all docs

224
docs citations

224
times ranked

3489
citing authors

#	ARTICLE	IF	CITATIONS
1	Birmingham Vasculitis Activity Score und der Short Form 36-Item Health Survey als Prädiktoren aktueller depressiver Störungen bei Patienten mit antineutrophile-zytoplasmatische-Antikörper-assoziiierter Vaskulitis während der SARS-CoV-2-Pandemie. Zeitschrift Fur Rheumatologie, 2024, 83, 222-229.	1.0	1
2	Reclassification of previously diagnosed GPA patients using the 2022 ACR/EULAR classification criteria. Rheumatology, 2023, 62, 1179-1186.	1.9	8
3	2019 American College of Rheumatology/European League Against Rheumatism classification criteria for IgG4-related disease by Wallace et al. Annals of the Rheumatic Diseases, 2022, 81, e179-e179.	0.9	10
4	Increased prevalence rate of metabolic syndrome is an independent predictor of cardiovascular disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. Rheumatology International, 2022, 42, 291-302.	3.0	7
5	Rheumatoid factor positivity in antineutrophil cytoplasmic antibody-associated vasculitis: a distinct clinical entity or innocent bystander?. Rheumatology, 2022, 61, 1366-1375.	1.9	3
6	Nutrition Risk Index Score at Diagnosis Can Effectively Predict Poor Prognosis in Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. , 2022, 32, 423-431.		4
7	Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients. Journal of Clinical Rheumatology, 2022, 28, e81-e88.	0.9	4
8	A retrospective analysis of antineutrophil cytoplasmic antibody-associated vasculitis aiming for an equation prediction end-stage renal disease. Clinical Rheumatology, 2022, 41, 773-781.	2.2	5
9	Association Between Idiopathic Cutaneous Leukocytoclastic Angiitis and ANCA-negative Microscopic Polyangiitis. Journal of Rheumatic Diseases, 2022, 29, 40-45.	1.1	0
10	Clinical application of low erythrocyte sedimentation rate/high C-reactive protein to antineutrophil cytoplasmic antibody-associated vasculitis. Journal of Clinical Laboratory Analysis, 2022, 36, e24237.	2.1	2
11	Clinical characteristics and long-term outcomes in patients with mixed Class III/IV + V and pure proliferative lupus nephritis: A single-center experience. Lupus, 2022, 31, 588-595.	1.6	0
12	New body mass index for predicting prognosis in patients with antineutrophil cytoplasmic antibody-associated vasculitis. Journal of Clinical Laboratory Analysis, 2022, , e24357.	2.1	4
13	Clinical Features of Anti-Synthetase Syndrome Associated with Prognosis in Patients with Dermatomyositis and Polymyositis. Journal of Clinical Medicine, 2022, 11, 2052.	2.4	2
14	Incidence, prevalence and risk of stroke in patients with Takayasu arteritis: a nationwide population-based study in South Korea. Stroke and Vascular Neurology, 2022, 7, 149-157.	3.3	7
15	Anti-Citrullinated Peptide Antibody Expression and Its Association with Clinical Features and Outcomes in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. Medicina (Lithuania), 2022, 58, 558.	2.0	2
16	Effect of numbers of metabolic syndrome components on mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis with metabolic syndrome.. Clinical and Experimental Rheumatology, 2022, , .	0.8	0
17	Incidence and Patterns of Interstitial Lung Disease and Their Clinical Impact on Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Korean Single-Centre Observational Study. Journal of Immunology Research, 2022, 2022, 1-7.	2.2	3
18	Triglyceride and Glucose Index Predicts Acute Coronary Syndrome in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. Diagnostics, 2022, 12, 1486.	2.6	3

#	ARTICLE	IF	CITATIONS
19	Modified Body Mass Index at Diagnosis is a Useful Predictor of Mortality in Patients With Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , 2022, 29, 154-161.	1.1	0
20	Clinical Significance of Antineutrophil Cytoplasmic Antibody Positivity in Patients Infected with SARS-CoV-2. <i>Journal of Clinical Medicine</i> , 2022, 11, 4152.	2.4	3
21	Serum albumin, prealbumin, and ischemia-modified albumin levels in patients with ANCA-associated vasculitis: A prospective cohort study. <i>PLoS ONE</i> , 2022, 17, e0271055.	2.5	5
22	Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. <i>Clinical Rheumatology</i> , 2021, 40, 1423-1430.	2.2	2
23	Serological Biomarkers and Indices for the Current Activity and Prognosis of ANCA-Associated Vasculitis: Experience in a Single Centre in Korea. <i>Yonsei Medical Journal</i> , 2021, 62, 279.	2.2	8
24	Systemic inflammation response index predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Urology and Nephrology</i> , 2021, 53, 1631-1638.	1.4	4
25	Association Between Serum Alarmin Levels and Disease-specific Indices in Patients With Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis. <i>In Vivo</i> , 2021, 35, 1761-1768.	1.3	1
26	Fibrosis-5 predicts end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis without substantial liver diseases. <i>Clinical and Experimental Medicine</i> , 2021, 21, 399-406.	3.6	4
27	Fibrinogen to albumin ratio reflects the activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e23731.	2.1	7
28	Serum chitinase-3-like 1 protein is a useful biomarker to assess disease activity in ANCA-associated vasculitis: an observational study. <i>Arthritis Research and Therapy</i> , 2021, 23, 77.	3.5	8
29	Correlation between serum cysteine-rich protein 61 and disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2021, 40, 3703-3710.	2.2	1
30	Risk of Stroke in Systemic Necrotizing Vasculitis: A Nationwide Study Using the National Claims Database. <i>Frontiers in Immunology</i> , 2021, 12, 629902.	4.8	8
31	Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 441-446.	1.7	3
32	Unclassifiable repeated antineutrophil cytoplasmic antibody (ANCA) positivity in diseases other than ANCA-associated vasculitis. <i>Zeitschrift Fur Rheumatologie</i> , 2021, , 1.	1.0	4
33	Association between follistatin-related protein 1 and the functional status of patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Chinese Medical Journal</i> , 2021, 134, 1168-1174.	2.3	0
34	Total Haemolytic Complement Activity at Diagnosis as an Indicator of the Baseline Activity of Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , 2021, 28, 85-93.	1.1	1
35	Clinical features of Korean elderly patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 731-741.	1.7	3
36	Novel mortality-predicting index at diagnosis can effectively predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e23885.	2.1	4

#	ARTICLE	IF	CITATIONS
37	Clinical significance of large unstained cell count in estimating the current activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021, 75, e14512.	1.7	3
38	Predictive Ability of Serum IL-27 Level for Assessing Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Mediators of Inflammation</i> , 2021, 2021, 1-8.	3.0	2
39	Serum adipokine profiles in patients with microscopic polyangiitis and granulomatosis with polyangiitis: An exploratory analysis. <i>PLoS ONE</i> , 2021, 16, e0254226.	2.5	1
40	Efficacy of tacrolimus as maintenance therapy after cyclophosphamide for treating antineutrophil cytoplasmic antibody-associated vasculitis. <i>Medicine (United States)</i> , 2021, 100, e26956.	1.0	0
41	Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 1221-1232.	1.7	1
42	The novel fibrosis index at diagnosis may predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis without substantial liver diseases. <i>Clinics</i> , 2021, 76, e2501.	1.5	2
43	Antineutrophil Cytoplasmic Antibody Positivity Is Associated with Vascular Involvement in Behçet's Disease. <i>Yonsei Medical Journal</i> , 2021, 62, 149.	2.2	6
44	Male Sex Is a Significant Predictor of All-cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Korean Medical Science</i> , 2021, 36, e120.	2.5	8
45	The Efficacy of Mycophenolate Mofetil in Remission Maintenance Therapy for Microscopic Polyangiitis and Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2021, 62, 494.	2.2	2
46	Efficacy of the fibrosis index for predicting end-stage renal disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021, 75, e13929.	1.7	4
47	Serum Clusterin Level Could Reflect the Current Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2021, 62, 1016.	2.2	3
48	Serum progranulin as a predictive marker for high activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e24048.	2.1	1
49	Prevalence of Osteopenia in Drug-Naive Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Journal of Clinical Rheumatology</i> , 2021, 27, e330-e335.	0.9	2
50	Serum granzyme B is associated with otorhinolaryngological, pulmonary, and renal involvement of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Investigative Medicine</i> , 2021, 69, 91-95.	1.6	0
51	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39 Suppl 129, 88-93.	0.8	1
52	D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39 Suppl 129, 94-100.	0.8	0
53	Serum galectin-9 could be a potential biomarker in assessing the disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, . .	0.8	0
54	Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. <i>Clinical and Experimental Rheumatology</i> , 2021, 39 Suppl 131, 111-118.	0.8	0

#	ARTICLE	IF	CITATIONS
55	D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 94-100.	0.8	3
56	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 88-93.	0.8	14
57	Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 111-118.	0.8	0
58	Clinical impact of proteinase 3-antineutrophil cytoplasmic antibody positivity in eosinophilic granulomatosis with polyangiitis. <i>Korean Journal of Internal Medicine</i> , 2021, , .	1.7	1
59	Metabolic Syndrome Severity Score, Comparable to Serum Creatinine, Could Predict the Occurrence of End-Stage Kidney Disease in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Journal of Clinical Medicine</i> , 2021, 10, 5744.	2.4	5
60	Fibrosis-4 index at diagnosis can predict all-cause mortality in patients with rheumatoid arthritis: A retrospective monocentric study. <i>Modern Rheumatology</i> , 2020, 30, 70-77.	1.8	17
61	Multivariable index for assessing the activity and predicting all-cause mortality in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2020, 34, e23022.	2.1	13
62	Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjögren's syndrome?. <i>Rheumatology International</i> , 2020, 40, 1063-1070.	3.0	4
63	Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , 2020, 74, e13431.	1.7	2
64	Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , 2020, 39, 831-840.	2.2	2
65	Punicalagin Ameliorates Lupus Nephritis via Inhibition of PAR2. <i>International Journal of Molecular Sciences</i> , 2020, 21, 4975.	4.1	14
66	Incidence of Tuberculosis in Systemic Necrotizing Vasculitides: A Population-Based Study From an Intermediate-Burden Country. <i>Frontiers in Medicine</i> , 2020, 7, 550004.	2.6	0
67	Atherogenic index of plasma predicts cerebrovascular accident occurrence in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Lipids in Health and Disease</i> , 2020, 19, 184.	3.0	7
68	Serum Amyloid A Is a Biomarker of Disease Activity and Health-Related Quality-of-Life in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Disease Markers</i> , 2020, 2020, 1-9.	1.3	4
69	Definite IgG4-related disease had no overlap with eosinophilic granulomatosis with polyangiitis in Korean patients: a pilot study in one centre. <i>Clinical Rheumatology</i> , 2020, 39, 3009-3015.	2.2	3
70	Association between the antineutrophil cytoplasmic antibody and late coronary arterial occlusive disease in patients with Takayasu arteritis. <i>Journal of Cardiology</i> , 2020, 76, 407-412.	1.9	3
71	Clinical implication of plasma exchange on life-threatening antineutrophil cytoplasmic antibody-associated vasculitis. <i>BMC Pulmonary Medicine</i> , 2020, 20, 147.	2.0	2
72	Clinical characteristics and long-term outcomes of Libman's Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020, 29, 1115-1120.	1.6	11

#	ARTICLE	IF	CITATIONS
73	Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , 2020, 22, 138.	3.5	5
74	Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , 2020, 21, 258.	1.8	5
75	Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , 2020, 40, 1089-1099.	3.0	7
76	Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). <i>Seminars in Arthritis and Rheumatism</i> , 2020, 50, 1414-1420.	3.4	3
77	Serum interleukin-16 significantly correlates with the Vasculitis Damage Index in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Arthritis Research and Therapy</i> , 2020, 22, 73.	3.5	6
78	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 251, 117-123.	1.2	7
79	Worse Renal Presentation and Prognosis in Initial-Onset Lupus Nephritis than Early-Onset Lupus Nephritis. <i>Yonsei Medical Journal</i> , 2020, 61, 951.	2.2	4
80	Rituximab Biosimilar Prevents Poor Outcomes of Microscopic Polyangiitis and Granulomatosis with Polyangiitis as Effectively as Rituximab Originator. <i>Yonsei Medical Journal</i> , 2020, 61, 712.	2.2	5
81	Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor 1 Is Inversely Correlated with the Activity of ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2020, 61, 720.	2.2	2
82	Lipid Profiles in Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis: A Cross-sectional Analysis. <i>Journal of Rheumatic Diseases</i> , 2020, 27, 261-269.	1.1	4
83	Pregnancy Morbidities in Korean Patients with Takayasu Arteritis: A Monocentric Pilot Study. <i>Yonsei Medical Journal</i> , 2020, 61, 970.	2.2	3
84	Will the HALP score help to assess the activity and predict the prognosis of antineutrophil cytoplasmic antibody-associated vasculitis?. <i>Clinical and Experimental Rheumatology</i> , 2020, 38 Suppl 124, 236-237.	0.8	2
85	Clinical significance of ANCA positivity in patients with IgA vasculitis: a retrospective monocentric study. <i>Rheumatology International</i> , 2019, 39, 1927-1936.	3.0	4
86	Atializumab, a humanized anti-aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1) antibody significantly improves nephritis in (NZB/NZW) F1 mice. <i>Biomaterials</i> , 2019, 220, 119408.	11.4	7
87	Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2019, 38, 3485-3491.	2.2	7
88	Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , 2019, 14, e0219970.	2.5	8
89	Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naïve patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , 2019, 39, 1899-1905.	3.0	3
90	Serum Aminoacyl-tRNA Synthetase-Interacting Multifunctional Protein-1 Can Predict Severe Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Pilot Monocentric Study. <i>BioMed Research International</i> , 2019, 2019, 1-6.	1.9	7

#	ARTICLE	IF	CITATIONS
91	Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , 2019, 11, 1456.	4.1	14
92	Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010–2018. <i>Journal of Clinical Medicine</i> , 2019, 8, 1871.	2.4	10
93	Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , 2019, 39, 1859-1866.	3.0	4
94	Classification of Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , 2019, 26, 156.	1.1	11
95	Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , 2019, 39, 1937-1944.	3.0	26
96	Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , 2019, 19, 90.	2.0	13
97	Anti-phospholipid antibody syndrome occurrence in patients with persistent anti-phospholipid antibodies. <i>Rheumatology International</i> , 2019, 39, 1359-1367.	3.0	4
98	Thyroid Dysfunction in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis: A Monocentric Retrospective Study. <i>Journal of Rheumatology</i> , 2019, 46, 1248-1250.	2.0	2
99	Comparison of Radiological and Histological Findings of Lung Parenchyma in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 454.	2.2	10
100	Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , 2019, 39, 885-892.	3.0	3
101	Ex Vivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , 2019, 20, e299-e308.	2.6	2
102	Serum interleukin-21 positivity could indicate the current activity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical Rheumatology</i> , 2019, 38, 1685-1690.	2.2	5
103	Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 797-804.	1.9	16
104	ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , 2019, 39, 879-884.	3.0	9
105	Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , 2019, 39, 395-401.	0.4	6
106	Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 487-494.	2.2	11
107	Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 223-230.	1.6	20
108	Systemic immune-inflammation index could estimate the cross-sectional high activity and the poor outcomes in immunosuppressive drug-naïve patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Nephrology</i> , 2019, 24, 711-717.	1.6	42

#	ARTICLE	IF	CITATIONS
109	No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 1147-1154.	2.2	21
110	Antineutrophil Cytoplasmic Antibody-Associated Vasculitis in Korea: A Narrative Review. <i>Yonsei Medical Journal</i> , 2019, 60, 10.	2.2	27
111	Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 1164.	2.2	8
112	Eosinophilic Granulomatosis with Polyangiitis: Experiences in Korean Patients. <i>Yonsei Medical Journal</i> , 2019, 60, 705.	2.2	11
113	Subclinical but significant liver fibrosis in patients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 117, 26-31.	0.8	1
114	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 117, 116-121.	0.8	2
115	Intranuclear delivery of the transcription modulation domain of Tbet-improved lupus nephritis in (NZB/NZW) F1 lupus-prone mice. <i>Kidney International</i> , 2018, 93, 1118-1130.	5.2	9
116	Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , 2018, 37, 2133-2141.	2.2	12
117	Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , 2018, 38, 1095-1101.	3.0	9
118	Increased serum interleukin-32 levels in patients with Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 2167-2174.	1.9	20
119	Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010-2011). <i>Clinical Rheumatology</i> , 2018, 37, 1333-1340.	2.2	15
120	Rheumatoid factor false positivity in patients with ANCA-associated vasculitis not having medical conditions producing rheumatoid factor. <i>Clinical Rheumatology</i> , 2018, 37, 2771-2779.	2.2	10
121	Hemoglobin A1c, Not Glycated Albumin, Can Independently Reflect the Ankylosing Spondylitis Disease Activity Score. <i>Journal of Rheumatic Diseases</i> , 2018, 25, 131.	1.1	2
122	Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide Korean College of Rheumatology BIOlogics (KOBIO) registry. <i>PLoS ONE</i> , 2018, 13, e0205651.	2.5	13
123	Application of 2012 EULAR/ACR criteria for polymyalgia rheumatica to Korean patients previously classified by Chuang and Hunder criteria or Healey criteria. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 1838-1843.	1.9	2
124	HBsAg-negative and anti-HBc-positive in eosinophilic granulomatosis with polyangiitis: a retrospective pilot study. <i>Rheumatology International</i> , 2018, 38, 1531-1538.	3.0	5
125	Red Blood Cell Distribution Width Can Predict Vasculitis Activity and Poor Prognosis in Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2018, 59, 294.	2.2	6
126	Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , 2018, 19, 187.	1.8	32

#	ARTICLE	IF	CITATIONS
127	Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , 2018, 59, 452.	2.2	27
128	Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018, 59, 397.	2.2	16
129	Extravascular manifestations of Takayasu arteritis: focusing on the features shared with spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2018, 20, 142.	3.5	20
130	Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , 2018, 38, 1865-1871.	3.0	28
131	C-Reactive Protein to Serum Albumin Ratio Is an Independent Predictor of All-Cause Mortality in Patients with ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018, 59, 865.	2.2	36
132	Serum leucine-rich Î±2-glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , 2018, 486, 253-258.	1.1	10
133	Red blood cell distribution width is useful in discriminating adult onset Still's disease and sepsis within 24 hours after hospitalization. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 1234-1240.	1.7	8
134	Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 165-166.	0.8	2
135	The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 85-87.	0.8	14
136	Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 115-120.	0.8	2
137	Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , 2018, 36, 533-539.	0.8	2
138	Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 73-77.	0.8	3
139	The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 65-72.	0.8	9
140	Glycated albumin increases with disease activity in rheumatoid factor positive rheumatoid arthritis patients with normal fasting glucose and HbA1c. <i>Joint Bone Spine</i> , 2017, 84, 115-118.	1.6	5
141	Serum anti-lysozyme is associated with disease activity of Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 261-268.	1.9	8
142	Five factor score of more than 1 is associated with relapse during the first 2 year follow up in patients with eosinophilic granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1261-1268.	1.9	12
143	In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , 2017, 47, 216-221.	3.4	28
144	Evaluation of Spleen Glucose Metabolism Using ¹⁸ F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , 2017, 58, 507-513.	5.0	33

#	ARTICLE	IF	CITATIONS
145	Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2017, 44, 996-1003.	2.0	43
146	Delta neutrophil index contributes to the differential diagnosis between acute gout attack and cellulitis within 24 hours after hospitalization. <i>Rheumatology</i> , 2017, 56, kew471.	1.9	11
147	Malignancies in Korean patients with immunoglobulin G4-related disease. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1028-1035.	1.9	35
148	Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , 2017, 37, 685-694.	3.0	11
149	Echocardiographic features in patients with ANCA-associated vasculitis within 3 months before and after diagnosis. <i>Clinical Rheumatology</i> , 2017, 36, 2751-2759.	2.2	9
150	Cell-penetrating interactomic inhibition of nuclear factor-kappa B in a mouse model of postoperative cognitive dysfunction. <i>Scientific Reports</i> , 2017, 7, 13482.	3.3	27
151	The clinical utility of splenic fluorodeoxyglucose uptake for diagnosis and prognosis in patients with macrophage activation syndrome. <i>Medicine (United States)</i> , 2017, 96, e7901.	1.0	6
152	Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1593-1605.	1.9	17
153	Clinical significance of delta neutrophil index in the differential diagnosis between septic arthritis and acute gout attack within 24 hours after hospitalization. <i>Medicine (United States)</i> , 2017, 96, e7431.	1.0	7
154	Defective autophagy activity and its association with spinal damage in patients with ankylosing spondylitis. <i>Joint Bone Spine</i> , 2017, 84, 583-587.	1.6	12
155	Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , 2017, 12, e0184323.	2.5	5
156	Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , 2017, 19, 193.	3.5	14
157	Platelet Distribution Width and Mean Platelet Volume Are Not Correlated with the Disease Activity Indices of Ankylosing Spondylitis. <i>Journal of Rheumatic Diseases</i> , 2017, 24, 143.	1.1	0
158	Subsequent Thrombotic Outcomes in Patients with Ischemic Stroke with Antiphospholipid Antibody Positivity. <i>Yonsei Medical Journal</i> , 2017, 58, 1128.	2.2	6
159	Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , 2017, 12, e0181590.	2.5	9
160	Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 47-54.	0.8	11
161	Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 111-118.	0.8	16
162	The Presence of Anti-ribonucleoprotein at Diagnosis Is Associated with the Flare during the First Follow-up Year in Korean Patients with Systemic Lupus Erythematosus. <i>Journal of Rheumatic Diseases</i> , 2016, 23, 154.	1.1	1

#	ARTICLE	IF	CITATIONS
163	Anti- Sm is associated with the early poor outcome of lupus nephritis. <i>International Journal of Rheumatic Diseases</i> , 2016, 19, 897-902.	1.9	22
164	Clinical predictors of silent but substantial liver fibrosis in primary Sjogren's syndrome. <i>Modern Rheumatology</i> , 2016, 26, 576-582.	1.8	6
165	Treatment of Collagen-Induced Arthritis Using Immune Modulatory Properties of Human Mesenchymal Stem Cells. <i>Cell Transplantation</i> , 2016, 25, 1057-1072.	2.5	39
166	Brain meningioma in a patient with systemic lupus erythematosus. <i>Yeungnam University Journal of Medicine</i> , 2016, 33, 159.	1.4	1
167	Red nail folds and Gottron's sign in a patient with leukemic infiltration of pelvic muscles. <i>JDDG - Journal of the German Society of Dermatology</i> , 2015, 13, 581-582.	0.8	0
168	Hepatic Sarcoidosis in a Patient with Chronic Hepatitis B Virus Infection. <i>Journal of Rheumatic Diseases</i> , 2015, 22, 200.	1.1	0
169	The antibody atliximab attenuates collagen-induced arthritis by neutralizing AIMP1, an inflammatory cytokine that enhances osteoclastogenesis. <i>Biomaterials</i> , 2015, 44, 45-54.	11.4	16
170	Application of the 2013 ACR/EULAR classification criteria for systemic sclerosis to patients with Raynaud's phenomenon. <i>Arthritis Research and Therapy</i> , 2015, 17, 77.	3.5	20
171	Dermatomyositis Associated with Gallbladder Cancer. <i>Journal of Rheumatic Diseases</i> , 2014, 21, 261.	1.1	0
172	Delta Neutrophil Index as an Early Marker for Differential Diagnosis of Adult-Onset Still's Disease and Sepsis. <i>Yonsei Medical Journal</i> , 2014, 55, 753.	2.2	26
173	Differences in Clinical Manifestations and Outcomes between Adult and Child Patients with Henoch-Schönlein Purpura. <i>Journal of Korean Medical Science</i> , 2014, 29, 198.	2.5	109
174	Usefulness of Serum Leucine-Rich Alpha-2 Glycoprotein as a Disease Activity Biomarker in Patients with Rheumatoid Arthritis. <i>Journal of Korean Medical Science</i> , 2014, 29, 1199.	2.5	37
175	ROR γ -specific transcriptional interactomic inhibition suppresses autoimmunity associated with T _H 17 cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 18673-18678.	7.1	33
176	Usefulness of FDG PET/CT in the Diagnosis of Eosinophilic Fasciitis. <i>Clinical Nuclear Medicine</i> , 2014, 39, 801-802.	1.3	18
177	Active Tuberculosis Risk With Tumor Necrosis Factor Inhibitors After Treating Latent Tuberculosis. <i>Journal of Clinical Rheumatology</i> , 2014, 20, 68-73.	0.9	12
178	Inhibition of glycogen synthase kinase-3 β suppresses inflammatory responses in rheumatoid arthritis fibroblast-like synoviocytes and collagen-induced arthritis. <i>Joint Bone Spine</i> , 2014, 81, 240-246.	1.6	29
179	The concomitant use of meloxicam and methotrexate does not clearly increase the risk of silent kidney and liver damages in patients with rheumatoid arthritis. <i>Rheumatology International</i> , 2014, 34, 833-840.	3.0	4
180	Concurrent use of methotrexate and celecoxib increases risk of silent liver fibrosis in rheumatoid arthritis patients with subclinical reduced kidney function. <i>Clinical Rheumatology</i> , 2014, 33, 1415-1423.	2.2	9

#	ARTICLE	IF	CITATIONS
181	Secreted frizzled-related protein 5 suppresses inflammatory response in rheumatoid arthritis fibroblast-like synoviocytes through down-regulation of c-Jun N-terminal kinase. <i>Rheumatology</i> , 2014, 53, 1704-1711.	1.9	24
182	Activation of Liver X Receptors Suppresses Inflammatory Gene Expressions and Transcriptional Corepressor Clearance in Rheumatoid Arthritis Fibroblast Like Synoviocytes. <i>Journal of Clinical Immunology</i> , 2013, 33, 190-199.	3.8	11
183	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2013, 23, 509-515.	1.8	4
184	Gallic acid, a natural polyphenolic acid, induces apoptosis and inhibits proinflammatory gene expressions in rheumatoid arthritis fibroblast-like synoviocytes. <i>Joint Bone Spine</i> , 2013, 80, 274-279.	1.6	116
185	Osteoporotic Calcaneal Stress Fractures Mistaken for Aggravation of Rheumatoid Arthritis. <i>Arthritis and Rheumatism</i> , 2013, 65, 2983-2983.	6.7	6
186	Soluble Receptor for Advanced Glycation End Products Alleviates Nephritis in (NZB/NZW)F1 Mice. <i>Arthritis and Rheumatism</i> , 2013, 65, 1902-1912.	6.7	23
187	Adipokines, inflammation, insulin resistance, and carotid atherosclerosis in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2013, 15, R194.	3.5	36
188	Typical 18-FDG-PET/CT Findings of Polymyalgia Rheumatica: A Case Report. <i>Journal of Rheumatic Diseases</i> , 2013, 20, 113.	1.1	4
189	Two Cases of Refractory Thrombocytopenia in Systemic Lupus Erythematosus that Responded to Intravenous Low-Dose Cyclophosphamide. <i>Journal of Korean Medical Science</i> , 2013, 28, 472.	2.5	8
190	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2013, 23, 509-515.	1.8	2
191	Attenuation of nephritis in lupus-prone mice by thalidomide. <i>Rheumatology</i> , 2012, 51, 2131-2140.	1.9	12
192	Patients with early arthritis who fulfil the 1987 ACR classification criteria for rheumatoid arthritis but not the 2010 ACR/EULAR criteria. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 1097-1098.	0.9	16
193	Leflunomide increases the risk of silent liver fibrosis in patients with rheumatoid arthritis receiving methotrexate. <i>Arthritis Research and Therapy</i> , 2012, 14, R232.	3.5	45
194	Alleviation of rheumatoid arthritis by cell-transducible methotrexate upon transcutaneous delivery. <i>Biomaterials</i> , 2012, 33, 1563-1572.	11.4	29
195	The role of ¹⁸ F-fluorodeoxyglucose positron emission tomography in the assessment of disease activity in patients with Takayasu arteritis. <i>Arthritis and Rheumatism</i> , 2012, 64, 866-875.	6.7	88
196	The prevalence and the clinical relevance of anti-Ro52 in Korean patients with primary Sjögren's syndrome. <i>Rheumatology International</i> , 2012, 32, 491-495.	3.0	19
197	A Case of Systemic Lupus Erythematosus Manifested as Anemia and Thrombocytopenia during Antithyroid Drug Treatment. <i>Korean Journal of Medicine</i> , 2012, 82, 232.	0.3	0
198	Retroperitoneal Fibrosis in 27 Korean Patients: Single Center Experience. <i>Journal of Korean Medical Science</i> , 2011, 26, 985.	2.5	19

#	ARTICLE	IF	CITATIONS
199	A Case of Ankylosing Spondylitis with Follicular Lymphoma. <i>Journal of Rheumatic Diseases</i> , 2011, 18, 315.	1.1	0
200	Clinical Manifestations of Korean Adult Patients with Henoch-Schönlein Purpura. <i>The Journal of the Korean Rheumatism Association</i> , 2010, 17, 133.	0.1	3
201	Inflammatory Polyarthritis in a Patient with Psoriasis: Is It Psoriatic Arthritis or Rheumatoid Arthritis?. <i>Korean Journal of Internal Medicine</i> , 2010, 25, 224.	1.7	17
202	Comparison of therapeutic efficacy between bortezomib and combination treatment of prednisolone and mycophenolate mofetil on nephritis in NZB/WF1 mice. <i>Clinical and Experimental Rheumatology</i> , 2010, 28, 393-6.	0.8	7
203	E/E' ratio is more sensitive than E/A ratio for detection of left ventricular diastolic dysfunction in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2010, 28, S12-7.	0.8	17
204	The Mid-Range of the Adjusted Level of Ferritin Can Predict the Chronic Course in Patients with Adult Onset Still's Disease. <i>Journal of Rheumatology</i> , 2009, 36, 156-162.	2.0	30
205	Kikuchi-Fujimoto's Disease with Adult Onset Still's Disease. <i>The Journal of the Korean Rheumatism Association</i> , 2008, 15, 58.	0.1	0
206	Measurement of the serum leptin level could assist disease activity monitoring in rheumatoid arthritis. <i>Rheumatology International</i> , 2007, 27, 537-540.	3.0	89
207	Malignancies in Korean Patients with Inflammatory Myopathy. <i>Yonsei Medical Journal</i> , 2006, 47, 519.	2.2	34
208	Serum galectin-9 could be a potential biomarker in assessing the disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.8	1
209	Effect of numbers of metabolic syndrome components on mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis with metabolic syndrome. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.8	1
210	Application of the 2022 ACR/EULAR criteria for microscopic polyangiitis to patients with previously diagnosed microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.8	2
211	Fatty Liver Index Independently Predicts All-Cause Mortality in Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis but No Substantial Liver Disease. <i>Frontiers in Cardiovascular Medicine</i> , 0, 9, .	2.4	3
212	Predicting the depressive status using empirical dietary inflammatory index in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 0, , .	2.1	1
213	BVAS version 3 and BVAS/GPA: standing on the same line?. <i>Clinical Rheumatology</i> , 0, , .	2.2	1