

# Sten Fredrikson

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6244001/publications.pdf>

Version: 2024-02-01

158  
papers

6,480  
citations

61984

43  
h-index

79698

73  
g-index

164  
all docs

164  
docs citations

164  
times ranked

6300  
citing authors

#	ARTICLE	IF	CITATIONS
1	Picornavirus May Be Linked to Parkinson's Disease through Viral Antigen in Dopamine-Containing Neurons of Substantia Nigra. <i>Microorganisms</i> , 2022, 10, 599.	3.6	2
2	Narcolepsy among first- and second-generation immigrants in Sweden: A study of the total population. <i>Acta Neurologica Scandinavica</i> , 2022, 146, 160-166.	2.1	2
3	Telephone validation of an Urdu translated version of the extended disability severity scale in multiple sclerosis patients. <i>Multiple Sclerosis and Related Disorders</i> , 2021, 48, 102684.	2.0	6
4	Associations Between Fluctuations in Daytime Sleepiness and Motor and Non-Motor Symptoms in Parkinson's Disease. <i>Movement Disorders Clinical Practice</i> , 2021, 8, 44-50.	1.5	7
5	Huntington's disease among immigrant groups and Swedish-born individuals: a cohort study of all adults 18 years of age and older in Sweden. <i>Neurological Sciences</i> , 2021, 42, 3851-3856.	1.9	1
6	Amyotrophic lateral sclerosis (ALS) among immigrant groups and Swedish-born individuals: a cohort study of all adults 18 years of age and older in Sweden. <i>Journal of Neurology</i> , 2021, , 1.	3.6	0
7	Multiple sclerosis among first- and second-generation immigrant groups in Sweden. <i>Acta Neurologica Scandinavica</i> , 2020, 142, 339-349.	2.1	5
8	Multiple sclerosis in Pakistan: Current status and future perspective. <i>Journal of the Neurological Sciences</i> , 2020, 418, 117066.	0.6	8
9	Parkinson's Disease Among Immigrant Groups and Swedish-Born Individuals: A Cohort Study of All Adults 50 Years of Age and Older in Sweden. <i>Journal of Parkinson's Disease</i> , 2020, 10, 1133-1141.	2.8	4
10	Epilepsy in immigrants and Swedish-born: A cohort study of all adults 18 years of age and older in Sweden. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2020, 76, 116-122.	2.0	1
11	Validation of Rapid Magnetic Resonance Myelin Imaging in Multiple Sclerosis. <i>Annals of Neurology</i> , 2020, 87, 710-724.	5.3	42
12	RebiQoL: A randomized trial of telemedicine patient support program for health-related quality of life and adherence in people with MS treated with Rebif. <i>PLoS ONE</i> , 2019, 14, e0218453.	2.5	11
13	Gadolinium Retention in the Brain: An MRI Relaxometry Study of Linear and Macrocyclic Gadolinium-Based Contrast Agents in Multiple Sclerosis. <i>American Journal of Neuroradiology</i> , 2019, 40, 1265-1273.	2.4	24
14	A 10-Year Follow-Up of Excessive Daytime Sleepiness in Parkinson's Disease. <i>Parkinson's Disease</i> , 2019, 2019, 1-7.	1.1	10
15	Lesion accumulation is predictive of long-term cognitive decline in multiple sclerosis. <i>Multiple Sclerosis and Related Disorders</i> , 2018, 21, 110-116.	2.0	20
16	Survey of diagnostic and treatment practices for multiple sclerosis (MS) in Europe. Part 2: Progressive MS, paediatric MS, pregnancy and general management. <i>European Journal of Neurology</i> , 2018, 25, 739-746.	3.3	12
17	Detection of Leukocortical Lesions in Multiple Sclerosis and Their Association with Physical and Cognitive Impairment: A Comparison of Conventional and Synthetic Phase-Sensitive Inversion Recovery MRI. <i>American Journal of Neuroradiology</i> , 2018, 39, 1995-2000.	2.4	17
18	Survey of diagnostic and treatment practices for multiple sclerosis in Europe. <i>European Journal of Neurology</i> , 2017, 24, 516-522.	3.3	34

#	ARTICLE	IF	CITATIONS
19	Treatment with disease-modifying drugs for people with a first clinical attack suggestive of multiple sclerosis. <i>The Cochrane Library</i> , 2017, 4, CD012200.	2.8	20
20	Retention of Gadolinium-Based Contrast Agents in Multiple Sclerosis: Retrospective Analysis of an 18-Year Longitudinal Study. <i>American Journal of Neuroradiology</i> , 2017, 38, 1311-1316.	2.4	48
21	Alemtuzumab Use in Clinical Practice: Recommendations from European Multiple Sclerosis Experts. <i>CNS Drugs</i> , 2017, 31, 33-50.	5.9	57
22	Clinical Feasibility of Synthetic MRI in Multiple Sclerosis: A Diagnostic and Volumetric Validation Study. <i>American Journal of Neuroradiology</i> , 2016, 37, 1023-1029.	2.4	104
23	A single-group pilot feasibility study of cognitive behavioural therapy in people with multiple sclerosis with depressive symptoms. <i>Disability and Rehabilitation</i> , 2016, 38, 2383-2391.	1.8	7
24	Incidence of Radiologically Isolated Syndrome: A Population-Based Study. <i>American Journal of Neuroradiology</i> , 2016, 37, 1017-1022.	2.4	40
25	A 10-year follow-up of the European multicenter trial of interferon $\beta$ -1b in secondary-progressive multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2016, 22, 533-543.	3.0	24
26	Is excessive daytime sleepiness a separate manifestation in Parkinson's disease?. <i>Acta Neurologica Scandinavica</i> , 2015, 132, 97-104.	2.1	20
27	MRI-defined Corpus Callosal Atrophy in Multiple Sclerosis: A Comparison of Volumetric Measurements, Corpus Callosum Area and Index. <i>Journal of Neuroimaging</i> , 2015, 25, 996-1001.	2.0	40
28	Corpus callosum atrophy is strongly associated with cognitive impairment in multiple sclerosis: Results of a 17-year longitudinal study. <i>Multiple Sclerosis Journal</i> , 2015, 21, 1151-1158.	3.0	63
29	Autologous haematopoietic stem cell transplantation for aggressive multiple sclerosis: the Swedish experience. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2014, 85, 1116-1121.	1.9	139
30	Health-related quality of life in partners of persons with MS: a longitudinal 10-year perspective. <i>BMJ Open</i> , 2014, 4, e006097.	1.9	5
31	A 10-year follow-up of a population-based study of people with multiple sclerosis in Stockholm, Sweden: Changes in disability and the value of different factors in predicting disability and mortality. <i>Journal of the Neurological Sciences</i> , 2013, 332, 121-127.	0.6	40
32	Reports of Patients and Relatives from the CogniCIS Study about Cognition in Clinically Isolated Syndrome: What Are Our Patients Telling Us?. <i>European Neurology</i> , 2013, 69, 346-351.	1.4	5
33	Radiologically isolated syndrome " incidental magnetic resonance imaging findings suggestive of multiple sclerosis, a systematic review. <i>Multiple Sclerosis Journal</i> , 2013, 19, 271-280.	3.0	116
34	Callosal atrophy in multiple sclerosis is related to cognitive speed. <i>Acta Neurologica Scandinavica</i> , 2013, 127, 281-289.	2.1	28
35	People with multiple sclerosis in Denmark who use complementary and alternative medicine "Do subgroups of patients differ?. <i>European Journal of Integrative Medicine</i> , 2013, 5, 365-373.	1.7	4
36	The utility of cerebrospinal fluid analysis in patients with multiple sclerosis. <i>Nature Reviews Neurology</i> , 2013, 9, 267-276.	10.1	181

#	ARTICLE	IF	CITATIONS
37	A cost-effectiveness analysis of subcutaneous interferon beta-1a 44mcg 3-times a week vs no treatment for patients with clinically isolated syndrome in Sweden. <i>Journal of Medical Economics</i> , 2013, 16, 756-762.	2.1	5
38	Interferon $\beta$ for secondary progressive multiple sclerosis: a systematic review. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2013, 84, 420-426.	1.9	47
39	Radiologically isolated syndrome: an uncommon finding at a university clinic in a high-prevalence region for multiple sclerosis. <i>BMJ Open</i> , 2013, 3, e003531.	1.9	13
40	Time to secondary progression in patients with multiple sclerosis who were treated with first generation immunomodulating drugs. <i>Multiple Sclerosis Journal</i> , 2013, 19, 765-774.	3.0	66
41	Differences between users and non-users of complementary and alternative medicine among people with multiple sclerosis in Denmark: A comparison of descriptive characteristics. <i>Scandinavian Journal of Public Health</i> , 2013, 41, 492-499.	2.3	23
42	Multiple sclerosis in Pakistan: histocompatibility antigen composition and disability. <i>Multiple Sclerosis Journal</i> , 2013, 19, 254-255.	3.0	5
43	Cost-minimization analysis of fingolimod compared with natalizumab for the treatment of relapsing-remitting multiple sclerosis in Sweden. <i>Journal of Medical Economics</i> , 2013, 16, 349-357.	2.1	8
44	Emerging Oral Medications for Multiple Sclerosis. , 2012, , .		0
45	Recommendations for a Brief International Cognitive Assessment for Multiple Sclerosis (BICAMS). <i>Multiple Sclerosis Journal</i> , 2012, 18, 891-898.	3.0	654
46	Altered cerebrospinal fluid index of prealbumin, fibrinogen, and haptoglobin in patients with Guillain-Barré syndrome and chronic inflammatory demyelinating polyneuropathy. <i>Acta Neurologica Scandinavica</i> , 2012, 125, 129-135.	2.1	27
47	Cost-Effectiveness Analysis of Interferon Beta-1b for the Treatment of Patients With a First Clinical Event Suggestive of Multiple Sclerosis. <i>Clinical Therapeutics</i> , 2012, 34, 1132-1144.	2.5	13
48	A longitudinal observational study of brain atrophy rate reflecting four decades of multiple sclerosis: a comparison of serial 1D, 2D, and volumetric measurements from MRI images. <i>Neuroradiology</i> , 2010, 52, 109-117.	2.2	28
49	The first case history of multiple sclerosis: Augustus Est (1794-1848). <i>Neurological Sciences</i> , 2010, 31, 29-33.	1.9	19
50	Evaluation of multiple sclerosis diagnostic criteria in Suzhou, China - risk of under-diagnosis in a low prevalence area. <i>Acta Neurologica Scandinavica</i> , 2010, 121, 24-29.	2.1	9
51	Health-related quality of life in relapsing remitting multiple sclerosis patients during treatment with glatiramer acetate: a prospective, observational, international, multi-centre study. <i>Health and Quality of Life Outcomes</i> , 2010, 8, 133.	2.4	44
52	"We noticed that suddenly the country has become full of MRI". Policy makers' views on diffusion and use of health technologies in Iran. <i>Health Research Policy and Systems</i> , 2010, 8, 9.	2.8	24
53	CSF immune variables in patients with narcolepsy. <i>Acta Neurologica Scandinavica</i> , 2009, 81, 253-254.	2.1	57
54	Lessons from randomised direct comparative trials. <i>Journal of the Neurological Sciences</i> , 2009, 277, S19-S24.	0.6	11

#	ARTICLE	IF	CITATIONS
55	Reduced cerebrospinal fluid BACE1 activity in multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2009, 15, 448-454.	3.0	55
56	Diffusion of interferon beta in Iran and its utilization in Tehran. <i>Pharmacoepidemiology and Drug Safety</i> , 2008, 17, 934-941.	1.9	8
57	Cord blood contains cells secreting antibodies to nervous system components. <i>Clinical and Experimental Immunology</i> , 2008, 84, 353-358.	2.6	7
58	The expression of TNF- $\alpha$ receptors 1 and 2 on peripheral blood mononuclear cells in chronic inflammatory demyelinating polyneuropathy. <i>Journal of Neuroimmunology</i> , 2008, 200, 129-132.	2.3	6
59	Use of health care services and satisfaction with care in people with multiple sclerosis in Stockholm County: A population-based study. <i>Multiple Sclerosis Journal</i> , 2008, 14, 962-971.	3.0	40
60	How is magnetic resonance imaging used in Iran?. <i>International Journal of Technology Assessment in Health Care</i> , 2008, 24, 452-458.	0.5	13
61	Clinical features of patients with multiple sclerosis from a survey in Shanghai, China. <i>Multiple Sclerosis Journal</i> , 2008, 14, 671-678.	3.0	28
62	Effects of inpatient rehabilitation in multiple sclerosis patients with moderate disability. <i>Advances in Physiotherapy</i> , 2008, 10, 58-65.	0.2	15
63	Diffusion of magnetic resonance imaging in Iran. <i>International Journal of Technology Assessment in Health Care</i> , 2007, 23, 278-285.	0.5	21
64	Multiple sclerosis in Pakistan. <i>Multiple Sclerosis Journal</i> , 2007, 13, 668-669.	3.0	21
65	Selective Decline in Information Processing in Subgroups of Multiple Sclerosis: An 8-Year Longitudinal Study. <i>European Neurology</i> , 2007, 57, 193-202.	1.4	119
66	A case-control study of Guillain-Barré syndrome in Harbin, China. <i>European Journal of Neurology</i> , 2006, 13, 953-957.	3.3	7
67	Costs and quality of life of multiple sclerosis in Sweden. <i>European Journal of Health Economics</i> , 2006, 7, 75-85.	2.8	63
68	Progression of non-age-related callosal brain atrophy in multiple sclerosis: a 9-year longitudinal MRI study representing four decades of disease development. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2006, 78, 375-380.	1.9	41
69	Cognitive and motor function in people with multiple sclerosis in Stockholm County. <i>Multiple Sclerosis Journal</i> , 2006, 12, 340-353.	3.0	52
70	Health-related quality of life in a population-based sample of people with multiple sclerosis in Stockholm County. <i>Multiple Sclerosis Journal</i> , 2006, 12, 605-612.	3.0	58
71	Activities of daily living and social activities in people with multiple sclerosis in Stockholm County. <i>Clinical Rehabilitation</i> , 2006, 20, 543-551.	2.2	88
72	Concordance for disease course and age of onset in Scandinavian multiple sclerosis coaffected sib pairs. <i>Multiple Sclerosis Journal</i> , 2004, 10, 5-8.	3.0	24

#	ARTICLE	IF	CITATIONS
73	Genes in the HLA class I region may contribute to the HLA class II-associated genetic susceptibility to multiple sclerosis. <i>Tissue Antigens</i> , 2004, 63, 237-247.	1.0	130
74	Cost-utility of interferon $\beta$ 1b in the treatment of patients with active relapsing-remitting or secondary progressive multiple sclerosis. <i>European Journal of Health Economics</i> , 2003, 4, 50-59.	2.8	39
75	Interferon- $\beta$ treatment in patients with multiple sclerosis does not alter CYP2C19 or CYP2D6 activity. <i>British Journal of Clinical Pharmacology</i> , 2003, 56, 337-340.	2.4	6
76	Prospective study of clinical epidemiology of Guillain-Barré syndrome in Harbin, China. <i>Journal of the Neurological Sciences</i> , 2003, 215, 63-69.	0.6	14
77	Multiple sclerosis in Stockholm County. A pilot study exploring the feasibility of assessment of impairment, disability and handicap by home visits. <i>Clinical Rehabilitation</i> , 2003, 17, 294-303.	2.2	30
78	Elevated Suicide Risk among Patients with Multiple Sclerosis in Sweden. <i>Neuroepidemiology</i> , 2003, 22, 146-152.	2.3	91
79	Altered phenotype and function of blood dendritic cells in multiple sclerosis are modulated by IFN- $\gamma$ and IL-10. <i>Clinical and Experimental Immunology</i> , 2002, 124, 306-314.	2.6	37
80	HHV-6 A- or B-specific P41 antigens do not reveal virus variant-specific IgG or IgM responses in human serum. <i>Journal of Medical Virology</i> , 2002, 66, 394-399.	5.0	19
81	Multiple sclerosis in Stockholm County. A pilot study of utilization of health-care resources, patient satisfaction with care and impact on family caregivers. <i>Acta Neurologica Scandinavica</i> , 2002, 106, 241-247.	2.1	16
82	A genome-wide screen for linkage in Nordic sib-pairs with multiple sclerosis. <i>Genes and Immunity</i> , 2002, 3, 279-285.	4.1	73
83	Distinct pattern of age-specific incidence of Guillain-Barré syndrome in Harbin, China. <i>Journal of Neurology</i> , 2002, 249, 25-32.	3.6	41
84	Costs, quality of life and disease severity in multiple sclerosis: a cross-sectional study in Sweden. <i>European Journal of Neurology</i> , 2001, 8, 27-35.	3.3	155
85	The T cell regulator gene SH2D2A contributes to the genetic susceptibility of multiple sclerosis. <i>Genes and Immunity</i> , 2001, 2, 263-268.	4.1	44
86	Linkage analysis suggests a region of importance for multiple sclerosis in 3p14-13. <i>Genes and Immunity</i> , 2001, 2, 451-454.	4.1	4
87	COST-UTILITY ANALYSIS OF INTERFERON BETA-1B IN SECONDARY PROGRESSIVE MULTIPLE SCLEROSIS. <i>International Journal of Technology Assessment in Health Care</i> , 2000, 16, 768-780.	0.5	35
88	Neutralizing and binding anti-interferon- $\beta$ (IFN- $\beta$ ) antibodies. A comparison between IFN- $\beta$ -1a and IFN- $\beta$ -1b treatment in multiple sclerosis. <i>European Journal of Neurology</i> , 2000, 7, 27-34.	3.3	67
89	Decreased mRNA expression of TNF-alpha and IL-10 in non-stimulated peripheral blood mononuclear cells in myasthenia gravis. <i>European Journal of Neurology</i> , 2000, 7, 195-202.	3.3	10
90	Clinical epidemiology of Guillain-Barré syndrome in adults in Sweden 1996-97: a prospective study. <i>European Journal of Neurology</i> , 2000, 7, 685-692.	3.3	43

#	ARTICLE	IF	CITATIONS
91	Epidemiological surveillance of Guillain-Barré syndrome in Sweden, 1996-1997. <i>Acta Neurologica Scandinavica</i> , 2000, 101, 104-111.	2.1	15
92	Linkage analysis of a candidate region in Scandinavian sib pairs with multiple sclerosis reveals linkage to chromosome 17q. <i>Genes and Immunity</i> , 2000, 1, 456-459.	4.1	29
93	Multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2000, 108, 236-243.	2.3	83
94	Multiple Sclerosis: Levels of Interleukin-10-Secreting Blood Mononuclear Cells are Low in Untreated Patients but Augmented During Interferon-beta-1b Treatment. <i>Scandinavian Journal of Immunology</i> , 1999, 49, 554-561.	2.7	67
95	High numbers of perforin mRNA expressing CSF cells in multiple sclerosis patients with gadolinium-enhancing brain MRI lesions. <i>Acta Neurologica Scandinavica</i> , 1999, 100, 18-24.	2.1	11
96	First hospital-admission rate as an epidemiological indicator for patients with multiple sclerosis in Stockholm, 1984-1993. <i>Acta Neurologica Scandinavica</i> , 1999, 100, 64-68.	2.1	5
97	Association and linkage analysis of candidate chromosomal regions in multiple sclerosis: indication of disease genes in 12q23 and 7p15. <i>European Journal of Human Genetics</i> , 1999, 7, 110-116.	2.8	23
98	Bilateral subdural haematomas following lumbar puncture in three haematopoietic stem cell transplant recipients. <i>Bone Marrow Transplantation</i> , 1999, 24, 1033-1035.	2.4	13
99	Multiple sclerosis is associated with high levels of circulating dendritic cells secreting pro-inflammatory cytokines. <i>Journal of Neuroimmunology</i> , 1999, 99, 82-90.	2.3	91
100	Linkage and association analysis of susceptibility regions on chromosomes 5 and 6 in 106 Scandinavian sibling pair families with multiple sclerosis. <i>Annals of Neurology</i> , 1999, 46, 612-616.	5.3	52
101	Linkage and association analysis of susceptibility regions on chromosomes 5 and 6 in 106 Scandinavian sibling pair families with multiple sclerosis. <i>Annals of Neurology</i> , 1999, 46, 612-616.	5.3	1
102	A New Cell Enzyme-Linked Immunosorbent Assay Demonstrates Gamma Interferon Suppression by Beta Interferon in Multiple Sclerosis. <i>Vaccine Journal</i> , 1999, 6, 415-419.	2.6	12
103	Similar Humoral and Cellular Immunological Reactivities to Human Herpesvirus 6 in Patients with Multiple Sclerosis and Controls. <i>Vaccine Journal</i> , 1999, 6, 545-549.	2.6	57
104	No evidence for increased frequency of autoantibodies during interferon- $\beta$ treatment of multiple sclerosis. <i>Acta Neurologica Scandinavica</i> , 1998, 97, 320-323.	2.1	23
105	The B-cell repertoire in myasthenia gravis includes all four acetylcholine receptor subunits. <i>Acta Neurologica Scandinavica</i> , 1998, 98, 422-426.	2.1	4
106	No evidence for elevated numbers of mononuclear cells expressing MCP-1 and RANTES mRNA in blood and CSF in multiple sclerosis. <i>Journal of Neuroimmunology</i> , 1998, 91, 108-112.	2.3	18
107	Influence of IFN-beta1b (Betaferon) on cytokine mRNA profiles in blood mononuclear cells and plasma levels of soluble VCAM-1 in multiple sclerosis. <i>European Journal of Neurology</i> , 1998, 5, 265-275.	3.3	21
108	Interleukin-12 and Perforin mRNA Expression is Augmented in Blood Mononuclear Cells in Multiple Sclerosis. <i>Scandinavian Journal of Immunology</i> , 1998, 47, 582-590.	2.7	32

#	ARTICLE	IF	CITATIONS
109	IL-15 mRNA expression is up-regulated in blood and cerebrospinal fluid mononuclear cells in multiple sclerosis (MS). <i>Clinical and Experimental Immunology</i> , 1998, 111, 193-197.	2.6	60
110	Linkage and association analysis of genes encoding cytokines and myelin proteins in multiple sclerosis. <i>Journal of Neuroimmunology</i> , 1998, 86, 13-19.	2.3	63
111	Chronic fatigue syndrome differs from fibromyalgia. No evidence for elevated substance P levels in cerebrospinal fluid of patients with chronic fatigue syndrome. <i>Pain</i> , 1998, 78, 153-155.	4.2	65
112	Soluble CD30 levels in plasma and cerebrospinal fluid in multiple sclerosis, HIV infection and other nervous system diseases. <i>Acta Neurologica Scandinavica</i> , 1997, 95, 99-102.	2.1	4
113	Absence of seven human herpesviruses, including HHV-6, by polymerase chain reaction in CSF and blood from patients with multiple sclerosis and optic neuritis. <i>Acta Neurologica Scandinavica</i> , 1997, 95, 280-283.	2.1	101
114	Multiple sclerosis and amyloid deposits in the white matter of the brain. <i>Acta Neuropathologica</i> , 1997, 93, 205-209.	7.7	4
115	Guillain-Barré syndrome in South-West Stockholm, 1973-1991, 2. <i>Clinical epidemiology. Italian Journal of Neurological Sciences</i> , 1997, 18, 49-53.	0.1	6
116	A zone immunoelectrophoresis assay method for quantification of apolipoprotein D in human cerebrospinal fluid. <i>Journal of Proteomics</i> , 1996, 33, 1-8.	2.4	4
117	Increased interleukin-6 mRNA expression in blood and cerebrospinal fluid mononuclear cells in multiple sclerosis. <i>Journal of Neuroimmunology</i> , 1996, 64, 63-69.	2.3	80
118	Multiple sclerosis: the proinflammatory cytokines lymphotoxin- $\alpha$ and tumour necrosis factor- $\alpha$ are upregulated in cerebrospinal fluid mononuclear cells. <i>Journal of Neuroimmunology</i> , 1996, 66, 115-123.	2.3	52
119	Tumor necrosis factor- $\alpha$ , lymphotoxin, interleukin (IL)-6, IL-10, IL-12 and perforin mRNA expression in mononuclear cells in response to acetylcholine receptor is augmented in myasthenia gravis. <i>Journal of Neuroimmunology</i> , 1996, 71, 191-198.	2.3	47
120	Lyme neuroborreliosis: cerebrospinal fluid contains myelin protein-reactive cells secreting interferon- $\gamma$ . <i>European Journal of Neurology</i> , 1996, 3, 122-129.	3.3	4
121	Nasal spray desmopressin treatment of bladder dysfunction in patients with multiple sclerosis. <i>Acta Neurologica Scandinavica</i> , 1996, 94, 31-34.	2.1	37
122	Increased mRNA Expression of IL-10 in Mononuclear Cells in Multiple Sclerosis and Optic Neuritis. <i>Scandinavian Journal of Immunology</i> , 1995, 41, 171-178.	2.7	64
123	Lyme Neuroborreliosis: Evidence for Persistent Up-Regulation of Borrelia Burgdorferi-Reactive Cells Secreting Interferon-gamma. <i>Scandinavian Journal of Immunology</i> , 1995, 42, 694-700.	2.7	27
124	Expression of IFN- $\gamma$ , IL-4, and TGF- $\beta$ in multiple sclerosis in relation to HLA-Dw2 phenotype and stage of disease. <i>Multiple Sclerosis Journal</i> , 1995, 1, 173-180.	3.0	19
125	Analysis of CD27 surface expression on T cell subsets in MS patients and control individuals. <i>Journal of Neuroimmunology</i> , 1995, 56, 99-105.	2.3	22
126	Transforming growth factor- $\beta$ 1 suppresses autoantigen-induced expression of pro-inflammatory cytokines but not of interleukin-10 in multiple sclerosis and myasthenia gravis. <i>Journal of Neuroimmunology</i> , 1995, 58, 21-35.	2.3	46



#	ARTICLE	IF	CITATIONS
127	Guillain-Barré syndrome in South-West Stockholm, 1973-1991, 1. Quality of registered hospital diagnoses and incidence. <i>Acta Neurologica Scandinavica</i> , 1995, 91, 109-117.	2.1	41
128	Organ-specific autoantigens induce transforming growth factor- $\beta$ mRNA expression in mononuclear cells in multiple sclerosis and myasthenia gravis. <i>Annals of Neurology</i> , 1994, 35, 197-203.	5.3	54
129	$\beta$ T cells are increased in patients with Parkinson's disease. <i>Journal of the Neurological Sciences</i> , 1994, 121, 39-45.	0.6	103
130	Augmented interferon- $\beta$ , interleukin-4 and transforming growth factor- $\beta$ mRNA expression in blood mononuclear cells in myasthenia gravis. <i>Journal of Neuroimmunology</i> , 1994, 51, 185-192.	2.3	36
131	V $\beta$ 1 gene usage, interleukin-2 receptors and adhesion molecules on $\beta$ T cells in inflammatory disease of the nervous system. <i>Journal of Neuroimmunology</i> , 1994, 49, 59-66.	2.3	9
132	The HLA-Dw2 haplotype segregates closely with multiple sclerosis in multiplex families. <i>Journal of Neuroimmunology</i> , 1994, 50, 95-100.	2.3	43
133	Multiple sclerosis: occurrence of myelin basic protein peptide-reactive T cells in healthy family members. <i>Acta Neurologica Scandinavica</i> , 1994, 89, 184-189.	2.1	10
134	Optic neuritis and multiple sclerosis: the T cell repertoires to myelin proteins and MBP peptides change with time. <i>Acta Neurologica Scandinavica</i> , 1994, 90, 10-18.	2.1	12
135	Parkinson's disease and immunological abnormalities: increase of HLA-DR expression on monocytes in cerebrospinal fluid and of CD45RO+ T cells in peripheral blood. <i>Acta Neurologica Scandinavica</i> , 1994, 90, 160-166.	2.1	111
136	T Cells Recognizing Multiple Peptides of Myelin Basic Protein are Found in Blood and Enriched in Cerebrospinal Fluid in Optic Neuritis and Multiple Sclerosis. <i>Scandinavian Journal of Immunology</i> , 1993, 37, 355-368.	2.7	41
137	Interleukin-2 secreting cells in multiple sclerosis and controls. <i>Journal of the Neurological Sciences</i> , 1993, 120, 99-106.	0.6	23
138	Myasthenia gravis: T and B cell reactivities to the $\beta$ -bungarotoxin binding protein presynaptic membrane receptor. <i>Journal of the Neurological Sciences</i> , 1992, 109, 173-181.	0.6	18
139	Virus-reactive and autoreactive T cells are accumulated in cerebrospinal fluid in multiple sclerosis. <i>Journal of Neuroimmunology</i> , 1992, 38, 63-73.	2.3	72
140	The T-Cell Repertoire in Myasthenia Gravis Involves Multiple Cholinergic Receptor Epitopes. <i>Scandinavian Journal of Immunology</i> , 1992, 36, 405-414.	2.7	19
141	T cell responses to human recombinant acetylcholine receptor- $\beta$ subunit in myasthenia gravis and controls. <i>European Journal of Immunology</i> , 1992, 22, 1553-1559.	2.9	28
142	CD5+ B cells and CD4 <sup>+</sup> 8 <sup>+</sup> T cells in neuroimmunological diseases. <i>Journal of Neuroimmunology</i> , 1991, 32, 123-132.	2.3	31
143	Interleukin-6 is elevated in plasma in multiple sclerosis. <i>Journal of Neuroimmunology</i> , 1991, 31, 147-153.	2.3	114
144	Autoreactive T and B cells responding to myelin proteolipid protein in multiple sclerosis and controls. <i>European Journal of Immunology</i> , 1991, 21, 1461-1468.	2.9	246

#	ARTICLE	IF	CITATIONS
145	Cells producing antibodies specific for myelin basic protein region 70â€“89 are predominant in cerebrospinal fluid from patients with multiple sclerosis. <i>European Journal of Immunology</i> , 1991, 21, 2971-2976.	2.9	36
146	The 150-Year Anniversary of Multiple Sclerosis: Does Its Early History Give an Etiological Clue?. <i>Perspectives in Biology and Medicine</i> , 1989, 32, 237-243.	0.5	18
147	Mononuclear Cell Types in Cerebrospinal Fluid and Blood of Patients With Multiple Sclerosis. <i>Archives of Neurology</i> , 1989, 46, 372.	4.5	8
148	Retrovirus in multiple sclerosis. <i>Acta Neurologica Scandinavica</i> , 1989, 80, 467-471.	2.1	4
149	Total, anti-viral, and anti-myelin IgG subclass reactivity in inflammatory diseases of the central nervous system. <i>Journal of Neurology</i> , 1989, 236, 238-242.	3.6	31
150	Bone marrow cells in patients with multiple sclerosis. <i>Journal of Neuroimmunology</i> , 1989, 24, 23-31.	2.3	4
151	Primarily chronic progressive and relapsing/remitting multiple sclerosis: two immunogenetically distinct disease entities.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 7113-7117.	7.1	241
152	CNS-borreliosis selectively affecting central motor neurons. <i>Acta Neurologica Scandinavica</i> , 1988, 78, 181-184.	2.1	34
153	Bone Marrow Cells in Multiple Sclerosis.. <i>Annals of the New York Academy of Sciences</i> , 1988, 540, 282-285.	3.8	3
154	Combination ELISAs for antiviral antibodies in CSF and serum in patients with neurological symptoms and in healthy controls. <i>Journal of Virological Methods</i> , 1988, 19, 169-179.	2.1	18
155	HLA CLASS II GENES IN CHRONIC PROGRESSIVE AND IN RELAPSING/REMITTING MULTIPLE SCLEROSIS. <i>Lancet, The</i> , 1987, 330, 327.	13.7	13
156	Increased reactivity to HTLV-I in inflammatory nervous system diseases. <i>Annals of Neurology</i> , 1987, 22, 67-71.	5.3	22
157	CSF neopterin as marker of disease activity in multiple sclerosis. <i>Acta Neurologica Scandinavica</i> , 1987, 75, 352-355.	2.1	45
158	Genomic HLAâ€“typing by RFLPâ€“analysis using DRÎ² and DQÎ² cDNA probes reveals normal DRâ€“DQ linkages in patients with multiple sclerosis. <i>Tissue Antigens</i> , 1987, 30, 135-138.	1.0	10