Shigeki Moriguchi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6294066/publications.pdf

Version: 2024-02-01

40 papers 1,210 citations

331670 21 h-index 34 g-index

42 all docs

42 docs citations 42 times ranked 1635 citing authors

#	Article	IF	CITATIONS
1	Propolis Promotes Memantine-Dependent Rescue of Cognitive Deficits in APP-KI Mice. Molecular Neurobiology, 2022, 59, 4630-4646.	4.0	4
2	Memantine improves cognitive deficits via KATP channel inhibition in olfactory bulbectomized mice. Molecular and Cellular Neurosciences, 2021, 117, 103680.	2.2	8
3	Kir6.1 Heterozygous Mice Exhibit Aberrant Amygdala-Dependent Cued Fear Memory. Molecular Neurobiology, 2020, 57, 1622-1635.	4.0	3
4	Nicotine Rescues Depressive-like Behaviors via α7-type Nicotinic Acetylcholine Receptor Activation in CaMKIV Null Mice. Molecular Neurobiology, 2020, 57, 4929-4940.	4.0	11
5	Memantine Improves Depressive-like Behaviors via Kir6.1 Channel Inhibition in Olfactory Bulbectomized Mice. Neuroscience, 2020, 442, 264-273.	2.3	9
6	Aberrant Amygdala-Dependent Cued Fear Memory in Na+/Ca2+ Exchanger 1 Heterozygous Mice. Molecular Neurobiology, 2019, 56, 4381-4394.	4.0	1
7	Reduced expression of Na+/Ca2+ exchangers is associated with cognitive deficits seen in Alzheimer's disease model mice. Neuropharmacology, 2018, 131, 291-303.	4.1	23
8	Aberrant Amygdala-dependent Fear Memory in Corticosterone-treated Mice. Neuroscience, 2018, 388, 448-459.	2.3	9
9	KATP channel is novel targets for Alzheimer's disease therapeutics. Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2018, WCP2018, PO1-1-40.	0.0	0
10	Combined Memantine and Donepezil Treatment Improves Behavioral and Psychological Symptoms of Dementia-Like Behaviors in Olfactory Bulbectomized Mice. Pharmacology, 2017, 99, 160-171.	2.2	17
11	Reduced CaM Kinase II and CaM Kinase IV Activities Underlie Cognitive Deficits in NCKX2 Heterozygous Mice. Molecular Neurobiology, 2017, 55, 3889-3900.	4.0	13
12	Stimulation of the Sigma-1 Receptor and the Effects on Neurogenesis and Depressive Behaviors in Mice. Advances in Experimental Medicine and Biology, 2017, 964, 201-211.	1.6	19
13	Functional Genomic Analyses Identify Pathways Dysregulated in Animal Model of Autism. CNS Neuroscience and Therapeutics, 2016, 22, 845-853.	3.9	22
14	Stimulation of Sigma-1 Receptor Ameliorates Depressive-like Behaviors in CaMKIV Null Mice. Molecular Neurobiology, 2015, 52, 1210-1222.	4.0	42
15	The extracellular fragment of <scp>GPNMB</scp> (Glycoprotein nonmelanosoma protein B,) Tj ETQq1 1 0.78431 Neurochemistry, 2015, 132, 583-594.	14 rgBT /O\ 3.9	verlock 10 Tf 16
16	Aberrant Behavioral Sensitization by Methamphetamine in Junctophilin-Deficient Mice. Molecular Neurobiology, 2015, 51, 533-542.	4.0	10
17	Ca <scp>MKII</scp> activity is essential for improvement of memoryâ€related behaviors by chronic rivastigmine treatment. Journal of Neurochemistry, 2014, 128, 927-937.	3.9	16
18	Novel nootropic drug sunifiram improves cognitive deficits via CaM kinase II and protein kinase C activation in olfactory bulbectomized mice. Behavioural Brain Research, 2013, 242, 150-157.	2.2	10

#	Article	IF	CITATIONS
19	Novel nootropic drug sunifiram enhances hippocampal synaptic efficacy via glycineâ€binding site of ⟨i>Nâ€methylâ€ <scp>D</scp> â€aspartate receptor. Hippocampus, 2013, 23, 942-951.	1.9	9
20	Stimulation of the Sigma-1 Receptor by DHEA Enhances Synaptic Efficacy and Neurogenesis in the Hippocampal Dentate Gyrus of Olfactory Bulbectomized Mice. PLoS ONE, 2013, 8, e60863.	2.5	75
21	Reduced calcium/calmodulinâ€dependent protein kinase II activity in the hippocampus is associated with impaired cognitive function in MPTPâ€treated mice. Journal of Neurochemistry, 2012, 120, 541-551.	3.9	71
22	The Tâ€type voltageâ€gated calcium channel as a molecular target of the novel cognitive enhancer ST101: enhancement of longâ€term potentiation and CaMKII autophosphorylation in rat cortical slices. Journal of Neurochemistry, 2012, 121, 44-53.	3.9	55
23	Ca2+/calmodulin-dependent protein kinase II and protein kinase C activities mediate extracellular glucose-regulated hippocampal synaptic efficacy. Molecular and Cellular Neurosciences, 2011, 46, 101-107.	2.2	10
24	Pharmacological Study on Alzheimer's Drugs Targeting Calcium/Calmodulin-Dependent Protein Kinase II. Journal of Pharmacological Sciences, 2011, 117, 6-11.	2.5	16
25	Sigma-1 receptor stimulation by dehydroepiandrosterone ameliorates cognitive impairment through activation of CaM kinase II, protein kinase C and extracellular signal-regulated kinase in olfactory bulbectomized mice. Journal of Neurochemistry, 2011, 117, 879-891.	3.9	57
26	A Novel Cognitive Enhancer, ZSET1446/ST101, Promotes Hippocampal Neurogenesis and Ameliorates Depressive Behavior in Olfactory Bulbectomized Mice. Journal of Pharmacology and Experimental Therapeutics, 2010, 333, 43-50.	2.5	45
27	Galantamine enhancement of longâ€ŧerm potentiation is mediated by calcium/calmodulinâ€dependent protein kinase II and protein kinase C activation. Hippocampus, 2009, 19, 844-854.	1.9	30
28	Nefiracetam activation of CaM kinase II and protein kinase C mediated by NMDA and metabotropic glutamate receptors in olfactory bulbectomized mice. Journal of Neurochemistry, 2009, 110, 170-181.	3.9	27
29	CaM kinase II and protein kinase C activations mediate enhancement of longâ€term potentiation by nefiracetam in the rat hippocampal CA1 region. Journal of Neurochemistry, 2008, 106, 1092-1103.	3.9	35
30	Spiro[imidazo[1,2- <i>a</i>]pyridine-3,2-indan]-2(3 <i>H</i>)-one (ZSET1446/ST101) Treatment Rescues Olfactory Bulbectomy-Induced Memory Impairment by Activating Ca ²⁺ /Calmodulin Kinase II and Protein Kinase C in Mouse Hippocampus. Journal of Pharmacology and Experimental Therapeutics, 2008, 326, 127-134.	2.5	43
31	Nefiracetam Potentiates N-Methyl-d-aspartate (NMDA) Receptor Function via Protein Kinase C Activation and Reduces Magnesium Block of NMDA Receptor. Molecular Pharmacology, 2007, 71, 580-587.	2.3	33
32	Effects of Ethanol on Excitatory and Inhibitory Synaptic Transmission in Rat Cortical Neurons. Alcoholism: Clinical and Experimental Research, 2007, 31, 89-99.	2.4	29
33	Constitutively active calcineurin mediates delayed neuronal death through Fas-ligand expression via activation of NFAT and FKHR transcriptional activities in mouse brain ischemia. Journal of Neurochemistry, 2007, 102, 1506-1517.	3.9	72
34	Decreased calcium/calmodulin-dependent protein kinase II and protein kinase C activities mediate impairment of hippocampal long-term potentiation in the olfactory bulbectomized mice. Journal of Neurochemistry, 2006, 97, 22-29.	3.9	72
35	Generation of constitutively active calcineurin by calpain contributes to delayed neuronal death following mouse brain ischemia. Journal of Neurochemistry, 2006, 98, 310-320.	3.9	78
36	Functional uncoupling between Ca2+ release and afterhyperpolarization in mutant hippocampal neurons lacking junctophilins. Proceedings of the National Academy of Sciences of the United States of America, 2006, 103, 10811-10816.	7.1	73

#	Article	IF	CITATION
37	Modulation of N-Methyl-d-aspartate Receptors by Donepezil in Rat Cortical Neurons. Journal of Pharmacology and Experimental Therapeutics, 2005, 315, 125-135.	2.5	26
38	Mechanism of Action of Galantamine on N-Methyl-d-Aspartate Receptors in Rat Cortical Neurons. Journal of Pharmacology and Experimental Therapeutics, 2004, 310, 933-942.	2.5	52
39	Potentiation of N-Methyl-d-aspartate-Induced Currents by the Nootropic Drug Nefiracetam in Rat Cortical Neurons. Journal of Pharmacology and Experimental Therapeutics, 2003, 307, 160-167.	2.5	36
40	Enhancement of N-methyl-D-aspartate receptor-mediated excitatory postsynaptic potentials in the neostriatum after methamphetamine sensitization. Experimental Brain Research, 2002, 144, 238-246.	1.5	31