

Pilar Prieto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6300646/publications.pdf>

Version: 2024-02-01

50
papers

5,559
citations

236612

25
h-index

214527

47
g-index

51
all docs

51
docs citations

51
times ranked

5791
citing authors

#	ARTICLE	IF	CITATIONS
1	Spectrophotometric Quantitation of Antioxidant Capacity through the Formation of a Phosphomolybdenum Complex: Specific Application to the Determination of Vitamin E. <i>Analytical Biochemistry</i> , 1999, 269, 337-341.	1.1	3,789
2	Root Hairs Play a Key Role in the Endophytic Colonization of Olive Roots by <i>Pseudomonas</i> spp. with Biocontrol Activity. <i>Microbial Ecology</i> , 2011, 62, 435-445.	1.4	142
3	Homologue recognition during meiosis is associated with a change in chromatin conformation. <i>Nature Cell Biology</i> , 2004, 6, 906-908.	4.6	135
4	Colonization process of olive tissues by <i>Verticillium dahliae</i> and its interaction with the biocontrol root endophyte <i>Pseudomonas fluorescens</i> PICF7. <i>Microbial Biotechnology</i> , 2009, 2, 499-511.	2.0	127
5	Effective chromosome pairing requires chromatin remodeling at the onset of meiosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 6075-6080.	3.3	97
6	Chromosomes associate premeiotically and in xylem vessel cells via their telomeres and centromeres in diploid rice (<i>Oryza sativa</i>). <i>Chromosoma</i> , 2004, 112, 300-307.	1.0	71
7	A core genetic map of <i>Hordeum chilense</i> and comparisons with maps of barley (<i>Hordeum vulgare</i>) and wheat (<i>Triticum aestivum</i>). <i>Theoretical and Applied Genetics</i> , 2001, 102, 1259-1264.	1.8	63
8	Molecular characterization of TaSTOP1 homoeologues and their response to aluminium and proton (H ⁺) toxicity in bread wheat (<i>Triticum aestivum</i> L.). <i>BMC Plant Biology</i> , 2013, 13, 134.	1.6	61
9	Complete genome sequence of <i>Pseudomonas fluorescens</i> strain PICF7, an indigenous root endophyte from olive (<i>Olea europaea</i> L.) and effective biocontrol agent against <i>Verticillium dahliae</i> . <i>Standards in Genomic Sciences</i> , 2015, 10, 10.	1.5	60
10	Endophytic colonization of olive roots by the biocontrol strain <i>Pseudomonas fluorescens</i> PICF7. <i>FEMS Microbiology Ecology</i> , 2008, 64, 297-306.	1.3	56
11	Fate of <i>Trichoderma harzianum</i> in the olive rhizosphere: time course of the root colonization process and interaction with the fungal pathogen <i>Verticillium dahliae</i> . <i>BioControl</i> , 2016, 61, 269-282.	0.9	56
12	<i>Arabidopsis thaliana</i> as a tool to identify traits involved in <i>Verticillium dahliae</i> biocontrol by the olive root endophyte <i>Pseudomonas fluorescens</i> PICF7. <i>Frontiers in Microbiology</i> , 2015, 06, 266.	1.5	55
13	Bacterial endophytes and root hairs. <i>Plant and Soil</i> , 2012, 361, 301-306.	1.8	54
14	Control of Seed Germination and Plant Development by Carbon and Nitrogen Availability. <i>Frontiers in Plant Science</i> , 2015, 6, 1023.	1.7	52
15	Endophytic colonization and biocontrol performance of <i>Pseudomonas fluorescens</i> PICF7 in olive (<i>Olea europaea</i> L.) are determined neither by pyoverdine production nor swimming motility. <i>Environmental Microbiology</i> , 2015, 17, 3139-3153.	1.8	51
16	Efficient colonization of the endophytes <i>Herbaspirillum huttiense</i> RCA24 and <i>Enterobacter cloacae</i> RCA25 influences the physiological parameters of <i>Oryza sativa</i> L. cv. Baldo rice. <i>Environmental Microbiology</i> , 2019, 21, 3489-3504.	1.8	47
17	Fluorescence in situ hybridization on vibratome sections of plant tissues. <i>Nature Protocols</i> , 2007, 2, 1831-1838.	5.5	42
18	The subtelomeric region is important for chromosome recognition and pairing during meiosis. <i>Scientific Reports</i> , 2014, 4, 6488.	1.6	39

#	ARTICLE	IF	CITATIONS
19	Characterization of two homeodomain transcription factors with critical but distinct roles in virulence in the vascular pathogen <i>Verticillium dahliae</i> . <i>Molecular Plant Pathology</i> , 2018, 19, 986-1004.	2.0	39
20	Control of conformation changes associated with homologue recognition during meiosis. <i>Theoretical and Applied Genetics</i> , 2005, 111, 505-510.	1.8	37
21	The use of the <i>ph1b</i> mutant to induce recombination between the chromosomes of wheat and barley. <i>Frontiers in Plant Science</i> , 2015, 6, 160.	1.7	36
22	Development of <i>Colletotrichum acutatum</i> on Tolerant and Susceptible <i>Olea europaea</i> L. cultivars: A Microscopic Analysis. <i>Mycopathologia</i> , 2009, 168, 203-211.	1.3	32
23	From the root to the stem: interaction between the biocontrol root endophyte <i>Pseudomonas fluorescens</i> PICF7 and the pathogen <i>Pseudomonas savastanoi</i> NCPPB 3335 in olive knots. <i>Microbial Biotechnology</i> , 2013, 6, 275-287.	2.0	31
24	Chromosomal distribution of telomeric and telomeric-associated sequences in <i>Hordeum chilense</i> by in situ hybridization. <i>Hereditas</i> , 2004, 141, 122-127.	0.5	30
25	Identification of Intergenomic Translocations Involving Wheat, <i>Hordeum Vulgare</i> and <i>Hordeum Chilense</i> Chromosomes by FISH. <i>Hereditas</i> , 2004, 135, 171-174.	0.5	29
26	Mammalian cell entry genes in <i>Streptomyces</i> may provide clues to the evolution of bacterial virulence. <i>Scientific Reports</i> , 2013, 3, 1109.	1.6	27
27	Unravelling the proteomic profile of rice meiocytes during early meiosis. <i>Frontiers in Plant Science</i> , 2014, 5, 356.	1.7	26
28	<i>Pseudomonas fluorescens</i> PICF7 displays an endophytic lifestyle in cultivated cereals and enhances yield in barley. <i>FEMS Microbiology Ecology</i> , 2016, 92, f1w092.	1.3	25
29	Molecular diversity of bacterial endosymbionts associated with dagger nematodes of the genus <i>Xiphinema</i> (Nematoda: Longidoridae) reveals a high degree of phylogenetic congruence with their host. <i>Molecular Ecology</i> , 2016, 25, 6225-6247.	2.0	23
30	Identification and validation of reference genes for RT-qPCR normalization in wheat meiosis. <i>Scientific Reports</i> , 2020, 10, 2726.	1.6	23
31	Novel Bread Wheat Lines Enriched in Carotenoids Carrying <i>Hordeum chilense</i> Chromosome Arms in the <i>ph1b</i> Background. <i>PLoS ONE</i> , 2015, 10, e0134598.	1.1	23
32	Genomic and Meiotic Changes Accompanying Polyploidization. <i>Plants</i> , 2022, 11, 125.	1.6	23
33	Development of <i>Hordeum chilense</i> 4Hch introgression lines in durum wheat: a tool for breeders and complex trait analysis. <i>Plant Breeding</i> , 2012, 131, 733-738.	1.0	20
34	Assessing the Involvement of Selected Phenotypes of <i>Pseudomonas simiae</i> PICF7 in Olive Root Colonization and Biological Control of <i>Verticillium dahliae</i> . <i>Plants</i> , 2021, 10, 412.	1.6	20
35	Detection of alien genetic introgressions in bread wheat using dot-blot genomic hybridisation. <i>Molecular Breeding</i> , 2017, 37, 32.	1.0	18
36	Telomeres and Subtelomeres Dynamics in the Context of Early Chromosome Interactions During Meiosis and Their Implications in Plant Breeding. <i>Frontiers in Plant Science</i> , 2021, 12, 672489.	1.7	17

#	ARTICLE	IF	CITATIONS
37	Sequence analysis of wheat subtelomeres reveals a high polymorphism among homoeologous chromosomes. <i>Plant Genome</i> , 2020, 13, e20065.	1.6	15
38	Homoeologous Chromosomes From Two <i>Hordeum</i> Species Can Recognize and Associate During Meiosis in Wheat in the Presence of the Ph1 Locus. <i>Frontiers in Plant Science</i> , 2018, 9, 585.	1.7	14
39	Mycovirus <i>Fusarium oxysporum</i> f. sp. <i>dianthi</i> Virus 1 Decreases the Colonizing Efficiency of Its Fungal Host. <i>Frontiers in Cellular and Infection Microbiology</i> , 2019, 9, 51.	1.8	13
40	Evaluation of Indigenous Olive Biocontrol Rhizobacteria as Protectants against Drought and Salt Stress. <i>Microorganisms</i> , 2021, 9, 1209.	1.6	8
41	Endophytes from African Rice (<i>Oryza glaberrima</i> L.) Efficiently Colonize Asian Rice (<i>Oryza sativa</i> L.) Stimulating the Activity of Its Antioxidant Enzymes and Increasing the Content of Nitrogen, Carbon, and Chlorophyll. <i>Microorganisms</i> , 2021, 9, 1714.	1.6	8
42	Development and cytogenetic characterisation of a double goat grass-barley chromosome substitution in tritordeum. <i>Euphytica</i> , 2006, 147, 337-342.	0.6	7
43	Wild and Cultivated Homoeologous Barley Chromosomes Can Associate and Recombine in Wheat in the Absence of the Ph1 Locus. <i>Agronomy</i> , 2021, 11, 147.	1.3	4
44	Chromosome Manipulation for Plant Breeding Purposes. <i>Agronomy</i> , 2020, 10, 1695.	1.3	3
45	Dynamics of DNA Replication during Premeiosis and Early Meiosis in Wheat. <i>PLoS ONE</i> , 2014, 9, e107714.	1.1	3
46	Analytical Methodology of Meiosis in Autopolyploid and Allopolyploid Plants. <i>Methods in Molecular Biology</i> , 2020, 2061, 141-168.	0.4	3
47	Homologous chromosome associations in domains before meiosis could facilitate chromosome recognition and pairing in wheat. <i>Scientific Reports</i> , 2022, 12, .	1.6	2
48	Olive “ Colletotrichum acutatum: An Example of Fruit-Fungal Interaction. , 2012, , .		1
49	Effect of <i>H^{ch}</i> <i>Hordeum chilense</i> Chromosome Introgressions on the Wheat Endosperm Proteomic Profile. <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 3793-3802.	2.4	0
50	Analysis of Chromosome Associations during Early Meiosis in Wheat Lines Carrying Chromosome Introgressions from <i>Agropyron cristatum</i> . <i>Plants</i> , 2021, 10, 2292.	1.6	0