
## Simon Gilbody

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6337009/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Shared goals for mental health research: what, why and when for the 2020s. Journal of Mental Health, 2023, 32, 997-1005.	1.9	39
2	Food insecurity and severe mental illness: understanding the hidden problem and how to ask about food access during routine healthcare. BJ Psych Advances, 2023, 29, 204-212.	0.7	8
3	How much change is enough? Evidence from a longitudinal study on depression in UK primary care. Psychological Medicine, 2022, 52, 1875-1882.	4.5	51
4	Can We Prevent Depression in At-Risk Older Adults Using Self-Help? The UK SHARD Trial of Behavioral Activation. American Journal of Geriatric Psychiatry, 2022, 30, 197-207.	1.2	10
5	Healthcare resource use and costs for people with type 2 diabetes mellitus with and without severe mental illness in England: longitudinal matched-cohort study using the Clinical Practice Research Datalink. British Journal of Psychiatry, 2022, 221, 402-409.	2.8	10
6	A systematic review of behavioural smoking cessation interventions for people with severe mental ill health—what works?. Addiction, 2022, 117, 1526-1542.	3.3	8
7	Cost-Utility Analysis of Discontinuing Antidepressants in England Primary Care Patients Compared with Long-Term Maintenance: The ANTLER Study. Applied Health Economics and Health Policy, 2022, 20, 269-282.	2.1	6
8	External validation of a shortened screening tool using individual participant data meta-analysis: A case study of the Patient Health Questionnaire-Dep-4. Methods, 2022, 204, 300-311.	3.8	3
9	Life events and treatment prognosis for depression: A systematic review and individual patient data meta-analysis. Journal of Affective Disorders, 2022, 299, 298-308.	4.1	7
10	Loneliness among people with severe mental illness during the COVID-19 pandemic: Results from a linked UK population cohort study. PLoS ONE, 2022, 17, e0262363.	2.5	22
11	Cost-utility analysis of LEGO based therapy for school children and young people with autism spectrum disorder: results from a randomised controlled trial. BMJ Open, 2022, 12, e056347.	1.9	5
12	Predicting relapse or recurrence of depression: systematic review of prognostic models. British Journal of Psychiatry, 2022, 221, 448-458.	2.8	13
13	The feasibility and acceptability of a brief psychological intervention for adults with long-term health conditions and subthreshold depression delivered via community pharmacies: a mixed methods evaluation—the Community Pharmacies Mood Intervention Study (CHEMIST). Pilot and Feasibility Studies, 2022, 8, 27.	1.2	3
14	A psychological intervention by community pharmacies to prevent depression in adults with subthreshold depression and long-term conditions: the CHEMIST pilot RCT. Public Health Research, 2022, 10, 1-160.	1.3	2
15	Socioeconomic Indicators of Treatment Prognosis for Adults With Depression. JAMA Psychiatry, 2022, 79, 406.	11.0	30
16	Behavioural Activation for Social IsoLation (BASIL+) trial (Behavioural activation to mitigate) Tj ETQq0 0 0 rgBT /( fully-powered pragmatic randomised controlled trial. PLoS ONE, 2022, 17, e0263856.	Overlock 1 2.5	0 Tf 50 147 1 7
17	A Qualitative Study Exploring the Barriers and Facilitators for Maintaining Oral Health and Using Dental Service in People with Severe Mental Illness: Perspectives from Service Users and Service Providers. International Journal of Environmental Research and Public Health, 2022, 19, 4344.	2.6	9
18	The DIAMONDS intervention to support self-management of type 2 diabetes in people with severe mental illness: Study protocol for a single-group feasibility study. SSM Mental Health, 2022, 2, 100086.	1.8	4

2

#	Article	IF	CITATIONS
19	Multiple versus single risk behaviour interventions for people with severe mental illness: a network meta-analysis and qualitative synthesis. , 2022, 10, 1-186.		1
20	Cognitive, behavioural or cognitive-behavioural self-help interventions for subclinical depression in older adults: A systematic review and meta-analysis. Journal of Affective Disorders, 2022, 308, 384-390.	4.1	4
21	Behavioral Activation in Nursing Homes to Treat Depression (BAN-Dep): Results From a Clustered, Randomized, Single-Blinded, Controlled Clinical Trial. American Journal of Geriatric Psychiatry, 2022, 30, 1313-1323.	1.2	6
22	Effectiveness of interventions to address obesity and health risk behaviours among people with severe mental illness in low- and middle-income countries (LMICs): a systematic review and meta analysis. Global Mental Health (Cambridge, England), 2022, 9, 264-273.	2.5	2
23	Depression prevalence based on the Edinburgh Postnatal Depression Scale compared to Structured Clinical Interview for DSM Disorders classification: Systematic review and individual participant data metaâ€analysis. International Journal of Methods in Psychiatric Research, 2021, 30, e1860.	2.1	30
24	Preventing depression among older people living in rural areas: A randomised controlled trial of behavioural activation in collaborative care. International Journal of Geriatric Psychiatry, 2021, 36, 530-539.	2.7	12
25	Probability of Major Depression Classification Based on the SCID, CIDI, and MINI Diagnostic Interviews: A Synthesis of Three Individual Participant Data Meta-Analyses. Psychotherapy and Psychosomatics, 2021, 90, 28-40.	8.8	20
26	Smoking cessation in severe mental illness: combined long-term quit rates from the UK SCIMITAR trials programme. British Journal of Psychiatry, 2021, 218, 95-97.	2.8	13
27	Impact of severe mental illness on healthcare use and health outcomes for people with type 2 diabetes: a longitudinal observational study in England. British Journal of General Practice, 2021, 71, e565-e573.	1.4	18
28	ls social support preâ€treatment associated with prognosis for adults with depression in primary care?. Acta Psychiatrica Scandinavica, 2021, 143, 392-405.	4.5	17
29	How can we estimate QALYs based on PHQ-9 scores? Equipercentile linking analysis of PHQ-9 and EQ-5D. Evidence-Based Mental Health, 2021, 24, 97-101.	4.5	11
30	Supporting smoking cessation and preventing relapse following a stay in a smokeâ€free setting: a metaâ€analysis and investigation of effective behaviour change techniques. Addiction, 2021, 116, 2978-2994.	3.3	17
31	Health of mothers of children with a life-limiting condition: a comparative cohort study. Archives of Disease in Childhood, 2021, 106, 987-993.	1.9	14
32	Investigating smoking and nicotine dependence among people with severe mental illness during the COVID-19 pandemic: analysis of linked data from a UK Closing the Gap cohort. BJPsych Open, 2021, 7, e86.	0.7	10
33	Internet-Based Cognitive Behavioral Therapy for Depression. JAMA Psychiatry, 2021, 78, 361.	11.0	398
34	The contribution of depressive †disorder characteristics' to determinations of prognosis for adults with depression: an individual patient data meta-analysis. Psychological Medicine, 2021, 51, 1068-1081.	4.5	28
35	The digital divide: amplifying health inequalities for people with severe mental illness in the time of COVID-19. British Journal of Psychiatry, 2021, 219, 529-531.	2.8	40
36	Predicting prognosis for adults with depression using individual symptom data: a comparison of modelling approaches. Psychological Medicine, 2021, , 1-11.	4.5	19

#	Article	IF	CITATIONS
37	Living with diabetes alongside a severe mental illness: A qualitative exploration with people with severe mental illness, family members and healthcare staff. Diabetic Medicine, 2021, 38, e14562.	2.3	25
38	The importance of transdiagnostic symptom level assessment to understanding prognosis for depressed adults: analysis of data from six randomised control trials. BMC Medicine, 2021, 19, 109.	5.5	20
39	Prognostic models for predicting relapse or recurrence of major depressive disorder in adults. The Cochrane Library, 2021, 2021, CD013491.	2.8	13
40	Identifying determinants of diabetes risk and outcomes for people with severe mental illness: a mixed-methods study. Health Services and Delivery Research, 2021, 9, 1-194.	1.4	2
41	Barriers and facilitators to implementing perinatal mental health care in health and social care settings: a systematic review. Lancet Psychiatry,the, 2021, 8, 521-534.	7.4	54
42	Dismantling, optimising, and personalising internet cognitive behavioural therapy for depression: a systematic review and component network meta-analysis using individual participant data. Lancet Psychiatry,the, 2021, 8, 500-511.	7.4	105
43	Cost-effectiveness of feedback-informed psychological treatment: Evidence from the IAPT-FIT trial. Behaviour Research and Therapy, 2021, 142, 103873.	3.1	16
44	Effectiveness of collaborative care in reducing suicidal ideation: An individual participant data meta-analysis General Hospital Psychiatry, 2021, 71, 27-35.	2.4	8
45	The development and validation of a prognostic model to PREDICT Relapse of depression in adult patients in primary care: protocol for the PREDICTR study. Diagnostic and Prognostic Research, 2021, 5, 12.	1.8	3
46	Home environmental assessments and modification delivered by occupational therapists to reduce falls in people aged 65 years and over: the OTIS RCT. Health Technology Assessment, 2021, 25, 1-118.	2.8	8
47	Still â€ <sup>-</sup> being bothered about Billy': managing the physical health of people with severe mental illness. British Journal of General Practice, 2021, 71, 373-376.	1.4	6
48	Mitigating the impacts of COVID-19: where are the mental health trials?. Lancet Psychiatry,the, 2021, 8, 647-650.	7.4	15
49	Relationship between Green and Blue Spaces with Mental and Physical Health: A Systematic Review of Longitudinal Observational Studies. International Journal of Environmental Research and Public Health, 2021, 18, 9010.	2.6	41
50	Long-Term Cost-Effectiveness of Smoking Cessation Interventions in People With Mental Disorders: A Dynamic Decision Analytical Model. Value in Health, 2021, 24, 1263-1272.	0.3	3
51	Data-driven methods distort optimal cutoffs and accuracy estimates of depression screening tools: a simulation study using individual participant data. Journal of Clinical Epidemiology, 2021, 137, 137-147.	5.0	9
52	Use of the Internet and Digital Devices Among People With Severe Mental III Health During the COVID-19 Pandemic Restrictions. Frontiers in Psychiatry, 2021, 12, 732735.	2.6	15
53	Maintenance or Discontinuation of Antidepressants in Primary Care. New England Journal of Medicine, 2021, 385, 1257-1267.	27.0	75
54	Nature-based outdoor activities for mental and physical health: Systematic review and meta-analysis. SSM - Population Health, 2021, 16, 100934.	2.7	96

#	Article	IF	CITATIONS
55	Role of age, gender and marital status in prognosis for adults with depression: An individual patient data meta-analysis. Epidemiology and Psychiatric Sciences, 2021, 30, e42.	3.9	43
56	Shortening the Edinburgh postnatal depression scale using optimal test assembly methods: Development of the EPDSâ€Đepâ€5. Acta Psychiatrica Scandinavica, 2021, 143, 348-362.	4.5	5
57	Health risk behaviours among people with severe mental ill health during the COVID-19 pandemic: Analysis of linked cohort data. PLoS ONE, 2021, 16, e0258349.	2.5	6
58	Behavioural activation to prevent depression and loneliness among socially isolated older people with long-term conditions: The BASIL COVID-19 pilot randomised controlled trial. PLoS Medicine, 2021, 18, e1003779.	8.4	31
59	Interventions for preventing relapse or recurrence of major depressive disorder in adults in a primary care setting: a network meta-analysis. The Cochrane Library, 2021, 2021, .	2.8	0
60	Antidepressant medication to prevent depression relapse in primary care: the ANTLER RCT. Health Technology Assessment, 2021, 25, 1-62.	2.8	9
61	A Year Into the Pandemic: The Diversity of Experience Amongst People With Severe Mental III Health. Frontiers in Psychiatry, 2021, 12, 794585.	2.6	4
62	A Patient Stratification Approach to Identifying the Likelihood of Continued Chronic Depression and Relapse Following Treatment for Depression. Journal of Personalized Medicine, 2021, 11, 1295.	2.5	10
63	Effectiveness and adherence of telephone-administered psychotherapy for depression: A systematic review and meta-analysis. Journal of Affective Disorders, 2020, 260, 514-526.	4.1	55
64	The Accuracy of the Patient Health Questionnaire-9 Algorithm for Screening to Detect Major Depression: An Individual Participant Data Meta-Analysis. Psychotherapy and Psychosomatics, 2020, 89, 25-37.	8.8	67
65	Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. Psychological Medicine, 2020, 50, 1368-1380.	4.5	175
66	What does the rise and rise of electronic nicotine delivery systems mean for mental health services?. Lancet Psychiatry,the, 2020, 7, 11-13.	7.4	5
67	Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale – Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. Journal of Psychosomatic Research, 2020, 129, 109892.	2.6	33
68	The role of relapse prevention for depression in collaborative care: A systematic review. Journal of Affective Disorders, 2020, 265, 618-644.	4.1	15
69	Standard smoking cessation services in sites participating in the SCIMITAR+ trial for people with severe mental ill health. BJPsych Bulletin, 2020, 44, 6-11.	1.1	1
70	Collaborative care for depression in older adults: How much is enough?. Behaviour Research and Therapy, 2020, 135, 103725.	3.1	0
71	Health of mothers of children with a life-limiting condition: a protocol for comparative cohort study using the Clinical Practice Research Datalink. BMJ Open, 2020, 10, e034024.	1.9	3
72	The associations between loneliness, social exclusion and pain in the general population: A N=502,528 cross-sectional UK Biobank study. Journal of Psychiatric Research, 2020, 130, 68-74.	3.1	29

#	Article	IF	CITATIONS
73	Conversion and neuro-inflammation disorder observational study (CANDO). Protocol of a feasibility study. European Journal of Psychiatry, 2020, 34, 164-172.	1.3	2
74	Overestimation of Postpartum Depression Prevalence Based on a 5-item Version of the EPDS: Systematic Review and Individual Participant Data Meta-analysis. Canadian Journal of Psychiatry, 2020, 65, 835-844.	1.9	9
75	A metaâ€review of "lifestyle psychiatryâ€r the role of exercise, smoking, diet and sleep in the prevention and treatment of mental disorders. World Psychiatry, 2020, 19, 360-380.	10.4	424
76	Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis. PLoS Medicine, 2020, 17, e1003262.	8.4	154
77	Challenges and solutions to nicotine replacement therapy access: observations from SCIMITAR+. BJPsych Open, 2020, 6, e120.	0.7	0
78	Out of pocket expenses in obsessive compulsive disorder. Journal of Mental Health, 2020, , 1-6.	1.9	0
79	REDUCE (Reviewing long-term antidepressant use by careful monitoring in everyday practice) internet and telephone support to people coming off long-term antidepressants: protocol for a randomised controlled trial. Trials, 2020, 21, 419.	1.6	11
80	Individual participant data (IPD) meta-analysis of psychological relapse prevention interventions versus control for patients in remission from depression: a protocol. BMJ Open, 2020, 10, e034158.	1.9	7
81	Accuracy of the PHQ-2 Alone and in Combination With the PHQ-9 for Screening to Detect Major Depression. JAMA - Journal of the American Medical Association, 2020, 323, 2290.	7.4	242
82	E-cigarette use and associated factors among smokers with severe mental illness. Addictive Behaviors, 2020, 108, 106456.	3.0	5
83	Factors that influence participation in physical activity for anxiety or depression: a synthesis of qualitative evidence. The Cochrane Library, 2020, , .	2.8	3
84	Patient Health Questionnaire-9 scores do not accurately estimate depression prevalence: individual participant data meta-analysis. Journal of Clinical Epidemiology, 2020, 122, 115-128.e1.	5.0	113
85	Predicting and preventing relapse of depression in primary care. British Journal of General Practice, 2020, 70, 54-55.	1.4	12
86	Estimating the sample mean and standard deviation from commonly reported quantiles in meta-analysis. Statistical Methods in Medical Research, 2020, 29, 2520-2537.	1.5	366
87	A systematic review of mental health measurement scales for evaluating the effects of mental health prevention interventions. European Journal of Public Health, 2020, 30, 510-516.	0.3	21
88	Costâ€effectiveness of a specialist smoking cessation package compared with standard smoking cessation services for people with severe mental illness in England: a trialâ€based economic evaluation from the SCIMITAR+ study. Addiction, 2020, 115, 2113-2122.	3.3	8
89	Optimised patient information materials and recruitment to a study of behavioural activation in older adults: an embedded study within a trial. F1000Research, 2020, 9, 417.	1.6	3
90	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort. PLoS ONE, 2020, 15, e0243972.	2.5	21

#	Article	IF	CITATIONS
91	The association between primary care quality and health-care use, costs and outcomes for people with serious mental illness: a retrospective observational study. Health Services and Delivery Research, 2020, 8, 1-126.	1.4	2
92	Cost-effectiveness of brief structured interventions to discontinue long-term benzodiazepine use: an economic analysis alongside a randomised controlled trial. HRB Open Research, 2020, 3, 33.	0.6	0
93	Interventions for adults with a history of complex traumatic events: the INCiTE mixed-methods systematic review. Health Technology Assessment, 2020, 24, 1-312.	2.8	19
94	Title is missing!. , 2020, 17, e1003262.		0
95	Title is missing!. , 2020, 17, e1003262.		0
96	Title is missing!. , 2020, 17, e1003262.		0
97	Title is missing!. , 2020, 17, e1003262.		0
98	Title is missing!. , 2020, 17, e1003262.		0
99	Title is missing!. , 2020, 15, e0243972.		0
100	Title is missing!. , 2020, 15, e0243972.		0
101	Title is missing!. , 2020, 15, e0243972.		0
102	Title is missing!. , 2020, 15, e0243972.		0
103	Title is missing!. , 2020, 15, e0243972.		0
104	Title is missing!. , 2020, 15, e0243972.		0
105	Commentary on: Suicide Screening Tools for Use with Incarcerated Offenders. Archives of Suicide Research, 2019, 23, 523-525.	2.3	1
106	Comparison of major depression diagnostic classification probability using the SCID, CIDI, and MINI diagnostic interviews among women in pregnancy or postpartum: An individual participant data metaâ€analysis. International Journal of Methods in Psychiatric Research, 2019, 28, e1803.	2.1	34
107	The clinical effectiveness of sertraline in primary care and the role of depression severity and duration (PANDA): a pragmatic, double-blind, placebo-controlled randomised trial. Lancet Psychiatry,the, 2019, 6, 903-914.	7.4	78
108	Can support workers from AgeUK deliver an intervention to support older people with anxiety and depression? A qualitative evaluation. BMC Family Practice, 2019, 20, 16.	2.9	4

#	Article	IF	CITATIONS
109	Impact of telephone delivered case-management on the effectiveness of collaborative care for depression and anti-depressant use: A systematic review and meta-regression. PLoS ONE, 2019, 14, e0217948.	2.5	15
110	A randomised controlled trial assessing the use of citalopram, sertraline, fluoxetine and mirtazapine in preventing relapse in primary care patients who are taking long-term maintenance antidepressants (ANTLER: ANTidepressants to prevent reLapse in dEpRession): study protocol for a randomised controlled trial. Trials, 2019, 20, 319.	1.6	12
111	Community Pharmacies Mood Intervention Study (CHEMIST): feasibility and external pilot randomised controlled trial protocol. Pilot and Feasibility Studies, 2019, 5, 71.	1.2	4
112	Everything you wanted to know about e-cigarettes and vaping but were afraid to ask: a guide for mental health clinicians. BJ Psych Advances, 2019, 25, 279-286.	0.7	3
113	Association Between Antipsychotic Polypharmacy and Outcomes for People With Serious Mental Illness in England. Psychiatric Services, 2019, 70, 650-656.	2.0	23
114	Smoking cessation for people with severe mental illness (SCIMITAR+): a pragmatic randomised controlled trial. Lancet Psychiatry,the, 2019, 6, 379-390.	7.4	126
115	Accuracy of Patient Health Questionnaire-9 (PHQ-9) for screening to detect major depression: individual participant data meta-analysis. BMJ: British Medical Journal, 2019, 365, 11476.	2.3	822
116	The Effects of Meditation, Yoga, and Mindfulness on Depression, Anxiety, and Stress in Tertiary Education Students: A Meta-Analysis. Frontiers in Psychiatry, 2019, 10, 193.	2.6	105
117	Behavioural activation in nursing homes to treat depression (BAN-Dep): study protocol for a pragmatic randomised controlled trial. BMJ Open, 2019, 9, e032421.	1.9	4
118	Antidepressants for major depression disorder in older people: a network meta-analysis. The Cochrane Library, 2019, , .	2.8	5
119	Pharmacy-based management for depression in adults. The Cochrane Library, 2019, 12, CD013299.	2.8	22
120	Impact of family practice continuity of care on unplanned hospital use for people with serious mental illness. Health Services Research, 2019, 54, 1316-1325.	2.0	27
121	Factors associated with regular physical activity participation among people with severe mental ill health. Social Psychiatry and Psychiatric Epidemiology, 2019, 54, 887-895.	3.1	24
122	Predicting persistent depressive symptoms in older adults: A machine learning approach to personalised mental healthcare. Journal of Affective Disorders, 2019, 246, 857-860.	4.1	69
123	GPs' attitudes towards digital technologies for depression: an online survey in primary care. British Journal of General Practice, 2019, 69, e164-e170.	1.4	19
124	Shortening self-report mental health symptom measures through optimal test assembly methods: Development and validation of the Patient Health Questionnaire-Depression-4. Depression and Anxiety, 2019, 36, 82-92.	4.1	16
125	Investigating SOcial Competence and Isolation in children with Autism taking part in LEGO-based therapy clubs In School Environments (I-SOCIALISE): study protocol. BMJ Open, 2019, 9, e030471.	1.9	9
126	The effectiveness of a contingent financial incentive to improve trial follow up; a randomised study within a trial (SWAT). F1000Research, 2019, 8, 1937.	1.6	4

#	Article	IF	CITATIONS
127	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. Wellcome Open Research, 2019, 4, 69.	1.8	11
128	Exploring Severe Mental Illness and Diabetes: Protocol for a Longitudinal, Observational, and Qualitative Mixed Methods Study. JMIR Research Protocols, 2019, 8, e13407.	1.0	6
129	A bespoke smoking cessation service compared with treatment as usual for people with severe mental ill health: the SCIMITAR+ RCT. Health Technology Assessment, 2019, 23, 1-116.	2.8	10
130	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. Programme Grants for Applied Research, 2019, 7, 1-108.	1.0	6
131	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. Wellcome Open Research, 2019, 4, 69.	1.8	11
132	Non-traditional support workers delivering a brief psychosocial intervention for older people with anxiety and depression: the NOTEPAD feasibility study. Health Services and Delivery Research, 2019, 7, 1-120.	1.4	1
133	SCIMITAR+ Trial: A randomised study within a trial (SWAT) of a contingent financial reward to improve trial follow-up. F1000Research, 2019, 8, 1937.	1.6	3
134	Relapse and Recurrence of Common Mental Health Problems after Low Intensity Cognitive Behavioural Therapy: The WYLOW Longitudinal Cohort Study. Psychotherapy and Psychosomatics, 2018, 87, 116-117.	8.8	14
135	A feasibility study for NOn-Traditional providers to support the management of Elderly People with Anxiety and Depression: The NOTEPAD study Protocol. Trials, 2018, 19, 172.	1.6	4
136	Probability of major depression diagnostic classification using semi-structured versus fully structured diagnostic interviews. British Journal of Psychiatry, 2018, 212, 377-385.	2.8	53
137	Is self-guided internet-based cognitive behavioural therapy (iCBT) harmful? An individual participant data meta-analysis. Psychological Medicine, 2018, 48, 2456-2466.	4.5	106
138	German EstSmoke: estimating adult smokingâ€related costs and consequences of smoking cessation for Germany. Addiction, 2018, 113, 125-136.	3.3	5
139	Do care plans and annual reviews of physical health influence unplanned hospital utilisation for people with serious mental illness? Analysis of linked longitudinal primary and secondary healthcare records in England. BMJ Open, 2018, 8, e023135.	1.9	8
140	Diagnostic accuracy of the Geriatric Depression Scale-30, Geriatric Depression Scale-15, Geriatric Depression Scale-5 and Geriatric Depression Scale-4 for detecting major depression: protocol for a systematic review and individual participant data meta-analysis. BMJ Open, 2018, 8, e026598.	1.9	24
141	The HARMONIC trial: study protocol for a randomised controlled feasibility trial of Shaping Healthy Minds—a modular transdiagnostic intervention for mood, stressor-related and anxiety disorders in adults. BMJ Open, 2018, 8, e024546.	1.9	18
142	Can occupational therapist-led home environmental assessment prevent falls in older people? A modified cohort randomised controlled trial protocol. BMJ Open, 2018, 8, e022488.	1.9	18
143	How should we implement collaborative care for older people with depression? A qualitative study using normalisation process theory within the CASPER plus trial. BMC Family Practice, 2018, 19, 116.	2.9	16
144	Feedback-informed treatment versus usual psychological treatment for depression and anxiety: a multisite, open-label, cluster randomised controlled trial. Lancet Psychiatry,the, 2018, 5, 564-572.	7.4	86

# ARTICLE IF CITATIONS Loneliness, social isolation and risk of cardiovascular disease in the English Longitudinal Study of 145 1.8 Ageing. European Journal of Preventive Cardiology, 2018, 25, 1387-1396. Successful recruitment to trials: findings from the SCIMITAR+ Trial. Trials, 2018, 19, 53. 146 17 1.6 Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA –) Tj ETQq1 1 0.784314 rgBT 1.4 Research, 2018, 6, 1-210. Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. Health 148 2.8 21 Technology Assessment, 2018, 22, 1-136. Collaborative case management to aid return to work after long-term sickness absence: a pilot 149 1.3 randomised controlled trial. Public Health Research, 2018, 6, 1-76. Effect of Collaborative Care vs Usual Care on Depressive Symptoms in Older Adults With 150 7.4 95 Subthreshold Depression. JAMA - Journal of the American Medical Association, 2017, 317, 728. Efficacy of Self-guided Internet-Based Cognitive Behavioral Therapy in the Treatment of Depressive 11.0 560 Symptoms. JAMA Psychiatry, 2017, 74, 351. Cost-effectiveness of computerized cognitiveâ€"behavioural therapy for the treatment of depression in primary care: findings from the Randomised Evaluation of the Effectiveness and Acceptability of 152 4.5 23 Computerised Therapy (REEACT) trial. Psychological Medicine, 2017, 47, 1825-1835. Telephone-supported computerised cognitive–behavioural therapy: REEACT-2 large-scale pragmatic 2.8 94 randomised controlled trial. British Journal of Psychiatry, 2017, 210, 362-367. How durable is the effect of low intensity CBT for depression and anxiety? Remission and relapse in a 154 3.1 97 longitudinal cohort study. Behaviour Research and Therapy, 2017, 94, 1-8. Selective Cutoff Reporting in Studies of Diagnostic Test Accuracy: A Comparison of Conventional and Individual-Patient-Data Meta-Analyses of the Patient Health Questionnaire-9 Depression Screening 3.4Tool. American Journal of Epidemiology, 2017, 185, 954-964. Fathers' views and experiences of their own mental health during pregnancy and the first postnatal 156 year: a qualitative interview study of men participating in the UK Born and Bred in Yorkshire (BaBY) 2.4 146 cohort. BMC Pregnancy and Childbirth, 2017, 17, 45. Impact of long-term medical conditions on the outcomes of psychological therapy for depression and 2.8 29 anxiety. British Journal of Psychiatry, 2017, 210, 47-53. Are there researcher allegiance effects in diagnostic validation studies of the PHQ-9? A systematic 158 1.9 18 review and meta-analysis. BMJ Open, 2017, 7, e015247. Improving the efficiency of psychological treatment using outcome feedback technology. Behaviour Research and Therapy, 2017, 99, 89-97. 3.1 58 Should Google offer an online screening test for depression?. BMJ: British Medical Journal, 2017, 358, 160 2.33 j4144. Identifying primary care quality indicators for people with serious mental illness: a systematic review. 161 1.4 British Journal of General Practice, 2017, 67, e519-e530. An randomized controlled trial of Postâ€it® notes did not increase postal response rates in older 162 1.8 6 depressed participants. Journal of Evaluation in Clinical Practice, 2017, 23, 102-107.

#	Article	IF	CITATIONS
163	Smoking Cessation Intervention for Severe Mental III Health Trial (SCIMITAR+): study protocol for a randomised controlled trial. Trials, 2017, 18, 44.	1.6	15
164	Informing efficient randomised controlled trials: exploration of challenges in developing progression criteria for internal pilot studies. BMJ Open, 2017, 7, e013537.	1.9	239
165	The Effectiveness of Pharmacological and Non-Pharmacological Interventions for Improving Glycaemic Control in Adults with Severe Mental Illness: A Systematic Review and Meta-Analysis. PLoS ONE, 2017, 12, e0168549.	2.5	35
166	Low-intensity cognitive-behaviour therapy interventions for obsessive-compulsive disorder compared to waiting list for therapist-led cognitive-behaviour therapy: 3-arm randomised controlled trial of clinical effectiveness. PLoS Medicine, 2017, 14, e1002337.	8.4	25
167	The ethics of â€~Trials within Cohorts' (TwiCs): 2nd international symposium. Trials, 2017, 18, .	1.6	7
168	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial. Trials, 2017, 18, 496.	1.6	15
169	Smoking cessation in severe mental ill health: what works? an updated systematic review and meta-analysis. BMC Psychiatry, 2017, 17, 252.	2.6	99
170	CollAborative care and active surveillance for Screen-Positive EldeRs with subthreshold depression (CASPER): a multicentred randomised controlled trial of clinical effectiveness and cost-effectiveness. Health Technology Assessment, 2017, 21, 1-196.	2.8	12
171	Clinical effectiveness, cost-effectiveness and acceptability of low-intensity interventions in the management of obsessive–compulsive disorder: the Obsessive–Compulsive Treatment Efficacy randomised controlled Trial (OCTET). Health Technology Assessment, 2017, 21, 1-132.	2.8	41
172	Cost and Outcome of BehaviouRal Activation (COBRA): a randomised controlled trial of behavioural activation versus cognitive–behavioural therapy for depression. Health Technology Assessment, 2017, 21, 1-366.	2.8	24
173	CollAborative care for Screen-Positive EldeRs with major depression (CASPER plus): a multicentred randomised controlled trial of clinical effectiveness and cost-effectiveness. Health Technology Assessment, 2017, 21, 1-252.	2.8	56
174	Making the journey with me: a qualitative study of experiences of a bespoke mental health smoking cessation intervention for service users with serious mental illness. BMC Psychiatry, 2016, 16, 193.	2.6	16
175	Identifying depression with the PHQ-2: A diagnostic meta-analysis. Journal of Affective Disorders, 2016, 203, 382-395.	4.1	144
176	The diagnostic accuracy of brief versions of the Geriatric Depression Scale: a systematic review and metaâ€analysis. International Journal of Geriatric Psychiatry, 2016, 31, 837-857.	2.7	186
177	Identification of depression in women during pregnancy and the early postnatal period using the Whooley questions and the Edinburgh Postnatal Depression Scale: protocol for the Born and Bred in Yorkshire: PeriNatal Depression Diagnostic Accuracy (BaBY PaNDA) study. BMJ Open, 2016, 6, e011223.	1.9	16
178	Varenicline for smoking cessation and reduction in people with severe mental illnesses: systematic review and meta-analysis. Addiction, 2016, 111, 1554-1567.	3.3	28
179	Loneliness and social isolation as risk factors for coronary heart disease and stroke: systematic review and meta-analysis of longitudinal observational studies. Heart, 2016, 102, 1009-1016.	2.9	1,109
180	Loneliness, social isolation and social relationships: what are we measuring? A novel framework for classifying and comparing tools. BMJ Open, 2016, 6, e010799.	1.9	198

#	Article	IF	CITATIONS
181	Disentangling psychopathology, substance use and dependence: a factor analysis. BMC Psychiatry, 2016, 16, 281.	2.6	7
182	Diagnostic accuracy of the Depression subscale of the Hospital Anxiety and Depression Scale (HADS-D) for detecting major depression: protocol for a systematic review and individual patient data meta-analyses. BMJ Open, 2016, 6, e011913.	1.9	22
183	Infection in people with severe mental illness – Authors' reply. Lancet Psychiatry,the, 2016, 3, 204-205.	7.4	0
184	Evaluation of ethnic disparities in detection of depression and anxiety in primary care during the maternal period: Combined analysis of routine and cohort data. British Journal of Psychiatry, 2016, 208, 453-461.	2.8	46
185	Life adversity in depressed and non-depressed older adults: A cross-sectional comparison of the brief LTE-Q questionnaire and life events and difficulties interview as part of the CASPER study. Journal of Affective Disorders, 2016, 193, 31-38.	4.1	14
186	Risperidone versus other atypical antipsychotic medication for schizophrenia. The Cochrane Library, 2016, 9, CD002306.	2.8	2
187	Routine use of patient reported outcome measures (PROMs) for improving treatment of common mental health disorders in adults. The Cochrane Library, 2016, 2016, CD011119.	2.8	133
188	Cost and Outcome of Behavioural Activation versus Cognitive Behavioural Therapy for Depression (COBRA): a randomised, controlled, non-inferiority trial. Lancet, The, 2016, 388, 871-880.	13.7	427
189	Prevalence of HIV, hepatitis B, and hepatitis C in people with severe mental illness: a systematic review and meta-analysis. Lancet Psychiatry,the, 2016, 3, 40-48.	7.4	193
190	Association Between Chronic Physical Conditions and the Effectiveness of Collaborative Care for Depression. JAMA Psychiatry, 2016, 73, 978.	11.0	76
191	On poverty, politics and psychology: the socioeconomic gradient of mental healthcare utilisation and outcomes. British Journal of Psychiatry, 2016, 209, 429-430.	2.8	57
192	Variation and ethnic inequalities in treatment of common mental disorders before, during and after pregnancy: combined analysis of routine and research data in the Born in Bradford cohort. BMC Psychiatry, 2016, 16, 99.	2.6	30
193	Mirtazapine added to selective serotonin reuptake inhibitors for treatment-resistant depression in primary care (MIR trial): study protocol for a randomised controlled trial. Trials, 2016, 17, 66.	1.6	10
194	Exploring why people with <scp>SMI</scp> smoke and why they may want to quit: baseline data from the <scp>SCIMITAR RCT</scp> . Journal of Psychiatric and Mental Health Nursing, 2016, 23, 282-289.	2.1	14
195	Usefulness of PHQ-9 in primary care to determine meaningful symptoms of low mood: a qualitative study. British Journal of General Practice, 2016, 66, e78-e84.	1.4	33
196	Maternal psychological distress in primary care and association with child behavioural outcomes at age three. European Child and Adolescent Psychiatry, 2016, 25, 601-613.	4.7	13
197	Clinical effectiveness and cost-effectiveness of collaborative care for depression in UK primary care (CADET): a cluster randomised controlled trial. Health Technology Assessment, 2016, 20, 1-192.	2.8	41
198	The second Randomised Evaluation of the Effectiveness, cost-effectiveness and Acceptability of Computerised Therapy (REEACT-2) trial: does the provision of telephone support enhance the effectiveness of computer-delivered cognitive behaviour therapy? A randomised controlled trial. Health Technology Assessment, 2016, 20, 1-64.	2.8	31

#	Article	IF	CITATIONS
199	The Management of Depression and Anxiety in Older People: Evidence-Based Psychological Interventions, Stepped Care and Collaborative Care. , 2016, , 79-92.		1
200	The effectiveness of self-management support interventions for men with long-term conditions: a systematic review and meta-analysis. BMJ Open, 2015, 5, e006620-e006620.	1.9	43
201	Diagnostic accuracy of the Whooley questions for the identification of depression: a diagnostic meta-analysis. BMJ Open, 2015, 5, e008913.	1.9	66
202	Determinants of hospital length of stay for people with serious mental illness in England and implications for payment systems: a regression analysis. BMC Health Services Research, 2015, 15, 439.	2.2	40
203	Revealing hidden depression in older people: a qualitative study within a randomised controlled trial. BMC Family Practice, 2015, 16, 142.	2.9	21
204	Does the quality and outcomes framework reduce psychiatric admissions in people with serious mental illness? A regression analysis. BMJ Open, 2015, 5, e007342-e007342.	1.9	15
205	M9â€Lung Health of OPiate Users (LHOp): A pilot study to assess the respiratory health of opiate misusers attending a community substance misuse clinic. Thorax, 2015, 70, A230.2-A231.	5.6	0
206	Post-it notes to improve questionnaire response rates in RCTs - findings from a randomised sub-study. Trials, 2015, 16, .	1.6	0
207	The acceptability of case-finding questions to identify perinatal depression. British Journal of Midwifery, 2015, 23, 630-638.	0.4	5
208	Evidence-based guidelines for treating depressive disorders with antidepressants: A revision of the 2008 British Association for Psychopharmacology guidelines. Journal of Psychopharmacology, 2015, 29, 459-525.	4.0	528
209	Systematic review and meta-analysis of transdiagnostic psychological treatments for anxiety and depressive disorders in adulthood. Clinical Psychology Review, 2015, 40, 91-110.	11.4	411
210	Patient experience of computerised therapy for depression in primary care. BMJ Open, 2015, 5, e008581.	1.9	65
211	Computerised cognitive behaviour therapy (cCBT) as treatment for depression in primary care (REEACT) Tj ETQq1	1.0.7843 6.0	1418BT /C
212	Screening and case finding for major depressive disorder using the Patient Health Questionnaire (PHQ-9): a meta-analysis. General Hospital Psychiatry, 2015, 37, 567-576.	2.4	234
213	Diagnostic accuracy of the Edinburgh Postnatal Depression Scale (EPDS) for detecting major depression in pregnant and postnatal women: protocol for a systematic review and individual patient data meta-analyses. BMJ Open, 2015, 5, e009742.	1.9	46
214	A diagnostic meta-analysis of the Patient Health Questionnaire-9 (PHQ-9) algorithm scoring method as a screen for depression. General Hospital Psychiatry, 2015, 37, 67-75.	2.4	490
215	Diagnostic accuracy of the Whooley depression tool in older adults in UK primary care. Journal of Affective Disorders, 2015, 182, 39-43.	4.1	8
216	Feasibility Randomized Controlled Trial of Cognitive and Behavioral Interventions for Depression Symptoms in Patients Accessing Drug and Alcohol Treatment. Journal of Substance Abuse Treatment, 2015, 55, 6-14.	2.8	24

#	Article	IF	CITATIONS
217	Bespoke smoking cessation for people with severe mental ill health (SCIMITAR): a pilot randomised controlled trial. Lancet Psychiatry,the, 2015, 2, 395-402.	7.4	68
218	Do higher primary care practice performance scores predict lower rates of emergency admissions for persons with serious mental illness? An analysis of secondary panel data. Health Services and Delivery Research, 2015, 3, 1-68.	1.4	6
219	A systematic review and metaethnography to identify how effective, cost-effective, accessible and acceptable self-management support interventions are for men with long-term conditions (SELF-MAN). Health Services and Delivery Research, 2015, 3, 1-302.	1.4	6
220	Screening for psychological and mental health difficulties in young people who offend: a systematic review and decision model. Health Technology Assessment, 2015, 19, 1-128.	2.8	5
221	A randomised controlled trial of computerised cognitive behaviour therapy for the treatment of depression in primary care: the Randomised Evaluation of the Effectiveness and Acceptability of Computerised Therapy (REEACT) trial. Health Technology Assessment, 2015, 19, 1-174.	2.8	55
222	Smoking Cessation Intervention for severe Mental III Health Trial (SCIMITAR): a pilot randomised control trial of the clinical effectiveness and cost-effectiveness of a bespoke smoking cessation service. Health Technology Assessment, 2015, 19, 1-148.	2.8	28
223	Clinical effectiveness and cost-effectiveness of parenting interventions for children with severe attachment problems: a systematic review and meta-analysis. Health Technology Assessment, 2015, 19, 1-348.	2.8	32
224	Heterogeneity in Patient-Reported Outcomes following Low-Intensity Mental Health Interventions: A Multilevel Analysis. PLoS ONE, 2014, 9, e99658.	2.5	16
225	Characteristics of Effective Collaborative Care for Treatment of Depression: A Systematic Review and Meta-Regression of 74 Randomised Controlled Trials. PLoS ONE, 2014, 9, e108114.	2.5	158
226	The accessibility and acceptability of self-management support interventions for men with long term conditions: a systematic review and meta-synthesis of qualitative studies. BMC Public Health, 2014, 14, 1230.	2.9	61
227	The diagnostic accuracy of the Patient Health Questionnaire-2 (PHQ-2), Patient Health Questionnaire-8 (PHQ-8), and Patient Health Questionnaire-9 (PHQ-9) for detecting major depression: protocol for a systematic review and individual patient data meta-analyses. Systematic Reviews, 2014, 3, 124.	5.3	71
228	Computerised cognitive behaviour therapy for depression in adolescents: study protocol for a feasibility randomised controlled trial. BMJ Open, 2014, 4, e006488.	1.9	16
229	Promoting <scp>HIV</scp> and sexual safety behaviour in people with severe mental illness: A systematic review of behavioural interventions. International Journal of Mental Health Nursing, 2014, 23, 344-354.	3.8	27
230	CASPER plus (CollAborative care in Screen-Positive EldeRs with major depressive disorder): study protocol for a randomised controlled trial. Trials, 2014, 15, 451.	1.6	16
231	Benchmarking Routine Psychological Services: A Discussion of Challenges and Methods. Behavioural and Cognitive Psychotherapy, 2014, 42, 16-30.	1.2	24
232	Adapting manualized Behavioural Activation treatment for older adults with depression. The Cognitive Behaviour Therapist, 2014, 7, .	1.0	21
233	Cost-effectiveness of combining systematic identification and treatment of co-morbid major depression for people with chronic diseases: the example of cancer. Psychological Medicine, 2014, 44, 1451-1460.	4.5	30
234	Depression screening and patient outcomes in pregnancy or postpartum: A systematic review. Journal of Psychosomatic Research, 2014, 76, 433-446.	2.6	88

#	Article	IF	CITATIONS
235	Cost and outcome of behavioural activation versus cognitive behaviour therapy for depression (COBRA): study protocol for a randomised controlled trial. Trials, 2014, 15, 29.	1.6	25
236	Early changes, attrition, and dose–response in low intensity psychological interventions. British Journal of Clinical Psychology, 2014, 53, 114-130.	3.5	87
237	Computerised Cognitive Behaviour Therapy for Depression Management: A Cost-Effectiveness Analysis. Value in Health, 2014, 17, A459.	0.3	0
238	Sudden gains in behavioural activation for depression. Behaviour Research and Therapy, 2014, 60, 34-38.	3.1	21
239	Obsessive Compulsive Treatment Efficacy Trial (OCTET) comparing the clinical and cost effectiveness of self-managed therapies: study protocol for a randomised controlled trial. Trials, 2014, 15, 278.	1.6	10
240	Qualitative Meta-Synthesis of User Experience of Computerised Therapy for Depression and Anxiety. PLoS ONE, 2014, 9, e84323.	2.5	148
241	Behavioural Activation for Depression; An Update of Meta-Analysis of Effectiveness and Sub Group Analysis. PLoS ONE, 2014, 9, e100100.	2.5	390
242	Cost-Effectiveness of Collaborative Care for Depression in UK Primary Care: Economic Evaluation of a Randomised Controlled Trial (CADET). PLoS ONE, 2014, 9, e104225.	2.5	38
243	A systematic review, psychometric analysis and qualitative assessment of generic preference-based measures of health in mental health populations and the estimation of mapping functions from widely used specific measures. Health Technology Assessment, 2014, 18, vii-viii, xiii-xxv, 1-188.	2.8	179
244	Assessing the risk of self-harm in an adult offender population: an incidence cohort study. Health Technology Assessment, 2014, 18, 1-152.	2.8	10
245	The psychometric properties of the subscales of the GHQ-28 in a multi-ethnic maternal sample: results from the Born in Bradford cohort. BMC Psychiatry, 2013, 13, 55.	2.6	20
246	Cost-Effectiveness of the Systematic Identification and Treatment of Comorbid Major Depression for People with Chronic Diseases: The Example of Cancer. Value in Health, 2013, 16, A414.	0.3	0
247	Nurse-delivered collaborative care for depression and long-term physical conditions: A systematic review and meta-analysis. Journal of Affective Disorders, 2013, 149, 14-22.	4.1	88
248	Psychometric properties of the Treatment Outcomes Profile (TOP) psychological health scale. Mental Health and Substance Use: Dual Diagnosis, 2013, 6, 140-149.	0.5	6
249	Decision curve analysis for assessing the usefulness of tests for making decisions to treat: An application to tests for prodromal psychosis Psychological Assessment, 2013, 25, 730-737.	1.5	7
250	Clinical effectiveness of collaborative care for depression in UK primary care (CADET): cluster randomised controlled trial. BMJ, The, 2013, 347, f4913-f4913.	6.0	173
251	Influence of initial severity of depression on effectiveness of low intensity interventions: meta-analysis of individual patient data. BMJ, The, 2013, 346, f540-f540.	6.0	251
252	Depression, anxiety and comorbid substance use: association patterns in outpatient addictions treatment. Mental Health and Substance Use: Dual Diagnosis, 2013, 6, 59-75.	0.5	10

#	Article	IF	CITATIONS
253	The COREâ€10: A short measure of psychological distress for routine use in the psychological therapies. Counselling and Psychotherapy Research, 2013, 13, 3-13.	3.2	229
254	Psychological Distress during Pregnancy in a Multi-Ethnic Community: Findings from the Born in Bradford Cohort Study. PLoS ONE, 2013, 8, e60693.	2.5	34
255	Rethinking recommendations for screening for depression in primary care. Cmaj, 2012, 184, 413-418.	2.0	184
256	Diagnostic accuracy of case-finding questions to identify perinatal depression. Cmaj, 2012, 184, E424-E430.	2.0	32
257	Collaborative care for depression and anxiety problems. The Cochrane Library, 2012, 10, CD006525.	2.8	624
258	Re-rethinking the article by Thombs and colleagues: Figure 1:. Cmaj, 2012, 184, 438.2-439.	2.0	2
259	Optimal cut-off score for diagnosing depression with the Patient Health Questionnaire (PHQ-9): a meta-analysis. Cmaj, 2012, 184, E191-E196.	2.0	1,336
260	Better care for depression in the workplace: Integrating occupational and mental health services. British Journal of Psychiatry, 2012, 200, 442-443.	2.8	13
261	Choosing a second generation antidepressant for treatment of major depressive disorder. BMJ: British Medical Journal, 2012, 344, e1014-e1014.	2.3	5
262	Brief case finding tools for anxiety disorders: Validation of GAD-7 and GAD-2 in addictions treatment. Drug and Alcohol Dependence, 2012, 125, 37-42.	3.2	66
263	Delivering stepped care: an analysis of implementation in routine practice. Implementation Science, 2012, 7, 3.	6.9	120
264	Acceptability of mental health screening in routine addictions treatment. General Hospital Psychiatry, 2012, 34, 415-422.	2.4	9
265	Should methodological filters for diagnostic test accuracy studies be used in systematic reviews of psychometric instruments? a case study involving screening for postnatal depression. Systematic Reviews, 2012, 1, 9.	5.3	8
266	How reliable is depression screening in alcohol and drug users? A validation of brief and ultra-brief questionnaires. Journal of Affective Disorders, 2011, 134, 266-271.	4.1	49
267	P1-294 Risk factors for poor mental health during pregnancy from the born in Bradford cohort: early results from a bi-ethnic sample. Journal of Epidemiology and Community Health, 2011, 65, A148-A148.	3.7	Ο
268	P1-496 An examination of the psychometric properties of the GHQ-28 subscales in a bi-ethnic antenatal sample in the UK: preliminary results from the born in Bradford cohort. Journal of Epidemiology and Community Health, 2011, 65, A204-A204.	3.7	0
269	Behavioural activation delivered by the non-specialist: phase II randomised controlled trial. British Journal of Psychiatry, 2011, 198, 66-72.	2.8	178
270	Cost utility of behavioural activation delivered by the non-specialist. British Journal of Psychiatry, 2011, 199, 510-511.	2.8	44

#	Article	IF	CITATIONS
271	Screening and case finding for depression in offender populations: A systematic review of diagnostic properties. Journal of Affective Disorders, 2011, 128, 72-82.	4.1	19
272	Validity of two case finding questions to detect postnatal depression: A review of diagnostic test accuracy. Journal of Affective Disorders, 2011, 133, 388-397.	4.1	15
273	A randomised evaluation of CollAborative care and active surveillance for Screen-Positive EldeRs with sub-threshold depression (CASPER): study protocol for a randomized controlled trial. Trials, 2011, 12, 225.	1.6	27
274	Can we improve recruitment to trials and informed consent by improving participant information sheets? - A nested RCT. Trials, 2011, 12, .	1.6	2
275	Recruiting ahead of target: What worked in the REEACT trial?. Trials, 2011, 12, .	1.6	2
276	The effectiveness of behavioural therapy for the treatment of depression in older adults: a metaâ€analysis. International Journal of Geriatric Psychiatry, 2011, 26, 1211-1220.	2.7	47
277	Prevalence and incidence of postnatal depression: what can systematic reviews tell us?. Archives of Women's Mental Health, 2010, 13, 295-305.	2.6	44
278	Improving health and productivity of depressed workers: a pilot randomized controlled trial of telephone cognitive behavioral therapy delivery in workplace settings. General Hospital Psychiatry, 2010, 32, 337-340.	2.4	27
279	Defining successful treatment outcome in depression using the PHQ-9: A comparison of methods. Journal of Affective Disorders, 2010, 127, 122-129.	4.1	179
280	What work has to be done to implement collaborative care for depression? Process evaluation of a trial utilizing the Normalization Process Model. Implementation Science, 2010, 5, 15.	6.9	53
281	Translating research into practice in Leeds and Bradford (TRiPLaB): a protocol for a programme of research. Implementation Science, 2010, 5, 37.	6.9	19
282	Smoking cessation in severe mental illness: what works?. Addiction, 2010, 105, 1176-1189.	3.3	208
283	Effectiveness of interventions designed to promote patient involvement to enhance safety: a systematic review. BMJ Quality and Safety, 2010, 19, e10-e10.	3.7	72
284	Instruments to identify post-natal depression: Which methods have been the most extensively validated, in what setting and in which language?. International Journal of Psychiatry in Clinical Practice, 2010, 14, 72-76.	2.4	64
285	Screening for postnatal depression – is it acceptable to women and healthcare professionals? A systematic review and metaâ€synthesis. Journal of Reproductive and Infant Psychology, 2010, 28, 328-344.	1.8	66
286	The Hospital Anxiety and Depression Scale: A diagnostic meta-analysis of case-finding ability. Journal of Psychosomatic Research, 2010, 69, 371-378.	2.6	285
287	Depression in primary care. , 2010, , 1-16.		2
288	User-defined outcomes in mental health: A qualitative study and consensus development exercise. Journal of Mental Health, 2009, 18, 415-423.	1.9	5

#	Article	IF	CITATIONS
289	A stepped care intervention is effective for elderly people with subthreshold depression or anxiety. Evidence-Based Mental Health, 2009, 12, 115-115.	4.5	0
290	Collaborative Depression Trial (CADET): multi-centre randomised controlled trial of collaborative care for depression - study protocol. BMC Health Services Research, 2009, 9, 188.	2.2	30
291	Assessing the quality of diagnostic studies using psychometric instruments: applying QUADAS. Social Psychiatry and Psychiatric Epidemiology, 2009, 44, 300-307.	3.1	26
292	Putting the â€~Q' in depression QALYs: a comparison of utility measurement using EQ-5D and SF-6D health related quality of life measures. Social Psychiatry and Psychiatric Epidemiology, 2009, 44, 569-578.	3.1	34
293	Detecting and predicting self-harm behaviour in prisoners: a prospective psychometric analysis of three instruments. Social Psychiatry and Psychiatric Epidemiology, 2009, 44, 853-861.	3.1	25
294	Is it clinically and cost effective to screen for postnatal depression: a systematic review of controlled clinical trials and economic evidence. BJOG: an International Journal of Obstetrics and Gynaecology, 2009, 116, 1019-1027.	2.3	53
295	Barriers to the uptake of computerized cognitive behavioural therapy: a systematic review of the quantitative and qualitative evidence. Psychological Medicine, 2009, 39, 705-712.	4.5	355
296	Collaborative care for depression in UK primary care: a randomized controlled trial – Corrigendum. Psychological Medicine, 2009, 39, 701-701.	4.5	3
297	Screening for postnatal depression in primary care: cost effectiveness analysis. BMJ: British Medical Journal, 2009, 339, b5203-b5203.	2.3	100
298	Methods to identify postnatal depression in primary care: an integrated evidence synthesis and value of information analysis. Health Technology Assessment, 2009, 13, 1-145, 147-230.	2.8	212
299	A review and critical appraisal of measures of therapist-patient interactions in mental health settings. Clinical Governance, 2009, 14, .	0.3	0
300	Psychotherapy mediated by remote communication technologies: a meta-analytic review. BMC Psychiatry, 2008, 8, 60.	2.6	151
301	Cluster randomized trials produced similar results to individually randomized trials in a meta-analysis of enhanced care for depression. Journal of Clinical Epidemiology, 2008, 61, 160-168.e2.	5.0	24
302	Telephone treatment support improves outcomes for depressed employees. Evidence-Based Mental Health, 2008, 11, 47-47.	4.5	1
303	Screening and case-finding instruments for depression: a meta-analysis. Cmaj, 2008, 178, 997-1003.	2.0	371
304	A meta-analysis of randomized trials of behavioural treatment of depression. Psychological Medicine, 2008, 38, 611-623.	4.5	324
305	Depression in older adults: collaborative care model seems effective. Evidence-Based Mental Health, 2008, 11, 44-44.	4.5	4
306	The European pharmaceutical industry: coming to a screen near you?. Journal of the Royal Society of Medicine, 2008, 101, 485-488.	2.0	1

#	Article	IF	CITATIONS
307	Collaborative care for depression in UK primary care: a randomized controlled trial. Psychological Medicine, 2008, 38, 279-287.	4.5	135
308	A review and critical appraisal of measures of therapist-patient interactions in mental health settings. Health Technology Assessment, 2008, 12, iii, ix-47.	2.8	36
309	A smoking ban in psychiatric units: threat or opportunity?. Mental Health in Family Medicine, 2008, 5, 129-33.	0.2	0
310	Can we predict suicide and non-fatal self-harm with the Beck Hopelessness Scale? A meta-analysis. Psychological Medicine, 2007, 37, 769-778.	4.5	206
311	Is low folate a risk factor for depression? A meta-analysis and exploration of heterogeneity. Journal of Epidemiology and Community Health, 2007, 61, 631-637.	3.7	197
312	IMPACT collaborative care programme reduces suicide ideation in depressed older adults. Evidence-Based Mental Health, 2007, 10, 51-51.	4.5	5
313	What makes self-help interventions effective in the management of depressive symptoms? Meta-analysis and meta-regression. Psychological Medicine, 2007, 37, 1217-1228.	4.5	387
314	Vitamin K and the Prevention of Fractures—Reply. Archives of Internal Medicine, 2007, 167, 94.	3.8	1
315	Clarification of Study and Citation—Reply. Archives of Internal Medicine, 2007, 167, 2531.	3.8	0
316	Randomized trials with concurrent economic evaluations reported unrepresentatively large clinical effect sizes. Journal of Clinical Epidemiology, 2007, 60, 781.e1-781.e10.	5.0	11
317	Screening for Depression in Medical Settings with the Patient Health Questionnaire (PHQ): A Diagnostic Meta-Analysis. Journal of General Internal Medicine, 2007, 22, 1596-1602.	2.6	991
318	A systematic review of quantitative and qualitative research on the role and effectiveness of written information available to patients about individual medicines. Health Technology Assessment, 2007, 11, iii, 1-160.	2.8	221
319	A systematic review and economic model of the clinical effectiveness and cost-effectiveness of interventions for preventing relapse in people with bipolar disorder. Health Technology Assessment, 2007, 11, iii-iv, ix-206.	2.8	83
320	Diagnosing depression in primary care using self-completed instruments: UK validation of PHQ-9 and CORE-OM. British Journal of General Practice, 2007, 57, 650-2.	1.4	136
321	Can we improve the morale of staff working in psychiatric units? A systematic review. Journal of Mental Health, 2006, 15, 7-17.	1.9	49
322	Collaborative care for depression in primary care. British Journal of Psychiatry, 2006, 189, 484-493.	2.8	343
323	The prevalence of nursing staff stress on adult acute psychiatric in-patient wards. Social Psychiatry and Psychiatric Epidemiology, 2006, 41, 34-43.	3.1	36
324	Developing a U.K. protocol for collaborative care: a qualitative study. General Hospital Psychiatry, 2006, 28, 296-305.	2.4	60

#	Article	IF	CITATIONS
325	Should we screen for depression?. BMJ: British Medical Journal, 2006, 332, 1027-1030.	2.3	172
326	Collaborative Care for Depression. Archives of Internal Medicine, 2006, 166, 2314.	3.8	1,141
327	Vitamin K and the Prevention of Fractures. Archives of Internal Medicine, 2006, 166, 1256.	3.8	317
328	Methylenetetrahydrofolate Reductase (MTHFR) Genetic Polymorphisms and Psychiatric Disorders: A HuGE Review. American Journal of Epidemiology, 2006, 165, 1-13.	3.4	290
329	Costs and consequences of enhanced primary care for depression. British Journal of Psychiatry, 2006, 189, 297-308.	2.8	150
330	NICE, but will they help people with depression? The new National Institute for Clinical Excellence depression guidelines. British Journal of Psychiatry, 2005, 186, 177-178.	2.8	28
331	Managing common mental health disorders in primary care: conceptual models and evidence base. BMJ: British Medical Journal, 2005, 330, 839-842.	2.3	184
332	Comparison of Large Versus Smaller Randomized Trials for Mental Health-Related Interventions. American Journal of Psychiatry, 2005, 162, 578-584.	7.2	32
333	Stepped care in psychological therapies: access, effectiveness and efficiency. British Journal of Psychiatry, 2005, 186, 11-17.	2.8	818
334	Screening and case finding instruments for depression. The Cochrane Library, 2005, , CD002792.	2.8	133
335	Benefits and harms of direct to consumer advertising: a systematic review. Quality and Safety in Health Care, 2005, 14, 246-250.	2.5	179
336	Direct-to-consumer advertising of psychotropics. British Journal of Psychiatry, 2004, 185, 1-2.	2.8	14
337	Review: disease management programmes improve detection and care of people with depression. Evidence-Based Mental Health, 2004, 7, 80-80.	4.5	11
338	Educational and organisational interventions can improve the management of depression in primary care. Evidence-Based Healthcare and Public Health, 2004, 8, 6-7.	0.0	0
339	A systematic review of risperidone for schizophrenia. Schizophrenia Research, 2003, 60, 286.	2.0	Ο
340	Improving the detection and management of depression in primary care. Quality and Safety in Health Care, 2003, 12, 149-155.	2.5	85
341	Review: complex organisational and educational interventions appear to be effective for managing depression in primary care. Evidence-Based Medicine, 2003, 8, 191-191.	0.6	1
342	Educational and Organizational Interventions to Improve the Management of Depression in Primary Care. JAMA - Journal of the American Medical Association, 2003, 289, 3145.	7.4	649

#	Article	IF	CITATIONS
343	Outcome measures and needs assessment tools for schizophrenia and related disorders. The Cochrane Library, 2003, , CD003081.	2.8	28
344	4 factors were predictive of extended duration in hospital in acute psychiatric admission. Evidence-Based Mental Health, 2002, 5, 64-64.	4.5	1
345	Routine administration of Health Related Quality of Life (HRQoL) and needs assessment instruments to improve psychological outcome–Âa systematic review. Psychological Medicine, 2002, 32, 1345-1356.	4.5	86
346	Outcomes research in mental health. British Journal of Psychiatry, 2002, 181, 8-16.	2.8	85
347	Newer atypical antipsychotic medication in comparison to clozapine: a systematic review of randomized trials. Schizophrenia Research, 2002, 56, 1-10.	2.0	93
348	Improving the delivery and organisation of mental health services: Beyond the conventional randomised controlled trial. British Journal of Psychiatry, 2002, 180, 13-18.	2.8	48
349	Psychiatrists in the UK do not use outcomes measures. British Journal of Psychiatry, 2002, 180, 101-103.	2.8	194
350	Randomized controlled trials in schizophrenia: a critical perspective on the literature. Acta Psychiatrica Scandinavica, 2002, 105, 243-251.	4.5	48
351	Routinely administered questionnaires for depression and anxiety: systematic review. BMJ: British Medical Journal, 2001, 322, 406-409.	2.3	272
352	"Nobody ever expects the Spanish Inquisition―(Python, 1991). Psychiatric Bulletin, 2001, 25, 291-292.	0.3	7
353	Risperidone versus other atypical antipsychotic medication for schizophrenia. , 2000, , CD002306.		34
354	Publication bias and the integrity of psychiatry research. Psychological Medicine, 2000, 30, 253-258.	4.5	59
355	The causes, consequences and detection of publication bias in psychiatry. Acta Psychiatrica Scandinavica, 2000, 102, 241-249.	4.5	85
356	Influence of Methodology on Outcomes of Randomised Clozapine Trials. Pharmacopsychiatry, 2000, 33, 54-59.	3.3	13
357	Drug treatments for schizophrenia. Quality in Health Care: QHC, 2000, 9, 73-79.	1.2	7
358	Efficacy of β-blocker supplementation for schizophrenia: a systematic review of randomized trials. Schizophrenia Research, 2000, 41, 341-347.	2.0	15
359	Modelling publication bias in meta-analysis: a review. Statistical Methods in Medical Research, 2000, 9, 421-445.	1.5	86
360	Publication and related biases Health Technology Assessment, 2000, 4, .	2.8	315

#	Article	IF	CITATIONS
361	Variations in psychiatric practice. British Journal of Psychiatry, 1999, 175, 303-305.	2.8	13
362	Lost information? The fate of papers presented at the 40th society for Social Medicine Conference. Journal of Epidemiology and Community Health, 1999, 53, 442-443.	3.7	14
363	The role of electronic journals in reducing publication bias. Informatics for Health and Social Care, 1999, 24, 223-229.	1.0	18
364	Vegetarianism in young women: Another means of weight control?. International Journal of Eating Disorders, 1999, 26, 87-90.	4.0	53
365	Relation between hostility and coronary heart disease. BMJ: British Medical Journal, 1999, 319, 917-917.	2.3	22
366	NICE work?. Psychiatric Bulletin, 1999, 23, 513-516.	0.3	2
367	Mental health promotion in high risk groups Quality and Safety in Health Care, 1997, 6, 219-225.	2.5	7
368	Rational Decision Making in Psychiatry: Evidence-Based Psychiatry is Just the Start. British Journal of Psychiatry, 1997, 171, 226-227.	2.8	3
369	Prescribing antidepressants in general practice. BMJ: British Medical Journal, 1997, 314, 826-826.	2.3	8
370	â€~Number needed to treat'. British Journal of Psychiatry, 1996, 169, 113-113.	2.8	1
371	Deliberate self harm Emergency Medicine Journal, 1996, 13, 432-432.	1.0	1
372	Differences in use of abbreviated mental test score by geriatricians and psychiatrists. BMJ: British Medical Journal, 1996, 313, 465-465.	2.3	17
373	Publication bias and meta-analysis. British Journal of Psychiatry, 1995, 167, 266-266.	2.8	5
374	Subject to many potential biases. BMJ: British Medical Journal, 1995, 311, 1303-1304.	2.3	3
375	Linking evidence to practice. , 0, , 31-54.		Ο
376	Improving Access to Care. , 0, , 386-400.		0
377	Collaborative Care and Stepped Care. , 0, , 401-419.		0
378	Prognostic models for predicting relapse or recurrence of depression. The Cochrane Library, 0, , .	2.8	1

0

#	Article	IF	CITATIONS
379	Factors that influence participation in physical activity for people with bipolar disorder: a synthesis of qualitative evidence. The Cochrane Library, 0, , .	2.8	1
380	Cognitive or behavioural interventions (or both) to prevent or mitigate loneliness in adolescents, adults, and older adults. The Cochrane Library, 0, , .	2.8	8
381	Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, may reduce social isolation and loneliness among older adults and how?. F1000Research, 0, 9, 1368.	1.6	19
382	Home hazard assessment and environmental modification to prevent falls in older people: the OTIS trial. F1000Research, 0, 10, 500.	1.6	2
383	Pharmacy-based management for depression in adults. The Cochrane Library, 0, , .	2.8	3
384	United Kingdom perspective. , 0, , 127-136.		2
385	Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work?. F1000Research, 0, 9, 1368.	1.6	7
386	Indicators of Prognosis Independent of Treatment for Adults with Depression in Primary Care, Going Beyond Baseline Symptom-Severity: A Systematic Review and Individual Patient Data Meta-Analysis. SSRN Electronic Journal, 0, , .	0.4	2
387	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. Wellcome Open Research, 0, 4, 69.	1.8	1
388	Models of care for depression. , 0, , 17-30.		0
389	Anatomy of a review. , 0, , 55-72.		0
390	Collaborative care. , 0, , 87-102.		0
391	Referral. , 0, , 103-110.		0
392	Summary of the evidence. , 0, , 111-116.		0
393	Making it happen. , 0, , 117-126.		0

United States perspective. , 0, , 137-144.