

# Albrecht E Melchinger

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6389567/publications.pdf>

Version: 2024-02-01

315  
papers

18,992  
citations

9786

73  
h-index

20961

115  
g-index

319  
all docs

319  
docs citations

319  
times ranked

10041  
citing authors

#	ARTICLE	IF	CITATIONS
1	High-resolution association mapping with libraries of immortalized lines from ancestral landraces. Theoretical and Applied Genetics, 2022, 135, 243-256.	3.6	5
2	Unraveling the potential of phenomic selection within and among diverse breeding material of maize (<i>Zea mays</i> L.). G3: Genes, Genomes, Genetics, 2022, 12, .	1.8	9
3	Genetic diversity of European maize landraces: Dataset on the molecular and phenotypic variation of derived doubled-haploid populations. Data in Brief, 2022, 42, 108164.	1.0	4
4	Theoretical and experimental assessment of genome-based prediction in landraces of allogamous crops. Proceedings of the National Academy of Sciences of the United States of America, 2022, 119, e2121797119.	7.1	4
5	Exploiting genetic diversity in two European maize landraces for improving Gibberella ear rot resistance using genomic tools. Theoretical and Applied Genetics, 2021, 134, 793-805.	3.6	18
6	Calibration and validation of predicted genomic breeding values in an advanced cycle maize population. Theoretical and Applied Genetics, 2021, 134, 3069-3081.	3.6	12
7	Optimum breeding strategies using genomic and phenotypic selection for the simultaneous improvement of two traits. Theoretical and Applied Genetics, 2021, 134, 4025-4042.	3.6	5
8	Genomic prediction with multiple biparental families. Theoretical and Applied Genetics, 2020, 133, 133-147.	3.6	22
9	Genetic dissection of maternal influence on in vivo haploid induction in maize. Crop Journal, 2020, 8, 287-298.	5.2	5
10	Discovery of beneficial haplotypes for complex traits in maize landraces. Nature Communications, 2020, 11, 4954.	12.8	38
11	Genome-wide association study to identify genomic regions influencing spontaneous fertility in maize haploids. Euphytica, 2019, 215, 138.	1.2	29
12	European maize landraces made accessible for plant breeding and genome-based studies. Theoretical and Applied Genetics, 2019, 132, 3333-3345.	3.6	52
13	Doubled haploid technology for line development in maize: technical advances and prospects. Theoretical and Applied Genetics, 2019, 132, 3227-3243.	3.6	126
14	Efficient genetic value prediction using incomplete omics data. Theoretical and Applied Genetics, 2019, 132, 1211-1222.	3.6	8
15	Reduced response diversity does not negatively impact wheat climate resilience. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116, 10623-10624.	7.1	11
16	Haploid male fertility and spontaneous chromosome doubling evaluated in a diallel and recurrent selection experiment in maize. Theoretical and Applied Genetics, 2019, 132, 2273-2284.	3.6	17
17	Testcross performance of doubled haploid lines from European flint maize landraces is promising for broadening the genetic base of elite germplasm. Theoretical and Applied Genetics, 2019, 132, 1897-1908.	3.6	28
18	Early diagnosis of ploidy status in doubled haploid production of maize by stomata length and flow cytometry measurements. Plant Breeding, 2019, 138, 266-276.	1.9	13

#	ARTICLE	IF	CITATIONS
19	Progress for testcross performance within the flint heterotic pool of a public maize breeding program since the onset of hybrid breeding. <i>Euphytica</i> , 2019, 215, 1.	1.2	9
20	Across-years prediction of hybrid performance in maize using genomics. <i>Theoretical and Applied Genetics</i> , 2019, 132, 933-946.	3.6	23
21	Beyond Genomic Prediction: Combining Different Types of <i>omics</i> Data Can Improve Prediction of Hybrid Performance in Maize. <i>Genetics</i> , 2018, 208, 1373-1385.	2.9	130
22	Selection on Expected Maximum Haploid Breeding Values Can Increase Genetic Gain in Recurrent Genomic Selection. <i>G3: Genes, Genomes, Genetics</i> , 2018, 8, 1173-1181.	1.8	42
23	Genomic prediction and GWAS of <i>Gibberella</i> ear rot resistance traits in dent and flint lines of a public maize breeding program. <i>Euphytica</i> , 2018, 214, 1.	1.2	32
24	Genomic Prediction Within and Among Doubled-Haploid Libraries from Maize Landraces. <i>Genetics</i> , 2018, 210, 1185-1196.	2.9	18
25	Marker-Assisted Breeding of Improved Maternal Haploid Inducers in Maize for the Tropical/Subtropical Regions. <i>Frontiers in Plant Science</i> , 2018, 9, 1527.	3.6	41
26	Small RNA-based prediction of hybrid performance in maize. <i>BMC Genomics</i> , 2018, 19, 371.	2.8	24
27	High-Throughput Precision Phenotyping of the Oil Content of Single Seeds of Various Oilseed Crops. <i>Crop Science</i> , 2018, 58, 670-678.	1.8	22
28	Nitrous Oxide-Induced Chromosome Doubling of Maize Haploids. <i>Crop Science</i> , 2018, 58, 650-659.	1.8	17
29	Parental Expression Variation of Small RNAs Is Negatively Correlated with Grain Yield Heterosis in a Maize Breeding Population. <i>Frontiers in Plant Science</i> , 2018, 9, 13.	3.6	21
30	Metabolic robustness in young roots underpins a predictive model of maize hybrid performance in the field. <i>Plant Journal</i> , 2017, 90, 319-329.	5.7	30
31	Tapping the genetic diversity of landraces in allogamous crops with doubled haploid lines: a case study from European flint maize. <i>Theoretical and Applied Genetics</i> , 2017, 130, 861-873.	3.6	41
32	Transcriptome-based prediction of hybrid performance with unbalanced data from a maize breeding programme. <i>Plant Breeding</i> , 2017, 136, 331-337.	1.9	22
33	Safeguarding Our Genetic Resources with Libraries of Doubled-Haploid Lines. <i>Genetics</i> , 2017, 206, 1611-1619.	2.9	24
34	Dissection of a major QTL <i>qhir1</i> conferring maternal haploid induction ability in maize. <i>Theoretical and Applied Genetics</i> , 2017, 130, 1113-1122.	3.6	20
35	Accuracy of Genomic Prediction in Synthetic Populations Depending on the Number of Parents, Relatedness, and Ancestral Linkage Disequilibrium. <i>Genetics</i> , 2017, 205, 441-454.	2.9	52
36	Genomic Prediction Within and Across Biparental Families: Means and Variances of Prediction Accuracy and Usefulness of Deterministic Equations. <i>G3: Genes, Genomes, Genetics</i> , 2017, 7, 3571-3586.	1.8	34

#	ARTICLE	IF	CITATIONS
37	Persistency of Prediction Accuracy and Genetic Gain in Synthetic Populations Under Recurrent Genomic Selection. <i>G3: Genes, Genomes, Genetics</i> , 2017, 7, 801-811.	1.8	37
38	Genotyping-by-sequencing highlights original diversity patterns within a European collection of 1191 maize flint lines, as compared to the maize USDA genebank. <i>Theoretical and Applied Genetics</i> , 2017, 130, 2165-2189.	3.6	39
39	High-throughput platform for automated sorting and selection of single seeds based on time-domain nuclear magnetic resonance (TD-NMR) measurement of oil content. <i>Biosystems Engineering</i> , 2017, 164, 213-220.	4.3	21
40	Omics-based hybrid prediction in maize. <i>Theoretical and Applied Genetics</i> , 2017, 130, 1927-1939.	3.6	90
41	Low validation rate of quantitative trait loci for <i>Gibberella</i> ear rot resistance in European maize. <i>Theoretical and Applied Genetics</i> , 2017, 130, 175-186.	3.6	20
42	Optimum breeding strategies using genomic selection for hybrid breeding in wheat, maize, rye, barley, rice and triticale. <i>Theoretical and Applied Genetics</i> , 2016, 129, 1901-1913.	3.6	69
43	Colchicine Alternatives for Chromosome Doubling in Maize Haploids for Doubled-Haploid Production. <i>Crop Science</i> , 2016, 56, 559-569.	1.8	47
44	Development and Validation of Red Root Marker-Based Haploid Inducers in Maize. <i>Crop Science</i> , 2016, 56, 1678-1688.	1.8	50
45	In Vivo Haploid Induction in Maize: Comparison of Different Testing Regimes for Measuring Haploid Induction Rates. <i>Crop Science</i> , 2016, 56, 1127-1135.	1.8	15
46	Association mapping for cold tolerance in two large maize inbred panels. <i>BMC Plant Biology</i> , 2016, 16, 127.	3.6	73
47	Domestication and Breeding of <i>Jatropha curcas</i> L. <i>Trends in Plant Science</i> , 2016, 21, 1045-1057.	8.8	84
48	Silage yield and quality traits in elite maize hybrids and their relationship to elemental concentrations in juvenile plants. <i>Plant Breeding</i> , 2016, 135, 55-62.	1.9	5
49	Prediction of hybrid performance in maize with a ridge regression model employed to DNA markers and mRNA transcription profiles. <i>BMC Genomics</i> , 2016, 17, 262.	2.8	28
50	Selectiongain: an R package for optimizing multi-stage selection. <i>Computational Statistics</i> , 2016, 31, 533-543.	1.5	5
51	The Genetic Basis of Haploid Induction in Maize Identified with a Novel Genome-Wide Association Method. <i>Genetics</i> , 2016, 202, 1267-1276.	2.9	61
52	Choice of models for QTL mapping with multiple families and design of the training set for prediction of <i>Fusarium</i> resistance traits in maize. <i>Theoretical and Applied Genetics</i> , 2016, 129, 431-444.	3.6	30
53	Controlling Misclassification Rates in Identification of Haploid Seeds from Induction Crosses in Maize with High-Oil Inducers. <i>Crop Science</i> , 2015, 55, 1076-1086.	1.8	10
54	Prediction of deoxynivalenol and zearalenone concentrations in <i>Fusarium graminearum</i> inoculated backcross populations of maize by symptom rating and near-infrared spectroscopy. <i>Plant Breeding</i> , 2015, 134, 529-534.	1.9	15

#	ARTICLE	IF	CITATIONS
55	Genomic selection in biparental populations: assessment of parameters for optimum estimation set design. <i>Plant Breeding</i> , 2015, 134, 623-630.	1.9	22
56	Oil Content is Superior to Oil Mass for Identification of Haploid Seeds in Maize Produced with Highâ€Oil Inducers. <i>Crop Science</i> , 2015, 55, 188-195.	1.8	23
57	Genetic diversity of Sudanese pearl millet ( <i>Pennisetum glaucum</i> (L.) R. Br.) landraces as revealed by SSR markers, and relationship between genetic and agro-morphological diversity. <i>Genetic Resources and Crop Evolution</i> , 2015, 62, 579-591.	1.6	32
58	Forecasting the accuracy of genomic prediction with different selection targets in the training and prediction set as well as truncation selection. <i>Theoretical and Applied Genetics</i> , 2015, 128, 2189-2201.	3.6	7
59	Shrinkage estimation of the genomic relationship matrix can improve genomic estimated breeding values in the training set. <i>Theoretical and Applied Genetics</i> , 2015, 128, 693-703.	3.6	16
60	Fine mapping of qhir8 affecting in vivo haploid induction in maize. <i>Theoretical and Applied Genetics</i> , 2015, 128, 2507-2515.	3.6	52
61	Cold Tolerance in Two Large Maize Inbred Panels Adapted to European Climates. <i>Crop Science</i> , 2014, 54, 1981-1991.	1.8	30
62	In Vivo Haploid Induction in Maize: Identification of Haploid Seeds by Their Oil Content. <i>Crop Science</i> , 2014, 54, 1497-1504.	1.8	59
63	Optimizing Resource Allocation for Multistage Selection in Plant Breeding with R Package <i>&lt;i&gt;Selectiongain&lt;/i&gt;</i> . <i>Crop Science</i> , 2014, 54, 1413-1418.	1.8	23
64	Breeding Potential of European Flint Maize Landraces Evaluated by their Testcross Performance. <i>Crop Science</i> , 2014, 54, 1665-1672.	1.8	25
65	Linkage Disequilibrium with Linkage Analysis of Multiline Crosses Reveals Different Multiallelic QTL for Hybrid Performance in the Flint and Dent Heterotic Groups of Maize. <i>Genetics</i> , 2014, 198, 1717-1734.	2.9	89
66	Characterization of Sudanese pearl millet germplasm for agro-morphological traits and grain nutritional values. <i>Plant Genetic Resources: Characterisation and Utilisation</i> , 2014, 12, 35-47.	0.8	38
67	Recovering Power in Association Mapping Panels with Variable Levels of Linkage Disequilibrium. <i>Genetics</i> , 2014, 197, 375-387.	2.9	89
68	Dent and Flint maize diversity panels reveal important genetic potential for increasing biomass production. <i>Theoretical and Applied Genetics</i> , 2014, 127, 2313-2331.	3.6	41
69	Optimum allocation of test resources and comparison of breeding strategies for hybrid wheat. <i>Theoretical and Applied Genetics</i> , 2014, 127, 2117-2126.	3.6	14
70	Genome Properties and Prospects of Genomic Prediction of Hybrid Performance in a Breeding Program of Maize. <i>Genetics</i> , 2014, 197, 1343-1355.	2.9	192
71	Identification of key ancestors of modern germplasm in a breeding program of maize. <i>Theoretical and Applied Genetics</i> , 2014, 127, 2545-2553.	3.6	22
72	Usefulness of Multiparental Populations of Maize ( <i>&lt;i&gt;Zea mays&lt;/i&gt;</i> L.) for Genome-Based Prediction. <i>Genetics</i> , 2014, 198, 3-16.	2.9	114

#	ARTICLE	IF	CITATIONS
73	Genome-wide meta-analysis of maize heterosis reveals the potential role of additive gene expression at pericentromeric loci. <i>BMC Plant Biology</i> , 2014, 14, 88.	3.6	54
74	Optimizing experimental procedures for quantitative evaluation of crop plant performance in high throughput phenotyping systems. <i>Frontiers in Plant Science</i> , 2014, 5, 770.	3.6	187
75	High-density genotyping: an overkill for QTL mapping? Lessons learned from a case study in maize and simulations. <i>Theoretical and Applied Genetics</i> , 2013, 126, 2563-2574.	3.6	63
76	Optimizing the allocation of resources for genomic selection in one breeding cycle. <i>Theoretical and Applied Genetics</i> , 2013, 126, 2835-2848.	3.6	74
77	Genomic Predictability of Interconnected Biparental Maize Populations. <i>Genetics</i> , 2013, 194, 493-503.	2.9	180
78	Intraspecific variation of recombination rate in maize. <i>Genome Biology</i> , 2013, 14, R103.	9.6	176
79	The maize leaf lipidome shows multilevel genetic control and high predictive value for agronomic traits. <i>Scientific Reports</i> , 2013, 3, 2479.	3.3	29
80	Genetic diversity analysis of elite European maize ( <i>Zea mays</i> L.) inbred lines using AFLP, SSR, and SNP markers reveals ascertainment bias for a subset of SNPs. <i>Theoretical and Applied Genetics</i> , 2013, 126, 133-141.	3.6	97
81	Association mapping for chilling tolerance in elite flint and dent maize inbred lines evaluated in growth chamber and field experiments. <i>Plant, Cell and Environment</i> , 2013, 36, 1871-1887.	5.7	73
82	Optimum design of family structure and allocation of resources in association mapping with lines from multiple crosses. <i>Heredity</i> , 2013, 110, 71-79.	2.6	34
83	High-density linkage mapping of yield components and epistatic interactions in maize with doubled haploid lines from four crosses. <i>Molecular Breeding</i> , 2013, 32, 533-546.	2.1	12
84	Effectiveness of selection at <scp>CIMMYT</scp>'s main maize breeding sites in Mexico for performance at sites in Africa and vice versa. <i>Plant Breeding</i> , 2013, 132, 299-304.	1.9	5
85	Genomic prediction of dichotomous traits with Bayesian logistic models. <i>Theoretical and Applied Genetics</i> , 2013, 126, 1133-1143.	3.6	12
86	QTL mapping of stalk bending strength in a recombinant inbred line maize population. <i>Theoretical and Applied Genetics</i> , 2013, 126, 2257-2266.	3.6	63
87	Rapid and accurate identification of in vivo-induced haploid seeds based on oil content in maize. <i>Scientific Reports</i> , 2013, 3, 2129.	3.3	95
88	Gametophytic and zygotic selection leads to segregation distortion through in vivo induction of a maternal haploid in maize. <i>Journal of Experimental Botany</i> , 2013, 64, 1083-1096.	4.8	107
89	Precision phenotyping of biomass accumulation in triticale reveals temporal genetic patterns of regulation. <i>Scientific Reports</i> , 2013, 3, 2442.	3.3	99
90	Mapping quantitative trait loci for freezing tolerance in a recombinant inbred line population of <i><scp>A</scp>rabidopsis thaliana</i> accessions Tenela and <scp>C24</scp> reveals <scp>REVEILLE1</scp> as negative regulator of cold acclimation. <i>Plant, Cell and Environment</i> , 2013, 36, 1256-1267.	5.7	48

#	ARTICLE	IF	CITATIONS
91	Genomic Prediction of Northern Corn Leaf Blight Resistance in Maize with Combined or Separated Training Sets for Heterotic Groups. G3: Genes, Genomes, Genetics, 2013, 3, 197-203.	1.8	112
92	Unlocking the Genetic Diversity of Maize Landraces with Doubled Haploids Opens New Avenues for Breeding. PLoS ONE, 2013, 8, e57234.	2.5	68
93	Relationship of Line per se and Testcross Performance for Grain Yield of Tropical Maize in Drought and Well-Watered Trials. Crop Science, 2013, 53, 1228-1236.	1.8	11
94	Comparative Quantitative Trait Loci Mapping for Gibberella Ear Rot Resistance and Reduced Deoxynivalenol Contamination across Connected Maize Populations. Crop Science, 2012, 52, 32-43.	1.8	27
95	Haploid Fertility in Temperate and Tropical Maize Germplasm. Crop Science, 2012, 52, 623-630.	1.8	61
96	Genetic Variation among Inbred Lines and Testcrosses of Maize for Early Growth Parameters and Their Relationship to Final Dry Matter Yield. Crop Science, 2012, 52, 1084-1092.	1.8	26
97	Strategies to Subdivide a Target Population of Environments: Results from the CIMMYT-Led Maize Hybrid Testing Programs in Africa. Crop Science, 2012, 52, 2143-2152.	1.8	49
98	Selection Strategy for Sorghum Targeting Phosphorus-Limited Environments in West Africa: Analysis of Multi-Environment Experiments. Crop Science, 2012, 52, 2517-2527.	1.8	41
99	Effectiveness of Genomic Prediction of Maize Hybrid Performance in Different Breeding Populations and Environments. G3: Genes, Genomes, Genetics, 2012, 2, 1427-1436.	1.8	242
100	Efficiency of Managed-Stress Screening of Elite Maize Hybrids under Drought and Low Nitrogen for Yield under Rainfed Conditions in Southern Africa. Crop Science, 2012, 52, 1011-1020.	1.8	84
101	Genome-wide association mapping of leaf metabolic profiles for dissecting complex traits in maize. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 8872-8877.	7.1	340
102	Genomic prediction of hybrid performance in maize with models incorporating dominance and population specific marker effects. Theoretical and Applied Genetics, 2012, 125, 1181-1194.	3.6	143
103	Genomic and metabolic prediction of complex heterotic traits in hybrid maize. Nature Genetics, 2012, 44, 217-220.	21.4	532
104	New Insights into the Genetics of <i>in Vivo</i> Induction of Maternal Haploids, the Backbone of Doubled Haploid Technology in Maize. Genetics, 2012, 190, 781-793.	2.9	143
105	Production of Haploids and Doubled Haploids in Maize. Methods in Molecular Biology, 2012, 877, 161-172.	0.9	69
106	Comparison of whole-genome prediction models for traits with contrasting genetic architecture in a diversity panel of maize inbred lines. BMC Genomics, 2012, 13, 452.	2.8	74
107	Association analysis of photoperiodic flowering time genes in west and central African sorghum [Sorghum bicolor (L.) Moench]. BMC Plant Biology, 2012, 12, 32.	3.6	30
108	Genome-wide association mapping of flowering time and northern corn leaf blight (Setosphaeria) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	3.6	70



#	ARTICLE	IF	CITATIONS
109	Maximizing the Reliability of Genomic Selection by Optimizing the Calibration Set of Reference Individuals: Comparison of Methods in Two Diverse Groups of Maize Inbreds ( <i>Zea mays</i> L.). Genetics, 2012, 192, 715-728.	2.9	258
110	Breeding maize as biogas substrate in Central Europe: I. Quantitative-genetic parameters for testcross performance. Theoretical and Applied Genetics, 2012, 124, 971-980.	3.6	34
111	Breeding maize as biogas substrate in Central Europe: II. Quantitative-genetic parameters for inbred lines and correlations with testcross performance. Theoretical and Applied Genetics, 2012, 124, 981-988.	3.6	24
112	Variation and covariation for Gibberella ear rot resistance and agronomic traits in testcrosses of doubled haploid maize lines. Euphytica, 2012, 185, 441-451.	1.2	7
113	Doubled haploids in tropical maize: II. Quantitative genetic parameters for testcross performance. Euphytica, 2012, 185, 453-463.	1.2	10
114	Development of in vivo haploid inducers for tropical maize breeding programs. Euphytica, 2012, 185, 481-490.	1.2	52
115	Inheritance of resistance to Gibberella ear rot and deoxynivalenol contamination in five flint maize crosses. Plant Breeding, 2012, 131, 28-32.	1.9	18
116	Prediction of grain yield using reflectance spectra of canopy and leaves in maize plants grown under different water regimes. Field Crops Research, 2012, 128, 82-90.	5.1	144
117	Kinetics of methane fermentation yield in biogas reactors: Genetic variation and association with chemical composition in maize. Biomass and Bioenergy, 2012, 37, 132-141.	5.7	27
118	QTL mapping under truncation selection in homozygous lines derived from biparental crosses. Theoretical and Applied Genetics, 2012, 124, 543-553.	3.6	5
119	Partial least squares regression, support vector machine regression, and transcriptome-based distances for prediction of maize hybrid performance with gene expression data. Theoretical and Applied Genetics, 2012, 124, 825-833.	3.6	33
120	Mapping of QTL for resistance to first and second generation of European corn borer using an integrated SNP and SSR linkage map. Euphytica, 2012, 183, 197-206.	1.2	16
121	Determination of Methane Fermentation Yield and its Kinetics by near Infrared Spectroscopy and Chemical Composition in Maize. Journal of Near Infrared Spectroscopy, 2011, 19, 463-477.	1.5	17
122	REML-Based Diallel Analysis. Crop Science, 2011, 51, 470-478.	1.8	47
123	High-throughput non-destructive biomass determination during early plant development in maize under field conditions. Field Crops Research, 2011, 121, 268-273.	5.1	139
124	Effect of source germplasm and season on the in vivo haploid induction rate in tropical maize. Euphytica, 2011, 180, 219-226.	1.2	59
125	Population structure in sorghum accessions from West Africa differing in race and maturity class. Genetica, 2011, 139, 453-463.	1.1	19
126	Best linear unbiased prediction and optimum allocation of test resources in maize breeding with doubled haploids. Theoretical and Applied Genetics, 2011, 123, 1-10.	3.6	42



#	ARTICLE	IF	CITATIONS
127	Extent and genome-wide distribution of linkage disequilibrium in commercial maize germplasm. Theoretical and Applied Genetics, 2011, 123, 11-20.	3.6	75
128	Gene stacking strategies with doubled haploids derived from biparental crosses: theory and simulations assuming a finite number of loci. Theoretical and Applied Genetics, 2011, 123, 1269-1279.	3.6	9
129	Doubled Haploids in Tropical Maize: I. Effects of Inducers and Source Germplasm on in vivo Haploid Induction Rates. Crop Science, 2011, 51, 1498-1506.	1.8	94
130	Colocalization of QTL for Gibberella Ear Rot Resistance and Low Mycotoxin Contamination in Early European Maize. Crop Science, 2011, 51, 1935-1945.	1.8	44
131	Broadening the genetic base of European maize heterotic pools with US Cornbelt germplasm using field and molecular marker data. Theoretical and Applied Genetics, 2010, 120, 301-310.	3.6	38
132	QTL analysis of early stage heterosis for biomass in Arabidopsis. Theoretical and Applied Genetics, 2010, 120, 227-237.	3.6	90
133	Molecular marker assisted broadening of the Central European heterotic groups in rye with Eastern European germplasm. Theoretical and Applied Genetics, 2010, 120, 291-299.	3.6	39
134	Optimum allocation of resources for QTL detection using a nested association mapping strategy in maize. Theoretical and Applied Genetics, 2010, 120, 553-561.	3.6	14
135	Hybrid maize breeding with doubled haploids: V. Selection strategies for testcross performance with variable sizes of crosses and S1 families. Theoretical and Applied Genetics, 2010, 120, 699-708.	3.6	6
136	Correlation between parental transcriptome and field data for the characterization of heterosis in Zea mays L.. Theoretical and Applied Genetics, 2010, 120, 401-413.	3.6	79
137	Variation of the parental genome contribution in segregating populations derived from biparental crosses and its relationship with heterosis of their Design III progenies. Theoretical and Applied Genetics, 2010, 120, 311-319.	3.6	5
138	Transcriptome-based distance measures for grouping of germplasm and prediction of hybrid performance in maize. Theoretical and Applied Genetics, 2010, 120, 441-450.	3.6	111
139	Prediction of hybrid performance in maize using molecular markers and joint analyses of hybrids and parental inbreds. Theoretical and Applied Genetics, 2010, 120, 451-461.	3.6	70
140	High congruency of QTL positions for heterosis of grain yield in three crosses of maize. Theoretical and Applied Genetics, 2010, 120, 321-332.	3.6	75
141	The International Conference on "Heterosis in Plants". Theoretical and Applied Genetics, 2010, 120, 201-203.	3.6	6
142	Dissection of the genetic basis of heterosis in an elite maize hybrid by QTL mapping in an immortalized F2 population. Theoretical and Applied Genetics, 2010, 120, 333-340.	3.6	132
143	Population structure and genetic diversity in a commercial maize breeding program assessed with SSR and SNP markers. Theoretical and Applied Genetics, 2010, 120, 1289-1299.	3.6	232
144	Patterns of molecular and phenotypic diversity in pearl millet [Pennisetum glaucum (L.) R. Br.] from West and Central Africa and their relation to geographical and environmental parameters. BMC Plant Biology, 2010, 10, 216.	3.6	55

#	ARTICLE	IF	CITATIONS
145	Dissecting grain yield pathways and their interactions with grain dry matter content by a two-step correlation approach with maize seedling transcriptome. BMC Plant Biology, 2010, 10, 63.	3.6	29
146	Genetic Variation in Testcrosses and Relationship between Line per se and Testcross Performance for Resistance to Gibberella Ear Rot in Maize. Crop Science, 2010, 50, 1691-1696.	1.8	12
147	Development of Heterotic Groups in Triticale. Crop Science, 2010, 50, 584-590.	1.8	76
148	Effect of N supply on stalk quality in maize hybrids. Field Crops Research, 2010, 118, 208-214.	5.1	25
149	Genetic Variation for Resistance to Ear Rots and Mycotoxins Contamination in Early European Maize Inbred Lines. Crop Science, 2009, 49, 2019-2028.	1.8	60
150	Impact of Genetic Divergence on the Ratio of Variance Due to Specific vs. General Combining Ability in Winter Triticale. Crop Science, 2009, 49, 2119-2122.	1.8	14
151	Unraveling Epistasis With Triple Testcross Progenies of Near-Isogenic Lines. Genetics, 2009, 181, 247-257.	2.9	28
152	Comparison of mixed-model approaches for association mapping in rapeseed, potato, sugar beet, maize, and Arabidopsis. BMC Genomics, 2009, 10, 94.	2.8	79
153	Molecular marker-based prediction of hybrid performance in maize using unbalanced data from multiple experiments with factorial crosses. Theoretical and Applied Genetics, 2009, 118, 741-751.	3.6	76
154	Efficiency gain of marker-assisted backcrossing by sequentially increasing marker densities over generations. Theoretical and Applied Genetics, 2009, 119, 23-32.	3.6	22
155	Identification of heterotic metabolite QTL in <i>Arabidopsis thaliana</i> RIL and IL populations. Plant Journal, 2009, 59, 777-788.	5.7	95
156	Optimum allocation of test resources and relative efficiency of alternative procedures of within-family selection in hybrid breeding. Plant Breeding, 2009, 128, 213-216.	1.9	3
157	A New near Infrared Spectroscopy Sample Presentation Unit for Measuring Feeding Quality of Maize Stover. Journal of Near Infrared Spectroscopy, 2009, 17, 195-201.	1.5	6
158	BLUP for phenotypic selection in plant breeding and variety testing. Euphytica, 2008, 161, 209-228.	1.2	569
159	The Plabsoft database: a comprehensive database management system for integrating phenotypic and genomic data in academic and commercial plant breeding programs. Euphytica, 2008, 161, 173-179.	1.2	11
160	Population genetic simulation and data analysis with Plabsoft. Euphytica, 2008, 161, 133-139.	1.2	62
161	Comparison of the observed with the simulated distributions of the parental genome contribution in two marker-assisted backcross programs in rice. Theoretical and Applied Genetics, 2008, 116, 739-744.	3.6	18
162	Hybrid maize breeding with doubled haploids. IV. Number versus size of crosses and importance of parental selection in two-stage selection for testcross performance. Theoretical and Applied Genetics, 2008, 117, 251-260.	3.6	20

#	ARTICLE	IF	CITATIONS
163	Multi-trait association mapping in sugar beet ( <i>Beta vulgaris</i> L.). Theoretical and Applied Genetics, 2008, 117, 947-954.	3.6	57
164	Association mapping in multiple segregating populations of sugar beet ( <i>Beta vulgaris</i> L.). Theoretical and Applied Genetics, 2008, 117, 1167-1179.	3.6	28
165	Trends in genetic variance components during 30 years of hybrid maize breeding at the University of Hohenheim. Plant Breeding, 2008, 127, 446-451.	1.9	44
166	Feeding quality assessment of fresh maize stover by means of near-infrared spectroscopy with a new sample presentation unit. Plant Breeding, 2008, 127, 214-216.	1.9	2
167	Comparison of Mixed-Model Approaches for Association Mapping. Genetics, 2008, 178, 1745-1754.	2.9	273
168	Genetic Expectations of Quantitative Trait Loci Main and Interaction Effects Obtained With the Triple Testcross Design and Their Relevance for the Analysis of Heterosis. Genetics, 2008, 178, 2265-2274.	2.9	30
169	Precision of Recombination Frequency Estimates After Random Intermating With Finite Population Sizes. Genetics, 2008, 178, 597-600.	2.9	5
170	Quantitative Trait Loci Mapping and The Genetic Basis of Heterosis in Maize and Rice. Genetics, 2008, 180, 1707-1724.	2.9	141
171	Genetic variation for resistance and mycotoxin content of European maize inoculated with <i>Fusarium graminearum</i> and <i>F. verticillioides</i> . Cereal Research Communications, 2008, 36, 45-48.	1.6	6
172	Genetic Diversity in CIMMYT Nontemperate Maize Germplasm: Landraces, Open Pollinated Varieties, and Inbred Lines. Crop Science, 2008, 48, 617-624.	1.8	93
173	The Role of Epistasis in the Manifestation of Heterosis: A Systems-Oriented Approach. Genetics, 2007, 177, 1815-1825.	2.9	125
174	Variance of the Parental Genome Contribution to Inbred Lines Derived From Biparental Crosses. Genetics, 2007, 176, 477-488.	2.9	15
175	Chilling Tolerance of Central European Maize Lines and their Factorial Crosses. Annals of Botany, 2007, 100, 1315-1321.	2.9	37
176	Analysis of a Triple Testcross Design With Recombinant Inbred Lines Reveals a Significant Role of Epistasis in Heterosis for Biomass-Related Traits in Arabidopsis. Genetics, 2007, 175, 2009-2017.	2.9	65
177	Genetic Basis of Heterosis for Growth-Related Traits in Arabidopsis Investigated by Testcross Progenies of Near-Isogenic Lines Reveals a Significant Role of Epistasis. Genetics, 2007, 177, 1827-1837.	2.9	95
178	Power to Detect Higher-Order Epistatic Interactions in a Metabolic Pathway Using a New Mapping Strategy. Genetics, 2007, 176, 563-570.	2.9	43
179	Heterosis for Biomass-Related Traits in Arabidopsis Investigated by Quantitative Trait Loci Analysis of the Triple Testcross Design With Recombinant Inbred Lines. Genetics, 2007, 177, 1839-1850.	2.9	55
180	Novel throughput phenotyping platforms in plant genetic studies. Trends in Plant Science, 2007, 12, 433-436.	8.8	220

#	ARTICLE	IF	CITATIONS
181	Quality assessment of rapeseed accessions by means of near-infrared spectroscopy on combine harvesters. <i>Plant Breeding</i> , 2007, 126, 329-330.	1.9	16
182	Determination of chemical composition and nutritional attributes of silage corn hybrids by near-infrared spectroscopy on chopper: evaluation of traits, sample presentation systems and calibration transferability. <i>Plant Breeding</i> , 2007, 126, 521-526.	1.9	10
183	Hybrid maize breeding with doubled haploids: II. Optimum type and number of testers in two-stage selection for general combining ability. <i>Theoretical and Applied Genetics</i> , 2007, 114, 393-402.	3.6	27
184	Temporal changes in allele frequencies in two European F2 flint maize populations under modified recurrent full-sib selection. <i>Theoretical and Applied Genetics</i> , 2007, 114, 765-776.	3.6	20
185	Prediction of single-cross hybrid performance in maize using haplotype blocks associated with QTL for grain yield. <i>Theoretical and Applied Genetics</i> , 2007, 114, 1345-1355.	3.6	33
186	Linkage disequilibrium in two European F2 flint maize populations under modified recurrent full-sib selection. <i>Theoretical and Applied Genetics</i> , 2007, 115, 289-297.	3.6	6
187	An incomplete enumeration algorithm for an exact test of Hardy-Weinberg proportions with multiple alleles. <i>Theoretical and Applied Genetics</i> , 2007, 115, 393-398.	3.6	7
188	Hybrid maize breeding with doubled haploids: III. Efficiency of early testing prior to doubled haploid production in two-stage selection for testcross performance. <i>Theoretical and Applied Genetics</i> , 2007, 115, 519-527.	3.6	27
189	Potential causes of linkage disequilibrium in a European maize breeding program investigated with computer simulations. <i>Theoretical and Applied Genetics</i> , 2007, 115, 529-536.	3.6	20
190	Comparison of transcript profiles between near-isogenic maize lines in association with SCMV resistance based on unigene-microarrays. <i>Plant Science</i> , 2006, 170, 159-169.	3.6	22
191	Modified full-sib selection and best linear unbiased prediction of progeny performance in a European F2 maize population. <i>Plant Breeding</i> , 2006, 125, 248-253.	1.9	7
192	Prospects for hybrid breeding in winter triticale: II. Relationship between parental genetic distance and specific combining ability. <i>Plant Breeding</i> , 2006, 125, 331-336.	1.9	24
193	Two chromosome segments confer multiple potyvirus resistance in maize. <i>Plant Breeding</i> , 2006, 125, 352-356.	1.9	25
194	Near-infrared spectroscopy on combine harvesters to measure maize grain dry matter content and quality parameters. <i>Plant Breeding</i> , 2006, 125, 591-595.	1.9	36
195	An extension of the Smith model for quantitative genetic analysis of selection response under recurrent selection. <i>Plant Breeding</i> , 2006, 125, 644-646.	1.9	1
196	Genetic Diversity and Relationships of Wheat Landraces from Oman Investigated with SSR Markers. <i>Genetic Resources and Crop Evolution</i> , 2006, 53, 1351-1360.	1.6	43
197	Identification of Essentially Derived Varieties Obtained from Biparental Crosses of Homozygous Lines. III. AFLP Data from Maize Inbreds and Comparison with SSR Data. <i>Molecular Breeding</i> , 2006, 17, 111-125.	2.1	26
198	Comparison of Linkage Disequilibrium in Elite European Maize Inbred Lines using AFLP and SSR Markers. <i>Molecular Breeding</i> , 2006, 17, 217-226.	2.1	52

#	ARTICLE	IF	CITATIONS
199	Trends in population parameters and best linear unbiased prediction of progeny performance in a European F2 maize population under modified recurrent full-sib selection. Theoretical and Applied Genetics, 2006, 112, 483-491.	3.6	11
200	Hybrid maize breeding with doubled haploids: I. One-stage versus two-stage selection for testcross performance. Theoretical and Applied Genetics, 2006, 112, 903-912.	3.6	35
201	Grouping of accessions of Mexican races of maize revisited with SSR markers. Theoretical and Applied Genetics, 2006, 113, 177-185.	3.6	81
202	Comparison of linkage maps from F2 and three times intermated generations in two populations of European flint maize ( <i>Zea mays</i> L.). Theoretical and Applied Genetics, 2006, 113, 857-866.	3.6	11
203	Prediction of single-cross hybrid performance for grain yield and grain dry matter content in maize using AFLP markers associated with QTL. Theoretical and Applied Genetics, 2006, 113, 1037-1047.	3.6	92
204	Genetic drift and selection effects of modified recurrent full-sib selection programs in two F2 populations of European flint maize. Theoretical and Applied Genetics, 2006, 113, 1113-1120.	3.6	7
205	A new test for family-based association mapping with inbred lines from plant breeding programs. Theoretical and Applied Genetics, 2006, 113, 1121-1130.	3.6	33
206	Marker-Based Prediction of the Parental Genome Contribution to Inbred Lines Derived From Biparental Crosses. Genetics, 2006, 174, 795-803.	2.9	8
207	Genetic diversity trends in Central European heterotic groups. Acta Agronomica Hungarica: an International Multidisciplinary Journal in Agricultural Science, 2006, 54, 315-320.	0.2	1
208	Hybrid maize breeding with doubled haploids: Comparison between selection criteria. Acta Agronomica Hungarica: an International Multidisciplinary Journal in Agricultural Science, 2006, 54, 343-350.	0.2	4
209	Prospects for Hybrid Breeding in Winter Triticale: I. Heterosis and Combining Ability for Agronomic Traits in European Elite Germplasm. Crop Science, 2005, 45, 1476-1482.	1.8	66
210	Genetic diversity in European perennial ryegrass cultivars investigated with RAPD markers. Plant Breeding, 2005, 124, 161-166.	1.9	56
211	Genetic similarity among European winter triticale elite germplasms assessed with AFLP and comparisons with SSR and pedigree data. Plant Breeding, 2005, 124, 154-160.	1.9	41
212	Molecular genetic diversity within and among German ecotypes in comparison to European perennial ryegrass cultivars. Plant Breeding, 2005, 124, 257-262.	1.9	40
213	Wheat genetic diversity trends during domestication and breeding. Theoretical and Applied Genetics, 2005, 110, 859-864.	3.6	365
214	Trends in genetic diversity among European maize cultivars and their parental components during the past 50 years. Theoretical and Applied Genetics, 2005, 111, 838-845.	3.6	93
215	Genetic structure and diversity of European flint maize populations determined with SSR analyses of individuals and bulks. Theoretical and Applied Genetics, 2005, 111, 906-913.	3.6	96
216	New editorial office. Theoretical and Applied Genetics, 2005, 111, 993-993.	3.6	0

#	ARTICLE	IF	CITATIONS
217	QTL mapping of resistance to Sclerotinia midstalk rot in RIL of sunflower population NDBLOSselAA—ACM625. Theoretical and Applied Genetics, 2005, 110, 1490-1498.	3.6	39
218	Validation of Dwarf8 polymorphisms associated with flowering time in elite European inbred lines of maize ( <i>Zea mays</i> L.). Theoretical and Applied Genetics, 2005, 111, 206-217.	3.6	115
219	Identification and validation of QTL for Sclerotinia midstalk rot resistance in sunflower by selective genotyping. Theoretical and Applied Genetics, 2005, 111, 233-242.	3.6	36
220	Identification of essentially derived varieties with molecular markers: an approach based on statistical test theory and computer simulations. Theoretical and Applied Genetics, 2005, 111, 598-608.	3.6	19
221	Linkage disequilibrium in European elite maize germplasm investigated with SSRs. Theoretical and Applied Genetics, 2005, 111, 723-730.	3.6	167
222	Identification by suppression subtractive hybridization of genes that are differentially expressed between near-isogenic maize lines in association with sugarcane mosaic virus resistance. Molecular Genetics and Genomics, 2005, 273, 450-461.	2.1	36
223	Hybrid performance and heterosis in spring bread wheat, and their relations to SSR-based genetic distances and coefficients of parentage. Euphytica, 2005, 144, 51-59.	1.2	52
224	Molecular characterization of genetic diversity in European germplasm of perennial ryegrass. Euphytica, 2005, 146, 39-44.	1.2	21
225	Quantifying novel sequence variation and selective advantage in synthetic hexaploid wheats and their backcross-derived lines using SSR markers. Molecular Breeding, 2005, 15, 1-10.	2.1	55
226	Association between SCMV Resistance and Macroarray-based Expression Patterns in Maize Inbreds. Molecular Breeding, 2005, 16, 173-184.	2.1	5
227	Identification of Essentially Derived Varieties Obtained from Biparental Crosses of Homozygous Lines: II. Morphological Distances and Heterosis in Comparison with Simple Sequence Repeat and Amplified Fragment Length Polymorphism Data in Maize. Crop Science, 2005, 45, 1132-1140.	1.8	26
228	Identification of Essentially Derived Varieties Obtained from Biparental Crosses of Homozygous Lines: I. Simple Sequence Repeat Data from Maize Inbreds. Crop Science, 2005, 45, 1120-1131.	1.8	21
229	Genetic Diversity among CIMMYT Maize Inbred Lines Investigated with SSR Markers: II. Subtropical, Tropical Midaltitude, and Highland Maize Inbred Lines and their Relationships with Elite U.S. and European Maize. Crop Science, 2005, 45, 2573-2582.	1.8	86
230	No Evidence for Epistasis in Hybrid and Per Se Performance of Elite European Flint Maize Inbreds from Generation Means and QTL Analyses. Crop Science, 2005, 45, 2605-2613.	1.8	69
231	Selection Theory for Marker-Assisted Backcrossing. Genetics, 2005, 170, 909-917.	2.9	99
232	Genetic Structure and Diversity among Radish Varieties as Inferred from AFLP and ISSR Analyses. Journal of the American Society for Horticultural Science, 2005, 130, 79-87.	1.0	27
233	Congruency of Quantitative Trait Loci Detected for Agronomic Traits in Testcrosses of Five Populations of European Maize. Crop Science, 2004, 44, 114-124.	1.8	41
234	Prospects for celeriac ( <i>Apium graveolens</i> var. <i>rapaceum</i> ) improvement by using genetic resources of <i>Apium</i> , as determined by AFLP markers and morphological characterization. Plant Genetic Resources: Characterisation and Utilisation, 2004, 2, 189-198.	0.8	9



#	ARTICLE	IF	CITATIONS
235	Quantitative Trait Locus Mapping Based on Resampling in a Vast Maize Testcross Experiment and Its Relevance to Quantitative Genetics for Complex Traits. <i>Genetics</i> , 2004, 167, 485-498.	2.9	231
236	Targeted BSA mapping of Scmv1 and Scmv2 conferring resistance to SCMV using PstI/MseI compared with EcoRI/MseI AFLP markers. <i>Plant Breeding</i> , 2004, 123, 434-437.	1.9	10
237	Genetic diversity in cornsalad ( <i>Valerianella locusta</i> ) and related species as determined by AFLP markers. <i>Plant Breeding</i> , 2004, 123, 460-466.	1.9	26
238	QTL analyses of complex traits with cross validation, bootstrapping and other biometric methods. <i>Euphytica</i> , 2004, 137, 1-11.	1.2	48
239	Genetic diversity in European winter triticale determined with SSR markers and coancestry coefficient. <i>Theoretical and Applied Genetics</i> , 2004, 108, 1385-1391.	3.6	58
240	QTL mapping for European corn borer resistance ( <i>Ostrinia nubilalis</i> Hb.), agronomic and forage quality traits of testcross progenies in early-maturing European maize ( <i>Zea mays</i> L.) germplasm. <i>Theoretical and Applied Genetics</i> , 2004, 108, 1545-1554.	3.6	54
241	QTL mapping of Sclerotinia midstalk-rot resistance in sunflower. <i>Theoretical and Applied Genetics</i> , 2004, 109, 1474-1484.	3.6	44
242	Congruency of Quantitative Trait Loci Detected for Agronomic Traits in Testcrosses of Five Populations of European Maize. <i>Crop Science</i> , 2004, 44, 114.	1.8	29
243	Title is missing!. <i>Molecular Breeding</i> , 2003, 12, 97-106.	2.1	28
244	Use of SSRs for establishing heterotic groups in subtropical maize. <i>Theoretical and Applied Genetics</i> , 2003, 107, 947-957.	3.6	110
245	Heterosis and combining ability for grain yield and other agronomic traits in winter triticale. <i>Plant Breeding</i> , 2003, 122, 318-321.	1.9	24
246	Concentration of moniliformin produced by <i>Fusarium</i> species in grains of transgenic Bt maize hybrids compared to their isogenic counterparts and commercial varieties under European corn borer pressure. <i>Plant Breeding</i> , 2003, 122, 322-327.	1.9	12
247	Heterosis for biomass yield and related traits in five hybrids of <i>Arabidopsis thaliana</i> L. Heynh. <i>Heredity</i> , 2003, 91, 36-42.	2.6	86
248	Molecular Genetic Diversity among Progenitors and Derived Elite Lines of BSSS and BSCB1 Maize Populations. <i>Crop Science</i> , 2003, 43, 474.	1.8	26
249	Genetic Characterization of CIMMYT Inbred Maize Lines and Open Pollinated Populations Using Large Scale Fingerprinting Methods. <i>Crop Science</i> , 2002, 42, 1832-1840.	1.8	141
250	Development of RGA-CAPS markers and genetic mapping of candidate genes for sugarcane mosaic virus resistance in maize. <i>Theoretical and Applied Genetics</i> , 2002, 105, 355-363.	3.6	41
251	Conversion of AFLP fragments tightly linked to SCMV resistance genes Scmv1 and Scmv2 into simple PCR-based markers. <i>Theoretical and Applied Genetics</i> , 2002, 105, 1190-1195.	3.6	28
252	Genetic diversity in <i>Arabidopsis thaliana</i> L. Heynh. investigated by cleaved amplified polymorphic sequence (CAPS) and inter-simple sequence repeat (ISSR) markers. <i>Molecular Ecology</i> , 2002, 11, 495-505.	3.9	80



#	ARTICLE	IF	CITATIONS
253	Relationship between European corn borer resistance and concentration of mycotoxins produced by <i>Fusarium</i> spp. in grains of transgenic Bt maize hybrids, their isogenic counterparts, and commercial varieties. <i>Plant Breeding</i> , 2002, 121, 146-154.	1.9	62
254	Title is missing!. <i>Molecular Breeding</i> , 2002, 10, 181-191.	2.1	74
255	Marker-Assisted Backcrossing for Simultaneous Introgression of Two Genes. <i>Crop Science</i> , 2001, 41, 1716-1725.	1.8	61
256	Marker-Assisted Backcrossing for Introgression of a Recessive Gene. <i>Crop Science</i> , 2001, 41, 1485-1494.	1.8	28
257	Re-evaluation of the prospects of marker-assisted selection for improving insect resistance against <i>Diatraea</i> spp. in tropical maize by cross validation and independent validation. <i>Theoretical and Applied Genetics</i> , 2001, 103, 1059-1067.	3.6	58
258	Comparison of Bt maize hybrids with their non-transgenic counterparts and commercial varieties for resistance to European corn borer and for agronomic traits. <i>Plant Breeding</i> , 2001, 120, 397-403.	1.9	28
259	Influence of genetic background and heterozygosity on meiotic recombination in <i>Arabidopsis thaliana</i> . <i>Genome</i> , 2001, 44, 971-978.	2.0	28
260	The Length of the Intact Donor Chromosome Segment Around a Target Gene in Marker-Assisted Backcrossing. <i>Genetics</i> , 2001, 157, 1343-1356.	2.9	37
261	Influence of genetic background and heterozygosity on meiotic recombination in <i>Arabidopsis thaliana</i> . <i>Genome</i> , 2001, 44, 971-978.	2.0	5
262	Molecular mapping and gene action of Scm1 and Scm2, two major QTL contributing to SCMV resistance in maize. <i>Plant Breeding</i> , 2000, 119, 299-303.	1.9	51
263	Origin of Scm1 and Scm2—two loci conferring resistance to sugarcane mosaic virus (SCMV) in maize. <i>Theoretical and Applied Genetics</i> , 2000, 100, 934-941.	3.6	19
264	A high-throughput system for genome-wide measurement of genetic recombination in <i>Arabidopsis thaliana</i> based on transgenic markers. <i>Functional and Integrative Genomics</i> , 2000, 1, 200-206.	3.5	19
265	Relationships among Early European Maize Inbreds: IV. Genetic Diversity Revealed with AFLP Markers and Comparison with RFLP, RAPD, and Pedigree Data. <i>Crop Science</i> , 2000, 40, 783-791.	1.8	79
266	Bias and Sampling Error of the Estimated Proportion of Genotypic Variance Explained by Quantitative Trait Loci Determined From Experimental Data in Maize Using Cross Validation and Validation With Independent Samples. <i>Genetics</i> , 2000, 154, 1839-1849.	2.9	305
267	Resistance in the leaf and stem of sunflower after infection with two isolates of <i>Phomopsis</i> . <i>Plant Breeding</i> , 1999, 118, 405-410.	1.9	6
268	Optimal allocation of resources in evaluating current sunflower inbred lines for resistance to <i>Sclerotinia</i> . <i>Plant Breeding</i> , 1999, 118, 157-160.	1.9	14
269	Mean, genetic variance, and usefulness of selfing progenies from intra- and inter-pool crosses in faba beans ( <i>Vicia faba</i> L.) and their prediction from parental parameters. <i>Theoretical and Applied Genetics</i> , 1999, 98, 569-580.	3.6	13
270	QTL mapping of resistance to <i>Sporisorium reilianii</i> in maize. <i>Theoretical and Applied Genetics</i> , 1999, 99, 593-598.	3.6	40

#	ARTICLE	IF	CITATIONS
271	QTLs for resistance to <i>Setosphaeria turcica</i> in an early maturing Dent—Flint maize population. <i>Theoretical and Applied Genetics</i> , 1999, 99, 649-655.	3.6	39
272	Two high-density AFLP® linkage maps of <i>Zea mays</i> L.: analysis of distribution of AFLP markers. <i>Theoretical and Applied Genetics</i> , 1999, 99, 921-935.	3.6	217
273	High-resolution mapping of loci conferring resistance to sugarcane mosaic virus in maize using RFLP, SSR, and AFLP markers. <i>Molecular Genetics and Genomics</i> , 1999, 261, 574-581.	2.4	88
274	Genetic Similarities among Winter Wheat Cultivars Determined on the Basis of RFLPs, AFLPs, and SSRs and Their Use for Predicting Progeny Variance. <i>Crop Science</i> , 1999, 39, 228-237.	1.8	232
275	Quantitative Trait Loci Mapping of Resistance to Sugarcane Mosaic Virus in Maize. <i>Phytopathology</i> , 1999, 89, 660-667.	2.2	84
276	Comparison of Selection Strategies for Marker-Assisted Backcrossing of a Gene. <i>Crop Science</i> , 1999, 39, 1295-1301.	1.8	206
277	Prediction of testcross means and variances among F3 progenies of F1 crosses from testcross means and genetic distances of their parents in maize. <i>Theoretical and Applied Genetics</i> , 1998, 96, 503-512.	3.6	25
278	Genetic basis of resistance to sugarcane mosaic virus in European maize germplasm. <i>Theoretical and Applied Genetics</i> , 1998, 96, 1151-1161.	3.6	63
279	Comparative QTL mapping of resistance to <i>Ustilago maydis</i> across four populations of European flint-maize. <i>Theoretical and Applied Genetics</i> , 1998, 97, 1321-1330.	3.6	43
280	Comparative Quantitative Trait Loci Mapping of Partial Resistance to <i>Puccinia sorghi</i> Across Four Populations of European Flint Maize. <i>Phytopathology</i> , 1998, 88, 1324-1329.	2.2	28
281	Apparatus for Dissecting Stalks to Evaluate Stem Borer Insect Resistance in Maize. <i>Agronomy Journal</i> , 1998, 90, 233-234.	1.8	2
282	QTL Mapping in Testcrosses of Flint Lines of Maize: III. Comparison across Populations for Forage Traits. <i>Crop Science</i> , 1998, 38, 1278-1289.	1.8	76
283	Quantitative Trait Locus (QTL) Mapping Using Different Testers and Independent Population Samples in Maize Reveals Low Power of QTL Detection and Large Bias in Estimates of QTL Effects. <i>Genetics</i> , 1998, 149, 383-403.	2.9	462
284	QTL Mapping in Testcrosses of European Flint Lines of Maize: II. Comparison of Different Testers for Forage Quality Traits. <i>Crop Science</i> , 1997, 37, 1913-1922.	1.8	66
285	QTL Mapping in Testcrosses of European Flint Lines of Maize: I. Comparison of Different Testers for Forage Yield Traits. <i>Crop Science</i> , 1997, 37, 921-931.	1.8	113
286	Genetic diversity in European and Mediterranean faba bean germ plasm revealed by RAPD markers. <i>Theoretical and Applied Genetics</i> , 1995, 90, 27-32.	3.6	217
287	In an elite cross of maize a major quantitative trait locus controls one-fourth of the genetic variation for grain yield. <i>Theoretical and Applied Genetics</i> , 1995, 90, 415-424.	3.6	67
288	Reciprocal differences for forage traits in single and three-way crosses of maize. <i>Plant Breeding</i> , 1995, 114, 231-234.	1.9	2

#	ARTICLE	IF	CITATIONS
289	Epistasis in an Elite Maize Hybrid and Choice of Generation for Inbred Line Development. <i>Crop Science</i> , 1995, 35, 1272-1281.	1.8	73
290	Relationships among European Barley Germplasm: II. Comparison of RFLP and Pedigree Data. <i>Crop Science</i> , 1994, 34, 1199-1205.	1.8	111
291	RFLP Mapping in Maize: Quantitative Trait Loci Affecting Testcross Performance of Elite European Flint Lines. <i>Crop Science</i> , 1994, 34, 378-389.	1.8	147
292	Heterosis and gene effects of multiplicative characters: theoretical relationships and experimental results from <i>Vicia faba</i> L.. <i>Theoretical and Applied Genetics</i> , 1994, 88-88, 343-348.	3.6	21
293	Economic Aspects of Breeding for Yield and Quality Traits in Forage Maize. I. Determination of Economic Weights. <i>Plant Breeding</i> , 1994, 112, 102-109.	1.9	10
294	Economic Aspects of Breeding for Yield and Quality Traits in Forage Maize. II. Derivation and Evaluation of Selection Indices. <i>Plant Breeding</i> , 1994, 112, 110-119.	1.9	12
295	The Use of DNA Fingerprinting in Ecological Studies of <i>Phragmites australis</i> (Cav.) Trin. ex Steudel. <i>Botanica Acta</i> , 1994, 107, 237-242.	1.6	31
296	Phylogenetic relationships between cultivated and wild species of the genus <i>Beta</i> revealed by DNA fingerprinting?. <i>Theoretical and Applied Genetics</i> , 1993, 86, 449-457.	3.6	93
297	Mapping and characterization of quantitative trait loci affecting resistance against second-generation European corn borer in maize with the aid of RFLPs. <i>Heredity</i> , 1993, 70, 648-659.	2.6	135
298	Graphical Genotype of Maize Inbred B86 Revealed by RFLPs. <i>Plant Breeding</i> , 1993, 110, 29-34.	1.9	2
299	Genetic Diversity for RFLPs in European Maize Inbreds. III. Performance of Crosses Within versus Between Heterotic Groups for Grain Traits. <i>Plant Breeding</i> , 1993, 111, 217-226.	1.9	41
300	Relationships among Early European Maize Inbreds: II. Comparison of Pedigree and RFLP Data. <i>Crop Science</i> , 1993, 33, 944-950.	1.8	127
301	Relationships among Early European Maize Inbreds: I. Genetic Diversity among Flint and Dent Lines Revealed by RFLPs. <i>Crop Science</i> , 1992, 32, 1301-1309.	1.8	73
302	Genetic diversity of maize inbred lines within and among heterotic groups revealed by RFLPs. <i>Theoretical and Applied Genetics</i> , 1992, 84, 17-25.	3.6	59
303	Genetic diversity for RFLPs in European maize inbreds. <i>Theoretical and Applied Genetics</i> , 1992, 84-84, 672-681.	3.6	86
304	RFLP analyses of early-maturing European maize germ plasm. <i>Theoretical and Applied Genetics</i> , 1992, 83, 1003-1012.	3.6	62
305	Cluster analysis of RFLP data from related maize inbred lines of the BSSS and LSC heterotic groups and comparison with pedigree data. <i>Euphytica</i> , 1992, 60, 139-148.	1.2	20
306	Genetic Diversity for RFLPs in European Maize Inbreds: I. Relation to Performance of Flint & Dent Crosses for Forage Traits. <i>Crop Science</i> , 1992, 32, 895-902.	1.8	56

#	ARTICLE	IF	CITATIONS
307	Genetic diversity among progenitors and elite lines from the Iowa Stiff Stalk Synthetic (BSSS) maize population: comparison of allozyme and RFLP data. Theoretical and Applied Genetics, 1991, 83, 97-107.	3.6	74
308	Use of Molecular Markers in Breeding for Oligogenic Disease Resistance. Plant Breeding, 1990, 104, 1-19.	1.9	191
309	Genetic diversity for restriction fragment length polymorphisms and heterosis for two diallel sets of maize inbreds. Theoretical and Applied Genetics, 1990, 80, 488-496.	3.6	124
310	Optimum prediction of three-way crosses from single crosses in forage maize (Zea mays L.). Theoretical and Applied Genetics, 1987, 74, 339-345.	3.6	40
311	Expectation of means and variances of testcrosses produced from F2 and backcross individuals and their selfed progenies. Heredity, 1987, 59, 105-115.	2.6	36
312	Epistasis in Maize (Zea mays L.) III. Comparison of Single and Three-Way Crosses for Forage Traits. Plant Breeding, 1987, 98, 185-193.	1.9	3
313	Epistasis in maize (Zea mays L.). Theoretical and Applied Genetics, 1986, 72, 231-239.	3.6	32
314	Evaluation of Near Infra-red Reflectance Spectroscopy for Predicting Grain and Stover Quality Traits in Maize. Plant Breeding, 1986, 97, 20-29.	1.9	33
315	Overview of Heterosis and Heterotic Groups in Agronomic Crops. CSSA Special Publication - Crop Science Society of America, 0, , 29-44.	0.1	109