

Gabriel Frampton

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6447430/publications.pdf>

Version: 2024-02-01

19
papers

654
citations

759233

12
h-index

888059

17
g-index

19
all docs

19
docs citations

19
times ranked

886
citing authors

#	ARTICLE	IF	CITATIONS
1	Bile acids permeabilize the blood brain barrier after bile duct ligation in rats via Rac1-dependent mechanisms. <i>Digestive and Liver Disease</i> , 2014, 46, 527-534.	0.9	167
2	Interleukin-6-driven progranulin expression increases cholangiocarcinoma growth by an Akt-dependent mechanism. <i>Gut</i> , 2012, 61, 268-277.	12.1	101
3	Bile Acid Signaling Is Involved in the Neurological Decline in a Murine Model of Acute Liver Failure. <i>American Journal of Pathology</i> , 2016, 186, 312-323.	3.8	76
4	Suppression of the HPA Axis During Cholestasis Can Be Attributed to Hypothalamic Bile Acid Signaling. <i>Molecular Endocrinology</i> , 2015, 29, 1720-1730.	3.7	65
5	Gli1 activation and protection against hepatic encephalopathy is suppressed by circulating transforming growth factor β 1 in mice. <i>Journal of Hepatology</i> , 2014, 61, 1260-1266.	3.7	47
6	Opposing actions of endocannabinoids on cholangiocarcinoma growth is via the differential activation of Notch signaling. <i>Experimental Cell Research</i> , 2010, 316, 1465-1478.	2.6	36
7	Direct Comparison of the Thioacetamide and Azoxymethane Models of Type A Hepatic Encephalopathy in Mice. <i>Gene Expression</i> , 2018, 18, 171-185.	1.2	29
8	Fractalkine suppression during hepatic encephalopathy promotes neuroinflammation in mice. <i>Journal of Neuroinflammation</i> , 2016, 13, 198.	7.2	25
9	Elevated circulating TGF β 1 during acute liver failure activates TGF β 2R2 on cortical neurons and exacerbates neuroinflammation and hepatic encephalopathy in mice. <i>Journal of Neuroinflammation</i> , 2019, 16, 69.	7.2	19
10	The novel growth factor, progranulin, stimulates mouse cholangiocyte proliferation via sirtuin-1-mediated inactivation of FOXO1. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 303, G1202-G1211.	3.4	17
11	The Neuropeptide Galanin Is Up-Regulated during Cholestasis and Contributes to Cholangiocyte Proliferation. <i>American Journal of Pathology</i> , 2017, 187, 819-830.	3.8	16
12	Glucocorticoids Cause Gender-Dependent Reversal of Hepatic Fibrosis in the MDR2-Knockout Mouse Model. <i>International Journal of Molecular Sciences</i> , 2017, 18, 2389.	4.1	15
13	The TGF β 1 Receptor Antagonist GW788388 Reduces JNK Activation and Protects Against Acetaminophen Hepatotoxicity in Mice. <i>Toxicological Sciences</i> , 2019, 170, 549-561.	3.1	12
14	Leptin Enhances Hepatic Fibrosis and Inflammation in a Mouse Model of Cholestasis. <i>American Journal of Pathology</i> , 2022, 192, 484-502.	3.8	12
15	Coordinated Targeting of Galanin Receptors on Cholangiocytes and Hepatic Stellate Cells Ameliorates Liver Fibrosis in Multidrug Resistance Protein 2 Knockout Mice. <i>American Journal of Pathology</i> , 2020, 190, 586-601.	3.8	8
16	Ghrelin reverses ductular reaction and hepatic fibrosis in a rodent model of cholestasis. <i>Scientific Reports</i> , 2020, 10, 16024.	3.3	7
17	Increased serum bile acids after extrahepatic biliary obstruction causes leakiness to the blood brain barrier via the disruption of tight junctions. <i>FASEB Journal</i> , 2012, 26, 1110.7.	0.5	2
18	Characterization of Hepatic Injury During the Azoxymethane Model of Acute Liver Failure. <i>FASEB Journal</i> , 2022, 36, .	0.5	0

#	ARTICLE	IF	CITATIONS
19	LPS-induced endotoxemia promotes blood-brain barrier permeability via TSP1-dependent TGF β 1 activation. FASEB Journal, 2022, 36, .	0.5	0