## Sugihiko Hoshizaki

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6497809/publications.pdf

Version: 2024-02-01

933447 839539 19 432 10 18 citations g-index h-index papers 19 19 19 367 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Perâ€stage growth rates of head and body sizes in scarab larvae (Coleoptera: Scarabaeidae) decrease across ontogeny following a speciesâ€specific pattern. Entomological Science, 2020, 23, 374-384.	0.6	1
2	Oviposition in the onion fly <i>Delia antiqua</i> (Diptera: Anthomyiidae) is socially facilitated by visual cues. Bulletin of Entomological Research, 2020, 110, 677-683.	1.0	2
3	Catchâ€up growth in the rhinoceros beetle <i>Trypoxylus dichotomus</i> (Coleoptera: Scarabaeidae): Smaller neonates gain relatively more body mass during larval development. Entomological Science, 2019, 22, 373-380.	0.6	7
4	A method for rearing the yellow-spotted longicorn beetle, Psacothea hilaris (Coleoptera:) Tj ETQq0 0 0 rgBT /Ove	erlock 10 T	f 50 622 Td ((
5	Insights into How Longicorn Beetle Larvae Determine the Timing of Metamorphosis: Starvation-Induced Mechanism Revisited. PLoS ONE, 2016, 11, e0158831.	2.5	15
6	Cloning, phylogeny, and expression analysis of the Broad-Complex gene in the longicorn beetle Psacothea hilaris. SpringerPlus, 2014, 3, 539.	1.2	8
7	The strepsipteran parasite Elenchus japonicus (Strepsiptera, Elenchidae) of planthoppers consists of three genotypes. Applied Entomology and Zoology, 2011, 46, 435-442.	1.2	10
8	Differences in timing of the emergence of the overwintering generation between rice and water-oats populations of the striped stem borer moth, Chilo suppressalis (Lepidoptera: Crambidae). Applied Entomology and Zoology, 2009, 44, 485-489.	1.2	10
9	Upregulation of a desaturase is associated with the enhancement of cold hardiness in the onion maggot, Delia antiqua. Insect Biochemistry and Molecular Biology, 2007, 37, 1160-1167.	2.7	52
10	Occurrence and density of Halobates micans (Hemiptera: Gerridae) in the eastern South Indian Ocean. Entomological Science, 2007, 10, 213-215.	0.6	4
11	Distribution of the oceanic insects Halobates (Hemiptera: Gerridae) off the south coast of Japan. Entomological Science, 2004, 7, 351-357.	0.6	12
12	Ostrinia spp. in Japan: their host plants and sex pheromones. Entomologia Experimentalis Et Applicata, 1999, 91, 237-244.	1.4	111
13	Geographic Variation in Sex Pheromone of Asian Corn Borer, Ostrinia furnacalis, in Japan. Journal of Chemical Ecology, 1998, 24, 2079-2088.	1.8	48
14	Identification of the sex pheromone of Ostrinia palustralis. Entomologia Experimentalis Et Applicata, 1998, 86, 313-318.	1.4	15
15	Sex pheromone of the butterbur borer, Ostrinia zaguliaevi. Entomologia Experimentalis Et Applicata, 1998, 89, 281-287.	1.4	23
16	Female-biased sex ratio in the Asian corn borer, Ostrinia furnacalis: evidence for the occurrence of feminizing bacteria in an insect. Heredity, 1998, 81, 311-316.	2.6	49
17	Female-biased sex ratio in the Asian corn borer, Ostrinia furnacalis: evidence for the occurrence of feminizing bacteria in an insect. Heredity, 1998, 81, 311-316.	2.6	8
18	Allozyme polymorphism and geographic variation in the small brown planthopper, Laodelphax striatellus (Homoptera: Delphacidae). Biochemical Genetics, 1997, 35, 383-393.	1.7	19

#	Article	IF	CITATIONS
19	Identification of Sex Pheromone of Adzuki Bean Borer,Ostrinia scapulalis. Journal of Chemical Ecology, 1997, 23, 2791-2802.	1.8	34