

# Sonja M Kessler

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6507056/publications.pdf>

Version: 2024-02-01

41  
papers

1,224  
citations

279798

23  
h-index

377865

34  
g-index

44  
all docs

44  
docs citations

44  
times ranked

1941  
citing authors

#	ARTICLE	IF	CITATIONS
1	First Small-Molecule Inhibitors Targeting the RNA-Binding Protein IGF2BP2/IMP2 for Cancer Therapy. <i>ACS Chemical Biology</i> , 2022, 17, 361-375.	3.4	23
2	Kupffer cells are protective in alcoholic steatosis. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2022, 1868, 166398.	3.8	1
3	Chemical composition and biological activities of <i>Valeriana dioscoridis</i> SM. roots. <i>South African Journal of Botany</i> , 2021, 141, 306-312.	2.5	4
4	The Good, the Bad, the Question—H19 in Hepatocellular Carcinoma. <i>Cancers</i> , 2020, 12, 1261.	3.7	30
5	Thioholgamide A, a New Anti-Proliferative Anti-Tumor Agent, Modulates Macrophage Polarization and Metabolism. <i>Cancers</i> , 2020, 12, 1288.	3.7	22
6	The Insulin-Like Growth Factor 2 mRNA Binding Protein IMP2/IGF2BP2 is Overexpressed and Correlates with Poor Survival in Pancreatic Cancer. <i>International Journal of Molecular Sciences</i> , 2019, 20, 3204.	4.1	53
7	CRUP: a comprehensive framework to predict condition-specific regulatory units. <i>Genome Biology</i> , 2019, 20, 227.	8.8	26
8	The mRNA-binding Protein TTP/ZFP36 in Hepatocarcinogenesis and Hepatocellular Carcinoma. <i>Cancers</i> , 2019, 11, 1754.	3.7	20
9	Chemical composition and antioxidant, cytotoxic, and insecticidal potential of <i>Valeriana alliariifolia</i> in Turkey. <i>Arhiv Za Higijenu Rada I Toksikologiju</i> , 2019, 70, 207-218.	0.7	5
10	IGF2 mRNA Binding Protein 2 Transgenic Mice Are More Prone to Develop a Ductular Reaction and to Progress Toward Cirrhosis. <i>Frontiers in Medicine</i> , 2019, 6, 179.	2.6	12
11	Lack of Kupffer cell depletion in diethylnitrosamine-induced hepatic inflammation. <i>Journal of Hepatology</i> , 2019, 70, 813-815.	3.7	11
12	Diethylnitrosamine (DENa) recapitulates formation of hepatic angiosarcoma in pigs. <i>PLoS ONE</i> , 2019, 14, e0214756.	2.5	3
13	High Keratin 8/18 Ratio Predicts Aggressive Hepatocellular Cancer Phenotype. <i>Translational Oncology</i> , 2019, 12, 256-268.	3.7	28
14	Hepatocellular Carcinoma and Nuclear Paraspeckles: Induction in Chemoresistance and Prediction for Poor Survival. <i>Cellular Physiology and Biochemistry</i> , 2019, 52, 787-801.	1.6	29
15	Hsp72 protects against liver injury via attenuation of hepatocellular death, oxidative stress, and JNK signaling. <i>Journal of Hepatology</i> , 2018, 68, 996-1005.	3.7	51
16	Transgenic expression of the RNA binding protein IMP2 stabilizes miRNA targets in murine microsteatosis. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018, 1864, 3099-3108.	3.8	10
17	Hepatic interleukin-6 production is maintained during endotoxin tolerance and facilitates lipid accumulation. <i>Immunobiology</i> , 2017, 222, 786-796.	1.9	26
18	The long non-coding RNA H19 suppresses carcinogenesis and chemoresistance in hepatocellular carcinoma. <i>Cell Stress</i> , 2017, 1, 37-54.	3.2	50

#	ARTICLE	IF	CITATIONS
19	IMP2/IGF2BP2 expression, but not IMP1 and IMP3, predicts poor outcome in patients and high tumor growth rate in xenograft models of gallbladder cancer. <i>Oncotarget</i> , 2017, 8, 89736-89745.	1.8	30
20	Insulin Signaling Linking Metabolism and Malignancy. , 2017, , 61-75.		0
21	Susceptibility of Different Mouse Wild Type Strains to Develop Diet-Induced NAFLD/AFLD-Associated Liver Disease. <i>PLoS ONE</i> , 2016, 11, e0155163.	2.5	62
22	Transient Hepatic Overexpression of Insulin-Like Growth Factor 2 Induces Free Cholesterol and Lipid Droplet Formation. <i>Frontiers in Physiology</i> , 2016, 7, 147.	2.8	19
23	Small BODIPY Probes for Combined Dual <sup>19</sup> Fâ€¦MRI and Fluorescence Imaging. <i>ChemMedChem</i> , 2016, 11, 1568-1575.	3.2	16
24	Hepatic Deletion of Janus Kinase 2 Counteracts Oxidative Stress in Mice. <i>Scientific Reports</i> , 2016, 6, 34719.	3.3	24
25	Elevated expression of the <i>IGF2</i> mRNA binding protein 2 (IGF2BP2/IMP2) is linked to short survival and metastasis in esophageal adenocarcinoma. <i>Oncotarget</i> , 2016, 7, 49743-49750.	1.8	45
26	Insulin-Like Growth Factor 2 - The Oncogene and its Accomplices. <i>Current Pharmaceutical Design</i> , 2016, 22, 5948-5961.	1.9	26
27	Overexpression of <i>IGF2</i> mRNA-Binding Protein 2 (IMP2/p62) as a Feature of Basal-like Breast Cancer Correlates with Short Survival. <i>Scandinavian Journal of Immunology</i> , 2015, 82, 142-143.	2.7	35
28	Hepatic hepcidin expression is decreased in cirrhosis and HCC. <i>Journal of Hepatology</i> , 2015, 62, 977-979.	3.7	28
29	Glucocorticoid-Induced Leucine Zipper: A Critical Factor in Macrophage Endotoxin Tolerance. <i>Journal of Immunology</i> , 2015, 194, 6057-6067.	0.8	76
30	IMP2/p62 induces genomic instability and an aggressive hepatocellular carcinoma phenotype. <i>Cell Death and Disease</i> , 2015, 6, e1894-e1894.	6.3	64
31	Growth hormone resistance exacerbates cholestasis-induced murine liver fibrosis. <i>Hepatology</i> , 2015, 61, 613-626.	7.3	27
32	Non-alcoholic Fatty Liver Disease. , 2015, , 1-21.		0
33	Fatty Acid Elongation in Non-Alcoholic Steatohepatitis and Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2014, 15, 5762-5773.	4.1	45
34	The insulin-like growth factor 2 ( <i>IGF2</i> ) mRNA-binding protein p62/IGF2BP2-2 as a promoter of NAFLD and HCC?. <i>Gut</i> , 2014, 63, 861-863.	12.1	54
35	Lipid Metabolism Signatures in NASH-Associated HCCâ€”Letter. <i>Cancer Research</i> , 2014, 74, 2903-2904.	0.9	12
36	The IGF2 mRNA binding protein p62/IGF2BP2-2 induces fatty acid elongation as a critical feature of steatosis. <i>Journal of Lipid Research</i> , 2014, 55, 1087-1097.	4.2	42

#	ARTICLE	IF	CITATIONS
37	Downregulation of the glucocorticoid-induced leucine zipper (GILZ) promotes vascular inflammation. <i>Atherosclerosis</i> , 2014, 234, 391-400.	0.8	53
38	Elevated free cholesterol in a p62 overexpression model of non-alcoholic steatohepatitis. <i>World Journal of Gastroenterology</i> , 2014, 20, 17839-17850.	3.3	28
39	IGF2 mRNA binding protein p62/IMP2-2 in hepatocellular carcinoma: antiapoptotic action is independent of IGF2/PI3K signaling. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 304, G328-G336.	3.4	49
40	Rapid chromatographic method to decipher distinct alterations in lipid classes in NAFLD/NASH. <i>World Journal of Hepatology</i> , 2013, 5, 558.	2.0	22
41	Overexpression of the IGF2-mRNA binding protein p62 in transgenic mice induces a steatotic phenotype. <i>Journal of Hepatology</i> , 2011, 54, 994-1001.	3.7	56