

Teik Chye Ooi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6511523/publications.pdf>

Version: 2024-02-01

19
papers

583
citations

933264

10
h-index

940416

16
g-index

19
all docs

19
docs citations

19
times ranked

977
citing authors

#	ARTICLE	IF	CITATIONS
1	Plasma PCSK9 levels are significantly modified by statins and fibrates in humans. <i>Lipids in Health and Disease</i> , 2008, 7, 22.	1.2	187
2	Novel Loss-of-Function PCSK9 Variant Is Associated with Low Plasma LDL Cholesterol in a French-Canadian Family and with Impaired Processing and Secretion in Cell Culture. <i>Clinical Chemistry</i> , 2011, 57, 1415-1423.	1.5	101
3	Plasma PCSK9 levels correlate with cholesterol in men but not in women. <i>Biochemical and Biophysical Research Communications</i> , 2007, 361, 451-456.	1.0	82
4	Nutritional and Lipid Modulation of PCSK9: Effects on Cardiometabolic Risk Factors. <i>Journal of Nutrition</i> , 2017, 147, 473-481.	1.3	32
5	Metformin in adults with type 1 diabetes: Design and methods of the REDucing with Metformin Versus Ascarbose Lower Lesions (REMOVAL): an international multicentre trial. <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 509-516.	2.2	32
6	The Effect of PCSK9 Loss-of-Function Variants on the Postprandial Lipid and ApoB-Lipoprotein Response. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 3452-3460.	1.8	32
7	Human Serum PCSK9 Is Elevated at Parturition in Comparison to Nonpregnant Subjects While Serum PCSK9 from Umbilical Cord Blood is Lower Compared to Maternal Blood. <i>Isrn Endocrinology</i> , 2013, 2013, 1-8.	2.0	26
8	Effect of hemodialysis on extracellular vesicles and circulating submicron particles. <i>BMC Nephrology</i> , 2019, 20, 294.	0.8	19
9	Methods to Study Postprandial Lipemia. <i>Current Vascular Pharmacology</i> , 2011, 9, 302-308.	0.8	14
10	Using a Structured Discharge Letter Template to Improve Communication During the Transition from a Specialized Outpatient Diabetes Clinic to a Primary Care Physician. <i>Canadian Journal of Diabetes</i> , 2015, 39, 457-466.	0.4	14
11	Associations Between Soluble LDLR and Lipoproteins in a White Cohort and the Effect of PCSK9 Loss-of-Function. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 3486-3495.	1.8	14
12	Proposing a 'Lipemic Index' As a Nutritional and Research Tool. <i>Current Vascular Pharmacology</i> , 2011, 9, 313-317.	0.8	8
13	Patient Perspectives on Discharge from Specialist Type 2 Diabetes Care Back to Primary Care: A Qualitative Study. <i>Canadian Journal of Diabetes</i> , 2014, 38, 191-197.	0.4	6
14	Relative effect of hypertriglyceridemia on non-HDLc and apolipoprotein B as cardiovascular disease risk markers. <i>Journal of Clinical Lipidology</i> , 2020, 14, 825-836.	0.6	6
15	Calculated Non-HDL Cholesterol Includes Cholesterol in Larger Triglyceride-Rich Lipoproteins in Hypertriglyceridemia. <i>Journal of the Endocrine Society</i> , 2020, 4, bvz010.	0.1	4
16	Circulating PCSK9 is lowered acutely following surgery. <i>Journal of Clinical Laboratory Analysis</i> , 2018, 32, e22358.	0.9	3
17	PCSK9 as a Biomarker of Cardiovascular Disease. , 2016, , 125-151.		1
18	PCSK9 as a Biomarker of Cardiovascular Disease. , 2015, , 1-27.		1

#	ARTICLE	IF	CITATIONS
19	A More Atherogenic Lipoprotein Status Is Present in Adults With Type 2 Diabetes Mellitus Than in Those Without With Equivalent Degrees of Hypertriglyceridemia. Canadian Journal of Diabetes, 2022, 46, 480-486.	0.4	1