Cleonice A M Bento

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6517544/publications.pdf

Version: 2024-02-01

42 papers 1,009 citations

394421 19 h-index 30 g-index

42 all docs 42 docs citations

42 times ranked 1587 citing authors

#	Article	IF	Citations
1	Elevated proportion of TLR2- and TLR4-expressing Th17-like cells and activated memory B cells was associated with clinical activity of cerebral cavernous malformations. Journal of Neuroinflammation, 2022, 19, 28.	7.2	6
2	Leptin favors Th $17/\text{Treg}$ cell subsets imbalance associated with allergic asthma severity. Clinical and Translational Allergy, 2022, 12, .	3.2	11
3	Human pregnancy levels of estrogen and progesterone contribute to humoral immunity by activating T _{FH} /B cell axis. European Journal of Immunology, 2021, 51, 167-179.	2.9	13
4	Selective serotonin reuptake inhibitor attenuates the hyperresponsiveness of TLR2 ⁺ and TLR4 ⁺ Th17/Tc17â€like cells in multiple sclerosis patients with major depression. Immunology, 2021, 162, 290-305.	4.4	17
5	Pregnancy favors circulating ILâ€21–secreting T FH â€like cell recovery in ARVâ€treated HIVâ€1–infected women. American Journal of Reproductive Immunology, 2020, 83, e13204.	1.2	1
6	Is Asian type MS an MS phenotype, an NMO spectrum disorder, or a MOG-IgG related disease?. Multiple Sclerosis and Related Disorders, 2020, 42, 102082.	2.0	8
7	Different core-specific T cell subsets are expanded in chronic hepatitis C with advanced liver disease. Cytokine, 2019, 124, 154456.	3.2	5
8	TLR-2 and TLR-4 agonists favor expansion of CD4+ T cell subsets implicated in the severity of neuromyelitis optica spectrum disorders. Multiple Sclerosis and Related Disorders, 2019, 34, 66-76.	2.0	12
9	Serum leptin levels correlate negatively with the capacity of vitamin D to modulate the in vitro cytokines production by CD4+ T cells in asthmatic patients. Clinical Immunology, 2019, 205, 93-105.	3.2	9
10	The expansion of circulating IL-6 and IL-17-secreting follicular helper T cells is associated with neurological disabilities in neuromyelitis optica spectrum disorders. Journal of Neuroimmunology, 2019, 330, 12-18.	2.3	25
11	Serotonin decreases the production of Th1/Th17 cytokines and elevates the frequency of regulatory CD4 ⁺ Tâ€eell subsets in multiple sclerosis patients. European Journal of Immunology, 2018, 48, 1376-1388.	2.9	58
12	Different interleukinâ€17â€secreting Tollâ€like receptor ⁺ Tâ€cell subsets are associated with disease activity in multiple sclerosis. Immunology, 2018, 154, 239-252.	4.4	20
13	Reactivation of latent HIV-1 in vitro using an ethanolic extract from Euphorbia umbellata (Euphorbiaceae) latex. PLoS ONE, 2018, 13, e0207664.	2.5	6
14	Lower frequency of antibodies to MOG in Brazilian patients with demyelinating diseases: An ethnicity influence?. Multiple Sclerosis and Related Disorders, 2018, 25, 87-94.	2.0	16
15	Cognitive changes in nurses working in intensive care units. Revista Brasileira De Enfermagem, 2018, 71, 73-79.	0.7	20
16	Case Report: Acute Transverse Myelitis after Zika Virus Infection. American Journal of Tropical Medicine and Hygiene, 2018, 99, 1419-1421.	1.4	21
17	The proportion of different interleukinâ€17â€producing Tâ€cell subsets is associated with liver fibrosis in chronic hepatitis C. Immunology, 2017, 151, 167-176.	4.4	15
18	Expansion of IL-6+ Th17-like cells expressing TLRs correlates with microbial translocation and neurological disabilities in NMOSD patients. Journal of Neuroimmunology, 2017, 307, 82-90.	2.3	14

#	Article	IF	CITATIONS
19	Pregnancy favors the expansion of circulating functional follicular helper T Cells. Journal of Reproductive Immunology, 2017, 121, 1-10.	1.9	27
20	Fatigue favors in vitro Th1 and Th17-like cell expansion and reduces corticoid sensitivity in MS patients. Journal of Neuroimmunology, 2017, 303, 81-89.	2.3	17
21	The effects of roasted yerba mate (llex paraguariensis A. ST. Hil.) consumption on glycemia and total serum creatine phosphokinase in patients with traumatic brain injury. Journal of Functional Foods, 2017, 28, 240-245.	3.4	12
22	Interleukinâ€17―and interleukinâ€22â€secreting myelinâ€specific <scp>CD</scp> 4 ⁺ T cells resista to corticoids are related with active brain lesions in multiple sclerosis patients. Immunology, 2016, 147, 212-220.	ant 4.4	37
23	Prediction of disease severity in neuromyelitis optica by the levels of interleukin (IL)-6 produced during remission phase. Clinical and Experimental Immunology, 2016, 183, 480-489.	2.6	60
24	Vitamin D modulates different IL-17-secreting T cell subsets in multiple sclerosis patients. Journal of Neuroimmunology, 2016, 299, 8-18.	2.3	47
25	Combined exercise training reduces fatigue and modulates the cytokine profile of T-cells from multiple sclerosis patients in response to neuromediators. Journal of Neuroimmunology, 2016, 293, 91-99.	2.3	54
26	Poor functional immune recovery in aged HIV-1-infected patients following successfully treatment with antiretroviral therapy. Human Immunology, 2015, 76, 701-710.	2.4	7
27	Endogenous interleukinâ€6 amplifies interleukinâ€17 production and corticoidâ€resistance in peripheral <scp>T</scp> cells from patients with multiple sclerosis. Immunology, 2014, 143, 560-568.	4.4	27
28	Dopamine favors expansion of glucocorticoid-resistant IL-17-producing T cells in multiple sclerosis. Brain, Behavior, and Immunity, 2014, 41, 182-190.	4.1	35
29	The Ex Vivo Production of IL-6 and IL-21 by CD4+ T Cells is Directly Associated with Neurological Disability in Neuromyelitis Optica Patients. Journal of Clinical Immunology, 2013, 33, 179-189.	3.8	64
30	High in vitro immune reactivity to Escherichia coli in neuromyelitis optica patients is correlated with both neurological disabilities and elevated plasma lipopolysaccharide levels. Human Immunology, 2013, 74, 1080-1087.	2.4	23
31	The impact of maternal anti-retroviral therapy on cytokine profile in the uninfected neonates. Human Immunology, 2013, 74, 1051-1056.	2.4	8
32	Low sensitivity to glucocorticoid inhibition of in vitro Th17-related cytokine production in multiple sclerosis patients is related to elevated plasma lipopolysaccharide levels. Clinical Immunology, 2013, 148, 209-218.	3.2	28
33	High IL-10 production by aged AIDS patients is related to high frequency of Tr-1 phenotype and low in vitro viral replication. Clinical Immunology, 2012, 145, 31-43.	3.2	10
34	The impact of pregnancy on the HIV-1-specific T cell function in infected pregnant women. Clinical Immunology, 2012, 145, 177-188.	3.2	10
35	Dopamine up-regulates Th17 phenotype from individuals with generalized anxiety disorder. Journal of Neuroimmunology, 2011, 238, 58-66.	2.3	36
36	Substance P Enhances Th17 Phenotype in Individuals with Generalized Anxiety Disorder: an Event Resistant to Glucocorticoid Inhibition. Journal of Clinical Immunology, 2011, 31, 51-59.	3.8	20

#	Article	IF	CITATION
37	Enhanced Th17 phenotype in individuals with generalized anxiety disorder. Journal of Neuroimmunology, 2010, 229, 212-218.	2.3	85
38	Failure of Highly Active Antiretroviral Therapy in Reconstituting Immune Response to Clostridium tetani Vaccine in Aged AIDS Patients. Journal of Acquired Immune Deficiency Syndromes (1999), 2010, 54, 10-17.	2.1	8
39	IL-10-secreting T cells from HIV-infected pregnant women downregulate HIV-1 replication: effect enhanced by antiretroviral treatment. Aids, 2009, 23, 9-18.	2.2	29
40	Altered immunological reactivity in HIV-1-exposed uninfected neonates. Clinical Immunology, 2008, 127, 340-347.	3.2	53
41	Interleukin-10-secreting CD4 cells from aged patients with AIDS decrease in-vitro HIV replication and tumour necrosis factor \hat{l}_{\pm} production. Aids, 2007, 21, 1763-1770.	2.2	11
42	Glucuronoxylomannan of Cryptococcus neoformans exacerbates in vitro yeast cell growth by interleukin 10-dependent inhibition of CD4+ T lymphocyte responses. Cellular Immunology, 2003, 222, 116-125.	3.0	24