## Yi-Chun Wu

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6521251/publications.pdf

Version: 2024-02-01

42 papers

3,635 citations

257450 24 h-index 289244 40 g-index

42 all docs 42 docs citations

42 times ranked 4459 citing authors

#	Article	IF	CITATIONS
1	In Vivo Imaging and Toxicity Assessments of Fluorescent Nanodiamonds in <i>Caenorhabditis elegans</i> . Nano Letters, 2010, 10, 3692-3699.	9.1	514
2	DNAâ^Gold Nanorod Conjugates for Remote Control of Localized Gene Expression by near Infrared Irradiation. Journal of the American Chemical Society, 2006, 128, 3709-3715.	13.7	411
3	C. elegans phagocytosis and cell-migration protein CED-5 is similar to human DOCK180. Nature, 1998, 392, 501-504.	27.8	346
4	Selective Binding of Mannose-Encapsulated Gold Nanoparticles to Type 1 Pili inEscherichia coli. Journal of the American Chemical Society, 2002, 124, 3508-3509.	13.7	280
5	The C. elegans Cell Corpse Engulfment Gene ced-7 Encodes a Protein Similar to ABC Transporters. Cell, 1998, 93, 951-960.	28.9	275
6	Phagocytosis of Apoptotic Cells Is Regulated by a UNC-73/TRIO-MIG-2/RhoG Signaling Module and Armadillo Repeats of CED-12/ELMO. Current Biology, 2004, 14, 2208-2216.	3.9	185
7	Cell Corpse Engulfment Mediated by C. elegans Phosphatidylserine Receptor Through CED-5 and CED-12. Science, 2003, 302, 1563-1566.	12.6	183
8	Preparation of Fluorescent Silica Nanotubes and Their Application in Gene Delivery. Advanced Materials, 2005, 17, 404-407.	21.0	166
9	C. elegans CED-12 Acts in the Conserved CrkII/DOCK180/Rac Pathway to Control Cell Migration and Cell Corpse Engulfment. Developmental Cell, 2001, 1, 491-502.	7.0	160
10	NUC-1, a <i>Caenorhabditis elegans</i> DNase II homolog, functions in an intermediate step of DNA degradation during apoptosis. Genes and Development, 2000, 14, 536-548.	5.9	146
11	Quantitative analysis of multivalent interactions of carbohydrate-encapsulated gold nanoparticles with concanavalin AElectronic supplementary information (ESI) available: detailed experimental procedures, SPR response curves and compound characterization data. See http://www.rsc.org/suppdata/cc/b3/b308995a/. Chemical Communications, 2003, , 2920.	4.1	125
12	Programmed Cell Death During <i>Caenorhabditis elegans </i> Development. Genetics, 2016, 203, 1533-1562.	2.9	88
13	Fluorescent nanodiamond as a probe for the intercellular transport ofÂproteins inÂvivo. Biomaterials, 2013, 34, 8352-8360.	11.4	83
14	Engulfment of Apoptotic Cells in C. elegans Is Mediated by Integrin α/SRC Signaling. Current Biology, 2010, 20, 477-486.	3.9	82
15	Fluorescence-Guided Probes of Aptamer-Targeted Gold Nanoparticles with Computed Tomography Imaging Accesses for in Vivo Tumor Resection. Scientific Reports, 2015, 5, 15675.	3.3	73
16	Negative regulation of <i>Caenorhabditis elegans</i> epidermal damage responses by death-associated protein kinase. Proceedings of the National Academy of Sciences of the United States of America, 2009, 106, 1457-1461.	7.1	70
17	Distinct Rac Activation Pathways Control Caenorhabditis elegans Cell Migration and Axon Outgrowth. Developmental Biology, 2002, 250, 145-155.	2.0	63
18	<i>Diacylglycerol lipase</i> regulates lifespan and oxidative stress response by inversely modulating <scp>TOR</scp> signaling in <i><scp>D</scp>rosophila</i> and <i><scp>C</scp>.Âelegans</i> Aging Cell, 2014, 13, 755-764.	6.7	53

#	Article	IF	CITATIONS
19	Noise propagation with interlinked feed-forward pathways. Scientific Reports, 2016, 6, 23607.	3.3	36
20	Restriction of vaccinia virus replication by a ced-3 and ced-4-dependent pathway in Caenorhabditis elegans. Proceedings of the National Academy of Sciences of the United States of America, 2006, 103, 4174-4179.	7.1	34
21	Fluorescence lifetime imaging microscopy of nanodiamonds in vivo. Proceedings of SPIE, 2013, , .	0.8	33
22	Integrin α PAT-2/CDC-42 Signaling Is Required for Muscle-Mediated Clearance of Apoptotic Cells in Caenorhabditis elegans. PLoS Genetics, 2012, 8, e1002663.	3.5	29
23	elF3k regulates apoptosis in epithelial cells by releasing caspase 3 from keratin-containing inclusions. Journal of Cell Science, 2008, 121, 2382-2393.	2.0	28
24	BLMP-1/Blimp-1 Regulates the Spatiotemporal Cell Migration Pattern in C. elegans. PLoS Genetics, 2014, 10, e1004428.	3.5	27
25	Length-dependent toxicity of untranslated CUG repeats on Caenorhabditis elegans. Biochemical and Biophysical Research Communications, 2007, 352, 774-779.	2.1	26
26	LIN-3/EGF Promotes the Programmed Cell Death of Specific Cells in Caenorhabditis elegans by Transcriptional Activation of the Pro-apoptotic Gene egl-1. PLoS Genetics, 2014, 10, e1004513.	3.5	18
27	Engulfing cells promote neuronal regeneration and remove neuronal debris through distinct biochemical functions of CED-1. Nature Communications, 2018, 9, 4842.	12.8	15
28	Analysis of Programmed Cell Death in the Nematode Caenorhabditis elegans. Methods in Enzymology, 2000, 322, 76-88.	1.0	12
29	Growth-dependent effect of muscleblind knockdown on Caenorhabditis elegans. Biochemical and Biophysical Research Communications, 2008, 366, 705-709.	2.1	12
30	Early developmental nanoplastics exposure disturbs circadian rhythms associated with stress resistance decline and modulated by DAF-16 and PRDX-2 in C. elegans. Journal of Hazardous Materials, 2022, 423, 127091.	12.4	9
31	<b>Fluorescent Nanodiamond – A Novel Nanomaterial for<i>In Vivo</i>Applications</b> . Materials Research Society Symposia Proceedings, 2011, 1362, 1.	0.1	8
32	Methods for Studying Programmed Cell Death in C. elegans. Methods in Cell Biology, 2012, 107, 295-320.	1.1	7
33	C. elegans EIF-3.K Promotes Programmed Cell Death through CED-3 Caspase. PLoS ONE, 2012, 7, e36584.	2.5	7
34	Ultrasensitive Detection of Alzheimer's Amyloids on a Plasmonic-Gold Platform. ACS Applied Materials & Lamp; Interfaces, 2021, 13, 57036-57042.	8.0	7
35	Quantum-dot-embedded silica nanotubes as nanoprobes for simple and sensitive DNA detection. Nanotechnology, 2011, 22, 155102.	2.6	6
36	Functional characterization of the meiosis-specific DNA double-strand break inducing factor SPO-11 from C. elegans. Scientific Reports, 2017, 7, 2370.	3.3	6

#	Article	IF	CITATIONS
37	Caenorhabditis elegans TLK-1 controls cytokinesis by localizing AIR-2/Aurora B to midzone microtubules. Biochemical and Biophysical Research Communications, 2010, 400, 187-193.	2.1	4
38	Development of a water refractive index-matched microneedle integrated into a light sheet microscopy system for continuous embryonic cell imaging. Lab on A Chip, 2022, 22, 584-591.	6.0	3
39	BLMP-1 promotes developmental cell death in C. elegans by timely repression of ced-9/bcl-2 transcription. Development (Cambridge), 2021, 148, .	2.5	2
40	C.Âelegans BLMP-1 controls apical epidermal cell morphology by repressing expression of mannosyltransferase bus-8 and molting signal mlt-8. Developmental Biology, 2022, 486, 96-108.	2.0	2
41	Programmed Cell Death in C. elegans. , 2009, , 355-373.		1
42	Programmed Cell Death in C. elegans. , 2003, , 135-144.		0