

Yiping Chen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6524782/publications.pdf>

Version: 2024-02-01

87
papers

4,203
citations

101543

36
h-index

123424

61
g-index

92
all docs

92
docs citations

92
times ranked

4011
citing authors

#	ARTICLE	IF	CITATIONS
1	FGF8-mediated signaling regulates tooth developmental pace during odontogenesis. <i>Journal of Genetics and Genomics</i> , 2022, 49, 40-53.	3.9	4
2	Discovery and functional assessment of a novel adipocyte population driven by intracellular Wnt/ β -catenin signaling in mammals. <i>ELife</i> , 2022, 11, .	6.0	5
3	Single-cell transcriptomic signatures and gene regulatory networks modulated by WIs in mammalian midline facial formation and clefts. <i>Development (Cambridge)</i> , 2022, 149, .	2.5	6
4	The Transcription Factor Shox2 Shapes Neuron Firing Properties and Suppresses Seizures by Regulation of Key Ion Channels in Thalamocortical Neurons. <i>Cerebral Cortex</i> , 2021, 31, 3194-3212.	2.9	2
5	A systematic dissection of human primary osteoblasts in vivo at single-cell resolution. <i>Aging</i> , 2021, 13, 20629-20650.	3.1	19
6	Single-cell RNA sequencing deconvolutes the <i>in vivo</i> heterogeneity of human bone marrow-derived mesenchymal stem cells. <i>International Journal of Biological Sciences</i> , 2021, 17, 4192-4206.	6.4	39
7	Olig2 regulates terminal differentiation and maturation of peripheral olfactory sensory neurons. <i>Cellular and Molecular Life Sciences</i> , 2020, 77, 3597-3609.	5.4	8
8	Exogenous FGF8 signaling in osteocytes leads to mandibular hypoplasia in mice. <i>Oral Diseases</i> , 2020, 26, 590-596.	3.0	4
9	Conjugated activation of myocardial-specific transcription of <i>Cja5</i> by a pair of <i>Nkx2-5-Shox2</i> co-responsive elements. <i>Developmental Biology</i> , 2020, 465, 79-87.	2.0	2
10	Cellular and developmental basis of orofacial clefts. <i>Birth Defects Research</i> , 2020, 112, 1558-1587.	1.5	40
11	The transcriptional regulator MEIS2 sets up the ground state for palatal osteogenesis in mice. <i>Journal of Biological Chemistry</i> , 2020, 295, 5449-5460.	3.4	15
12	<i>Nkx2-5</i> defines a subpopulation of pacemaker cells and is essential for the physiological function of the sinoatrial node in mice. <i>Development (Cambridge)</i> , 2019, 146, .	2.5	23
13	<i>Shox2</i> regulates osteogenic differentiation and pattern formation during hard palate development in mice. <i>Journal of Biological Chemistry</i> , 2019, 294, 18294-18305.	3.4	17
14	Regrowing a tooth: in vitro and in vivo approaches. <i>Current Opinion in Cell Biology</i> , 2019, 61, 126-131.	5.4	14
15	Opposing roles of TCF7/LEF1 and TCF7L2 in cyclin D2 and <i>Bmp4</i> expression and cardiomyocyte cell cycle control during late heart development. <i>Laboratory Investigation</i> , 2019, 99, 807-818.	3.7	20
16	Conditional deletion of <i>Bmp2</i> in cranial neural crest cells recapitulates Pierre Robin sequence in mice. <i>Cell and Tissue Research</i> , 2019, 376, 199-210.	2.9	30
17	TGF- β 2 signaling inhibits canonical BMP signaling pathway during palate development. <i>Cell and Tissue Research</i> , 2018, 371, 283-291.	2.9	26
18	Efficient induction of functional ameloblasts from human keratinocyte stem cells. <i>Stem Cell Research and Therapy</i> , 2018, 9, 126.	5.5	16

#	ARTICLE	IF	CITATIONS
19	ISLET1-Dependent β -Catenin/Hedgehog Signaling Is Required for Outgrowth of the Lower Jaw. <i>Molecular and Cellular Biology</i> , 2017, 37, .	2.3	19
20	A unique stylopod patterning mechanism by <i>Shox2</i> controlled osteogenesis. <i>Development</i> (Cambridge), 2016, 143, 2548-60.	2.5	15
21	LDL Receptor-Related Protein 6 Modulates Ret Proto-Oncogene Signaling in Renal Development and Cystic Dysplasia. <i>Journal of the American Society of Nephrology: JASN</i> , 2016, 27, 417-427.	6.1	12
22	Augmented Indian hedgehog signaling in cranial neural crest cells leads to craniofacial abnormalities and dysplastic temporomandibular joint in mice. <i>Cell and Tissue Research</i> , 2016, 364, 105-115.	2.9	7
23	Genetic Regulation of Sinoatrial Node Development and Pacemaker Program in the Venous Pole. <i>Journal of Cardiovascular Development and Disease</i> , 2015, 2, 282-298.	1.6	26
24	Altered FGF Signaling Pathways Impair Cell Proliferation and Elevation of Palate Shelves. <i>PLoS ONE</i> , 2015, 10, e0136951.	2.5	19
25	Reply to Kelder et al.: Does the Dorsal Mesenchymal Protrusion Act as a Temporary Pacemaker during Heart Development?. <i>Journal of Biological Chemistry</i> , 2015, 290, 8015.	3.4	0
26	The Short Stature Homeobox 2 (<i>Shox2</i>)-bone Morphogenetic Protein (BMP) Pathway Regulates Dorsal Mesenchymal Protrusion Development and Its Temporary Function as a Pacemaker during Cardiogenesis. <i>Journal of Biological Chemistry</i> , 2015, 290, 2007-2023.	3.4	26
27	Identification and analysis of a novel <i>bmp4</i> enhancer in Fugu genome. <i>Archives of Oral Biology</i> , 2015, 60, 540-545.	1.8	2
28	FGF8 signaling sustains progenitor status and multipotency of cranial neural crest-derived mesenchymal cells <i>in vivo</i> and <i>in vitro</i> . <i>Journal of Molecular Cell Biology</i> , 2015, 7, 441-454.	3.3	28
29	A common <i>Shox2</i> - <i>Nkx2-5</i> antagonistic mechanism primes the pacemaking cell fate in the pulmonary vein myocardium and sinoatrial node. <i>Development</i> (Cambridge), 2015, 142, 2521-32.	2.5	105
30	Persistent Noggin arrests cardiomyocyte morphogenesis and results in early in utero lethality. <i>Developmental Dynamics</i> , 2015, 244, 457-467.	1.8	5
31	The non-canonical BMP and Wnt/ β -catenin signaling pathways orchestrate early tooth development. <i>Development</i> (Cambridge), 2015, 142, 128-139.	2.5	60
32	BMPRIA Mediated Signaling Is Essential for Temporomandibular Joint Development in Mice. <i>PLoS ONE</i> , 2014, 9, e101000.	2.5	33
33	Expression patterns of genes critical for BMP signaling pathway in developing human primary tooth germs. <i>Histochemistry and Cell Biology</i> , 2014, 142, 657-665.	1.7	18
34	An Atypical Canonical Bone Morphogenetic Protein (BMP) Signaling Pathway Regulates Msh Homeobox 1 (<i>Msx1</i>) Expression during Odontogenesis. <i>Journal of Biological Chemistry</i> , 2014, 289, 31492-31502.	3.4	28
35	BMP-FGF Signaling Axis Mediates Wnt-Induced Epidermal Stratification in Developing Mammalian Skin. <i>PLoS Genetics</i> , 2014, 10, e1004687.	3.5	66
36	Phosphorylation of <i>Shox2</i> Is Required for Its Function to Control Sinoatrial Node Formation. <i>Journal of the American Heart Association</i> , 2014, 3, e000796.	3.7	16

#	ARTICLE	IF	CITATIONS
37	Directed Bmp4 expression in neural crest cells generates a genetic model for the rare human bony synnathia birth defect. <i>Developmental Biology</i> , 2014, 391, 170-181.	2.0	39
38	Replacing Shox2 with human SHOX leads to congenital disc degeneration of the temporomandibular joint in mice. <i>Cell and Tissue Research</i> , 2014, 355, 345-354.	2.9	17
39	Precise chronology of differentiation of developing human primary dentition. <i>Histochemistry and Cell Biology</i> , 2014, 141, 221-227.	1.7	12
40	Bioengineering of a human whole tooth: progress and challenge. <i>Cell Regeneration</i> , 2014, 3, 3:8.	2.6	15
41	<i>Pitx2</i> -microRNA pathway that delimits sinoatrial node development and inhibits predisposition to atrial fibrillation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 9181-9186.	7.1	109
42	Pten Loss Induces Autocrine FGF Signaling to Promote Skin Tumorigenesis. <i>Cell Reports</i> , 2014, 6, 818-826.	6.4	44
43	Expression of SHH signaling molecules in the developing human primary dentition. <i>BMC Developmental Biology</i> , 2013, 13, 11.	2.1	28
44	Exploring the effects of gene dosage on mandible shape in mice as a model for studying the genetic basis of natural variation. <i>Development Genes and Evolution</i> , 2013, 223, 279-287.	0.9	34
45	Enhanced BMP signaling prevents degeneration and leads to endochondral ossification of Meckel's cartilage in mice. <i>Developmental Biology</i> , 2013, 381, 301-311.	2.0	43
46	Mice with Tak1 Deficiency in Neural Crest Lineage Exhibit Cleft Palate Associated with Abnormal Tongue Development. <i>Journal of Biological Chemistry</i> , 2013, 288, 10440-10450.	3.4	50
47	FGF signaling sustains the odontogenic fate of dental mesenchyme by suppressing β -catenin signaling. <i>Development (Cambridge)</i> , 2013, 140, 4375-4385.	2.5	34
48	Generation of <i>Shox2</i> ^{Cre} allele for tissue specific manipulation of genes in the developing heart, palate, and limb. <i>Genesis</i> , 2013, 51, 515-522.	1.6	36
49	Intra-epithelial Requirement of Canonical Wnt Signaling for Tooth Morphogenesis. <i>Journal of Biological Chemistry</i> , 2013, 288, 12080-12089.	3.4	48
50	Augmented BMPRIA-Mediated BMP Signaling in Cranial Neural Crest Lineage Leads to Cleft Palate Formation and Delayed Tooth Differentiation. <i>PLoS ONE</i> , 2013, 8, e66107.	2.5	34
51	The Role of Shox2 in SAN Development and Function. <i>Pediatric Cardiology</i> , 2012, 33, 882-889.	1.3	32
52	Bmpr1a is required in mesenchymal tissue and has limited redundant function with Bmpr1b in tooth and palate development. <i>Developmental Biology</i> , 2011, 349, 451-461.	2.0	68
53	Epithelial Wnt/ β -catenin signaling regulates palatal shelf fusion through regulation of Tgfb3 expression. <i>Developmental Biology</i> , 2011, 350, 511-519.	2.0	83
54	Ectopic expression of Nkx2.5 suppresses the formation of the sinoatrial node in mice. <i>Developmental Biology</i> , 2011, 356, 359-369.	2.0	66

#	ARTICLE	IF	CITATIONS
55	Wnt5a regulates growth, patterning, and odontoblast differentiation of developing mouse tooth. <i>Developmental Dynamics</i> , 2011, 240, 432-440.	1.8	78
56	Exogenous fibroblast growth factor 8 rescues development of mouse diastemal vestigial tooth ex vivo. <i>Developmental Dynamics</i> , 2011, 240, 1344-1353.	1.8	13
57	Tissue interaction is required for glenoid fossa development during temporomandibular joint formation. <i>Developmental Dynamics</i> , 2011, 240, 2466-2473.	1.8	40
58	Functional Redundancy between Human SHOX and Mouse Shox2 Genes in the Regulation of Sinoatrial Node Formation and Pacemaking Function. <i>Journal of Biological Chemistry</i> , 2011, 286, 17029-17038.	3.4	44
59	Overexpression of constitutively active BMP-receptor-1B in mouse skin causes an ichthyosis-vulgaris-like disease. <i>Cell and Tissue Research</i> , 2010, 342, 401-410.	2.9	10
60	<i>Gsk3β</i> is required in the epithelium for palatal elevation in mice. <i>Developmental Dynamics</i> , 2010, 239, 3235-3246.	1.8	36
61	Genetic interactions between Pax9 and Msx1 regulate lip development and several stages of tooth morphogenesis. <i>Developmental Biology</i> , 2010, 340, 438-449.	2.0	125
62	Induction of human keratinocytes into enamel-secreting ameloblasts. <i>Developmental Biology</i> , 2010, 344, 795-799.	2.0	48
63	Modulation of BMP signaling by Noggin is required for the maintenance of palatal epithelial integrity during palatogenesis. <i>Developmental Biology</i> , 2010, 347, 109-121.	2.0	93
64	Shox2 is essential for the differentiation of cardiac pacemaker cells by repressing Nkx2-5. <i>Developmental Biology</i> , 2009, 327, 376-385.	2.0	209
65	Hand2 is required in the epithelium for palatogenesis in mice. <i>Developmental Biology</i> , 2009, 330, 131-141.	2.0	68
66	Wnt5a regulates directional cell migration and cell proliferation via Ror2-mediated noncanonical pathway in mammalian palatogenesis. <i>FASEB Journal</i> , 2009, 23, 308.4.	0.5	0
67	Mouse embryonic diastema region is an ideal site for the development of ectopically transplanted tooth germ. <i>Developmental Dynamics</i> , 2008, 237, 411-416.	1.8	10
68	Mice with an anterior cleft of the palate survive neonatal lethality. <i>Developmental Dynamics</i> , 2008, 237, 1509-1516.	1.8	25
69	Shox2-deficiency leads to dysplasia and ankylosis of the temporomandibular joint in mice. <i>Mechanisms of Development</i> , 2008, 125, 729-742.	1.7	61
70	Wnt5a regulates directional cell migration and cell proliferation via Ror2-mediated noncanonical pathway in mammalian palate development. <i>Development (Cambridge)</i> , 2008, 135, 3871-3879.	2.5	200
71	Shox2 is required for chondrocyte proliferation and maturation in proximal limb skeleton. <i>Developmental Biology</i> , 2007, 306, 549-559.	2.0	73
72	Expression survey of genes critical for tooth development in the human embryonic tooth germ. <i>Developmental Dynamics</i> , 2007, 236, 1307-1312.	1.8	53

#	ARTICLE	IF	CITATIONS
73	Application of lentivirus-mediated RNAi in studying gene function in mammalian tooth development. <i>Developmental Dynamics</i> , 2006, 235, 1347-1357.	1.8	52
74	Shox2-deficient mice exhibit a rare type of incomplete clefting of the secondary palate. <i>Development (Cambridge)</i> , 2005, 132, 4397-4406.	2.5	133
75	The cellular and molecular etiology of the cleft secondary palate in Fgf10 mutant mice. <i>Developmental Biology</i> , 2005, 277, 102-113.	2.0	117
76	Chick <i>Pcl2</i> regulates the left-right asymmetry by repressing <i>Shh</i> expression in Hensen's node. <i>Development (Cambridge)</i> , 2004, 131, 4381-4391.	2.5	32
77	Timing of odontogenic neural crest cell migration and tooth-forming capability in mice. <i>Developmental Dynamics</i> , 2003, 226, 713-718.	1.8	41
78	<i>Msx1/Bmp4</i> genetic pathway regulates mammalian alveolar bone formation via induction of <i>Dlx5</i> and <i>Cbfa1</i> . <i>Mechanisms of Development</i> , 2003, 120, 1469-1479.	1.7	53
79	Rescue of cleft palate in <i>Msx1</i> -deficient mice by transgenic <i>Bmp4</i> reveals a network of BMP and <i>Shh</i> signaling in the regulation of mammalian palatogenesis. <i>Development (Cambridge)</i> , 2002, 129, 4135-4146.	2.5	332
80	Rescue of cleft palate in <i>Msx1</i> -deficient mice by transgenic <i>Bmp4</i> reveals a network of BMP and <i>Shh</i> signaling in the regulation of mammalian palatogenesis. <i>Development (Cambridge)</i> , 2002, 129, 4135-46.	2.5	175
81	Evidence for the differential regulation of <i>Nkx-6.1</i> expression in the ventral spinal cord and foregut by <i>Shh</i> -dependent and -independent mechanisms. <i>Genesis</i> , 2000, 27, 6-11.	1.6	19
82	Antagonistic Signals between BMP4 and FGF8 Define the Expression of <i>Pitx1</i> and <i>Pitx2</i> in Mouse Tooth-Forming Anlage. <i>Developmental Biology</i> , 2000, 217, 323-332.	2.0	183
83	Targeted Misexpression of Constitutively Active BMP Receptor-IB Causes Bifurcation, Duplication, and Posterior Transformation of Digit in Mouse Limb. <i>Developmental Biology</i> , 2000, 220, 154-167.	2.0	45
84	Transgenically ectopic expression of <i>Bmp4</i> to the <i>Msx1</i> mutant dental mesenchyme restores downstream gene expression but represses <i>Shh</i> and <i>Bmp2</i> in the enamel knot of wild type tooth germ. <i>Mechanisms of Development</i> , 2000, 99, 29-38.	1.7	87
85	<i>Msx1</i> is required for the induction of <i>Patched</i> by <i>Sonic hedgehog</i> in the mammalian tooth germ. <i>Developmental Dynamics</i> , 1999, 215, 45-53.	1.8	76
86	Expression and regulation of the chicken <i>Nkx-6.2</i> homeobox gene suggest its possible involvement in the ventral neural patterning and cell fate specification. , 1999, 216, 459-468.		19
87	Shaping limbs by apoptosis. <i>The Journal of Experimental Zoology</i> , 1998, 282, 691-702.	1.4	81