

Yiru Fang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6582088/publications.pdf>

Version: 2024-02-01

171
papers

8,190
citations

109321

35
h-index

58581

82
g-index

173
all docs

173
docs citations

173
times ranked

17945
citing authors

#	ARTICLE	IF	CITATIONS
1	Peripheral biomarkers to predict the diagnosis of bipolar disorder from major depressive disorder in adolescents. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2022, 272, 817-826.	3.2	14
2	Difference in the regulation of biological rhythm symptoms of Major depressive disorder between escitalopram and mirtazapine. <i>Journal of Affective Disorders</i> , 2022, 296, 258-264.	4.1	3
3	Impaired robust interhemispheric function integration of depressive brain from REST meta-MDD database in China. <i>Bipolar Disorders</i> , 2022, 24, 400-411.	1.9	8
4	Evaluating the efficacy and moderators of algorithm-guided antidepressant treatments of major depressive disorder. <i>Journal of Affective Disorders</i> , 2022, 297, 68-75.	4.1	1
5	A Breakthrough in Understanding the Rapid Antidepressant Effect of Ketamine Based on Structural Analysis. <i>Neuroscience Bulletin</i> , 2022, 38, 229-231.	2.9	2
6	A Preliminary Study of Different Treatment Strategies for Anxious Depression. <i>Neuropsychiatric Disease and Treatment</i> , 2022, Volume 18, 11-18.	2.2	0
7	Bilateral Habenula deep brain stimulation for treatment-resistant depression: clinical findings and electrophysiological features. <i>Translational Psychiatry</i> , 2022, 12, 52.	4.8	21
8	Prevalence, clinical features and prescription patterns of psychotropic medications for patients with psychotic depression in China. <i>Journal of Affective Disorders</i> , 2022, 301, 248-252.	4.1	3
9	The mechanism underlying extrapulmonary complications of the coronavirus disease 2019 and its therapeutic implication. <i>Signal Transduction and Targeted Therapy</i> , 2022, 7, 57.	17.1	34
10	Phenotypes, mechanisms and therapeutics: insights from bipolar disorder GWAS findings. <i>Molecular Psychiatry</i> , 2022, 27, 2927-2939.	7.9	17
11	Analysis of Seasonal Clinical Characteristics in Patients With Bipolar or Unipolar Depression. <i>Frontiers in Psychiatry</i> , 2022, 13, 847485.	2.6	1
12	Employing biochemical biomarkers for building decision tree models to predict bipolar disorder from major depressive disorder. <i>Journal of Affective Disorders</i> , 2022, 308, 190-198.	4.1	10
13	Early Diagnosis of Bipolar Disorder Coming Soon: Application of an Oxidative Stress Injury Biomarker (BIOS) Model. <i>Neuroscience Bulletin</i> , 2022, 38, 979-991.	2.9	8
14	Exploring the Core Genes of Schizophrenia Based on Bioinformatics Analysis. <i>Genes</i> , 2022, 13, 967.	2.4	2
15	Short- and Long-Term Influences of Benzodiazepine and Z-Drug Use in Patients with Bipolar Disorder Combined Sleep Disturbance during Affective Period: A Nine-Month Follow-Up Analysis. <i>Disease Markers</i> , 2022, 2022, 1-7.	1.3	1
16	Reduced nucleus accumbens functional connectivity in reward network and default mode network in patients with recurrent major depressive disorder. <i>Translational Psychiatry</i> , 2022, 12, .	4.8	20
17	Novel Risk Loci Associated With Genetic Risk for Bipolar Disorder Among Han Chinese Individuals. <i>JAMA Psychiatry</i> , 2021, 78, 320.	11.0	35
18	Independent replications and integrative analyses confirm TRANK1 as a susceptibility gene for bipolar disorder. <i>Neuropsychopharmacology</i> , 2021, 46, 1103-1112.	5.4	20

#	ARTICLE	IF	CITATIONS
19	Probing the clinical and brain structural boundaries of bipolar and major depressive disorder. <i>Translational Psychiatry</i> , 2021, 11, 48.	4.8	9
20	Major Depressive Disorder: Advances in Neuroscience Research and Translational Applications. <i>Neuroscience Bulletin</i> , 2021, 37, 863-880.	2.9	107
21	Disturbances of affective cognition in mood disorders. <i>Science China Life Sciences</i> , 2021, 64, 938-941.	4.9	3
22	PTEN in prefrontal cortex is essential in regulating depression-like behaviors in mice. <i>Translational Psychiatry</i> , 2021, 11, 185.	4.8	21
23	Lower Health Literacy of Mania Than Depression Among Older People: A Random Survey of a Community Healthcare Service Center. <i>Frontiers in Psychiatry</i> , 2021, 12, 512689.	2.6	2
24	Disrupted hemispheric connectivity specialization in patients with major depressive disorder: Evidence from the REST-meta-MDD Project. <i>Journal of Affective Disorders</i> , 2021, 284, 217-228.	4.1	23
25	PAID study design on the role of PKC activation in immune/inflammation-related depression: a randomised placebo-controlled trial protocol. <i>Annals of General Psychiatry</i> , 2021, 34, e100440.	3.1	3
26	Hypothalamic-Pituitary-End-Organ Axes: Hormone Function in Female Patients with Major Depressive Disorder. <i>Neuroscience Bulletin</i> , 2021, 37, 1176-1187.	2.9	14
27	Can seizure therapies and noninvasive brain stimulations prevent suicidality? A systematic review. <i>Brain and Behavior</i> , 2021, 11, e02144.	2.2	8
28	Cognitive control and emotional response in attention-deficit/ hyperactivity disorder comorbidity with disruptive, impulse-control, and conduct disorders. <i>BMC Psychiatry</i> , 2021, 21, 232.	2.6	6
29	Gene expression signatures differentiating major depressive disorder from subsyndromal symptomatic depression. <i>Aging</i> , 2021, 13, 13124-13137.	3.1	2
30	Gastrointestinal Symptoms During Depressive Episodes in 3256 Patients with Major Depressive Disorders: Findings from the NSSD. <i>Journal of Affective Disorders</i> , 2021, 286, 27-32.	4.1	15
31	Clinical features of the patients with major depressive disorder co-occurring insomnia and hypersomnia symptoms: a report of NSSD study. <i>Sleep Medicine</i> , 2021, 81, 375-381.	1.6	13
32	Predictors and moderators of quality of life in patients with major depressive disorder: An AGTs-MDD study report. <i>Journal of Psychiatric Research</i> , 2021, 138, 96-102.	3.1	5
33	Schizophrenia, bipolar disorder, or intracranial aneurysm? A case report. <i>Brain and Behavior</i> , 2021, 11, e2245.	2.2	2
34	A Preliminary Randomized Controlled Trial of Different Treatment Regimens for Melancholic Depression. <i>Neuropsychiatric Disease and Treatment</i> , 2021, Volume 17, 2441-2449.	2.2	0
35	Disrupted intrinsic functional brain topology in patients with major depressive disorder. <i>Molecular Psychiatry</i> , 2021, 26, 7363-7371.	7.9	82
36	Barriers and facilitators to implementing measurement-based care for depression in Shanghai, China: a situational analysis. <i>BMC Psychiatry</i> , 2021, 21, 430.	2.6	4

#	ARTICLE	IF	CITATIONS
55	Functional Status of Hypothalamicâ€“Pituitaryâ€“Thyroid and Hypothalamicâ€“Pituitaryâ€“Adrenal Axes in Hospitalized Schizophrenics in Shanghai. <i>Frontiers in Psychiatry</i> , 2020, 11, 65.	2.6	19
56	The relationship of olfactory function and clinical traits in major depressive disorder. <i>Behavioural Brain Research</i> , 2020, 386, 112594.	2.2	10
57	Identification of a functional human-unique 351-bp Alu insertion polymorphism associated with major depressive disorder in the 1p31.1 GWAS risk loci. <i>Neuropsychopharmacology</i> , 2020, 45, 1196-1206.	5.4	17
58	Ifenprodil rapidly ameliorates depressive-like behaviors, activates mTOR signaling and modulates proinflammatory cytokines in the hippocampus of CUMS rats. <i>Psychopharmacology</i> , 2020, 237, 1421-1433.	3.1	22
59	Potential contribution of increased soluble IL-2R to lymphopenia in COVID-19 patients. <i>Cellular and Molecular Immunology</i> , 2020, 17, 878-880.	10.5	45
60	The Risk and Prevention of Novel Coronavirus Pneumonia Infections Among Inpatients in Psychiatric Hospitals. <i>Neuroscience Bulletin</i> , 2020, 36, 299-302.	2.9	178
61	<p>Variants in the Upstream Region of the Insulin Receptor Substrate-1 Gene Is Associated with Major Depressive Disorder in the Han Chinese Population</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 501-507.	2.2	1
62	Association between residual symptoms and social functioning in patients with depression. <i>Comprehensive Psychiatry</i> , 2020, 98, 152164.	3.1	10
63	Role of biological rhythm dysfunction in the development and management of bipolar disorders: a review. <i>Annals of General Psychiatry</i> , 2020, 33, e100127.	3.1	9
64	Clinical and immunological features of severe and moderate coronavirus disease 2019. <i>Journal of Clinical Investigation</i> , 2020, 130, 2620-2629.	8.2	3,820
65	Cortical thickness and subcortical volumes alterations in euthymic bipolar I patients treated with different mood stabilizers. <i>Brain Imaging and Behavior</i> , 2019, 13, 1255-1264.	2.1	8
66	Clinical characteristics associated with therapeutic nonadherence of the patients with major depressive disorder: A report on the National Survey on Symptomatology of Depression in China. <i>CNS Neuroscience and Therapeutics</i> , 2019, 25, 215-222.	3.9	8
67	<p>Cognitive symptoms in major depressive disorder: associations with clinical and functional outcomes in a 6-month, non-interventional, prospective study in China</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2019, Volume 15, 1723-1736.	2.2	25
68	Perspective on Etiology and Treatment of Bipolar Disorders in China: Clinical Implications and Future Directions. <i>Neuroscience Bulletin</i> , 2019, 35, 608-612.	2.9	5
69	Preliminary Clinical Investigation of Combinatorial Pharmacogenomic Testing for the Optimized Treatment of Depression: A Randomized Single-Blind Study. <i>Frontiers in Neuroscience</i> , 2019, 13, 960.	2.8	15
70	The Relationship Between Neuroimmunity and Bipolar Disorder: Mechanism and Translational Application. <i>Neuroscience Bulletin</i> , 2019, 35, 595-607.	2.9	19
71	Detection Study of Bipolar Depression Through the Application of a Model-Based Algorithm in Terms of Clinical Feature and Peripheral Biomarkers. <i>Frontiers in Psychiatry</i> , 2019, 10, 266.	2.6	7
72	Editorial: Involvement of Neuro-Immune Mechanism and Brainâ€“Gut Axis in Pathophysiology of Mood Disorders. <i>Frontiers in Psychiatry</i> , 2019, 10, 403.	2.6	2

#	ARTICLE	IF	CITATIONS
73	Perception of Stigma and Its Associated Factors Among Patients With Major Depressive Disorder: A Multicenter Survey From an Asian Population. <i>Frontiers in Psychiatry</i> , 2019, 10, 321.	2.6	26
74	Disagreement and factors between symptom on self-report and clinician rating of major depressive disorder: A report of a national survey in China. <i>Journal of Affective Disorders</i> , 2019, 253, 141-146.	4.1	10
75	Reduced default mode network functional connectivity in patients with recurrent major depressive disorder. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 9078-9083.	7.1	441
76	Integrative analyses of major histocompatibility complex loci in the genome-wide association studies of major depressive disorder. <i>Neuropsychopharmacology</i> , 2019, 44, 1552-1561.	5.4	27
77	Subtypes of treatment-resistant depression determined by a latent class analysis in a Chinese clinical population. <i>Journal of Affective Disorders</i> , 2019, 249, 82-89.	4.1	12
78	The depression GWAS risk allele predicts smaller cerebellar gray matter volume and reduced SIRT1 mRNA expression in Chinese population. <i>Translational Psychiatry</i> , 2019, 9, 333.	4.8	25
79	Prevalence and clinical features of atypical depression among patients with major depressive disorder in China. <i>Journal of Affective Disorders</i> , 2019, 246, 285-289.	4.1	7
80	Complement C7 is a novel risk gene for Alzheimer's disease in Han Chinese. <i>National Science Review</i> , 2019, 6, 257-274.	9.5	55
81	The association between somatic symptoms and suicidal ideation in Chinese first-episode major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 245, 17-21.	4.1	30
82	Introduction. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1180, 1-17.	1.6	3
83	Advance in Diagnosis of Depressive Disorder. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1180, 179-191.	1.6	4
84	Response Inhibition and Emotional Regulation in the Patients with Attention-Deficit/Hyperactivity Disorder and Comorbidity of Disruptive, Impulse-Control, and Conduct Disorders. <i>Psychiatry Investigation</i> , 2019, 16, 872-874.	1.6	4
85	Prevalence, risk factors and clinical characteristics of suicidal ideation in Chinese patients with depression. <i>Journal of Affective Disorders</i> , 2018, 235, 135-141.	4.1	40
86	Assessment and management of bipolar disorder: Principal summary of updated Chinese guidelines. <i>Bipolar Disorders</i> , 2018, 20, 289-292.	1.9	7
87	HTR1A/1B DNA methylation may predict escitalopram treatment response in depressed Chinese Han patients. <i>Journal of Affective Disorders</i> , 2018, 228, 222-228.	4.1	38
88	The Arc Gene Confers Genetic Susceptibility to Alzheimer's Disease in Han Chinese. <i>Molecular Neurobiology</i> , 2018, 55, 1217-1226.	4.0	30
89	Effects of tumor necrosis factor- β polymorphism on the brain structural changes of the patients with major depressive disorder. <i>Translational Psychiatry</i> , 2018, 8, 217.	4.8	25
90	Blood-based dynamic genomic signature for obsessive-compulsive disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2018, 177, 709-716.	1.7	5

#	ARTICLE	IF	CITATIONS
91	Clinical outcomes of patients with major depressive disorder treated with either duloxetine, escitalopram, fluoxetine, paroxetine, or sertraline. <i>Neuropsychiatric Disease and Treatment</i> , 2018, Volume 14, 2473-2484.	2.2	7
92	Somatic symptoms vary in major depressive disorder in China. <i>Comprehensive Psychiatry</i> , 2018, 87, 32-37.	3.1	37
93	The clinical correlates of comorbid anxiety symptoms and syndromal anxiety in patients with major depressive disorder. <i>Psychiatry Research</i> , 2018, 269, 251-257.	3.3	18
94	Association of DNA methylation in BDNF with escitalopram treatment response in depressed Chinese Han patients. <i>European Journal of Clinical Pharmacology</i> , 2018, 74, 1011-1020.	1.9	42
95	Risk Factors for Recent Suicide Attempts in Major Depressive Disorder Patients in China: Results From a National Study. <i>Frontiers in Psychiatry</i> , 2018, 9, 300.	2.6	18
96	Genetic association of the cytochrome c oxidase-related genes with Alzheimer's disease in Han Chinese. <i>Neuropsychopharmacology</i> , 2018, 43, 2264-2276.	5.4	29
97	Different levels of pro- and anti-inflammatory cytokines in patients with unipolar and bipolar depression. <i>Journal of Affective Disorders</i> , 2018, 237, 65-72.	4.1	47
98	Abnormal white matter integrity in Chinese young adults with first-episode medication-free anxious depression: a possible neurological biomarker of subtype major depressive disorder. <i>Neuropsychiatric Disease and Treatment</i> , 2018, Volume 14, 2017-2026.	2.2	13
99	The association of duration and severity of disease with executive function: Differences between drug-naïve patients with bipolar and unipolar depression. <i>Journal of Affective Disorders</i> , 2018, 238, 412-417.	4.1	12
100	Analysis of Misdiagnosis of Bipolar Disorder in An Outpatient Setting. <i>Shanghai Archives of Psychiatry</i> , 2018, 30, 93-101.	0.7	33
101	Female-specific effect of the BDNF gene on Alzheimer's disease. <i>Neurobiology of Aging</i> , 2017, 53, 192.e11-192.e19.	3.1	46
102	Increased ratio of high sensitivity C-reactive protein to interleukin-10 as a potential peripheral biomarker of schizophrenia and aggression. <i>International Journal of Psychophysiology</i> , 2017, 114, 9-15.	1.0	24
103	Gender differences in quality of life and functional disability for depression outpatients with or without residual symptoms after acute phase treatment in China. <i>Journal of Affective Disorders</i> , 2017, 219, 141-148.	4.1	13
104	Efficacy and safety of escitalopram in treatment of severe depression in Chinese population. <i>Metabolic Brain Disease</i> , 2017, 32, 891-901.	2.9	14
105	Reduced ENA78 levels as novel biomarker for major depressive disorder and venlafaxine efficiency: Result from a prospective longitudinal study. <i>Psychoneuroendocrinology</i> , 2017, 81, 113-121.	2.7	21
106	Duration of untreated bipolar disorder: a multicenter study. <i>Scientific Reports</i> , 2017, 7, 44811.	3.3	21
107	Identification of plasma biomarkers for distinguishing bipolar depression from major depressive disorder by iTRAQ-coupled LC-MS/MS and bioinformatics analysis. <i>Psychoneuroendocrinology</i> , 2017, 86, 17-24.	2.7	51
108	Common variants at 2q11.2, 8q21.3, and 11q13.2 are associated with major mood disorders. <i>Translational Psychiatry</i> , 2017, 7, 1273.	4.8	9

#	ARTICLE	IF	CITATIONS
109	Association analysis between mitogen-activated protein/extracellular signal-regulated kinase (MEK) gene polymorphisms and depressive disorder in the Han Chinese population. <i>Journal of Affective Disorders</i> , 2017, 222, 120-125.	4.1	6
110	Prescribing patterns of psychotropic medications and clinical features in patients with major depressive disorder with and without comorbid dysthymia in China. <i>Asia-Pacific Psychiatry</i> , 2017, 9, e12261.	2.2	1
111	Rare Genetic Variants of the Transthyretin Gene Are Associated with Alzheimer's Disease in Han Chinese. <i>Molecular Neurobiology</i> , 2017, 54, 5192-5200.	4.0	24
112	Ratio of mBDNF to proBDNF for Differential Diagnosis of Major Depressive Disorder and Bipolar Depression. <i>Molecular Neurobiology</i> , 2017, 54, 5573-5582.	4.0	62
113	Differential gene expression in patients with subsyndromal symptomatic depression and major depressive disorder. <i>PLoS ONE</i> , 2017, 12, e0172692.	2.5	10
114	Fatty acid amide hydrolase inhibitors produce rapid anti-anxiety responses through amygdala long-term depression in male rodents. <i>Journal of Psychiatry and Neuroscience</i> , 2017, 42, 230-241.	2.4	27
115	Complement factor H and susceptibility to major depressive disorder in Han Chinese. <i>British Journal of Psychiatry</i> , 2016, 208, 446-452.	2.8	21
116	Evaluations of treatment efficacy of depression from perspective of both patients' symptoms and general sense of mental health and wellbeing: A large scale, multi-centered, longitudinal study in China. <i>Psychiatry Research</i> , 2016, 241, 55-60.	3.3	4
117	Alterations of microRNA-124 expression in peripheral blood mononuclear cells in pre- and post-treatment patients with major depressive disorder. <i>Journal of Psychiatric Research</i> , 2016, 78, 65-71.	3.1	74
118	Identification of SLC25A37 as a major depressive disorder risk gene. <i>Journal of Psychiatric Research</i> , 2016, 83, 168-175.	3.1	24
119	Evaluating the association between the SHANK3 gene and bipolar disorder. <i>Psychiatry Research</i> , 2016, 244, 284-288.	3.3	10
120	Identification of IL6 as a susceptibility gene for major depressive disorder. <i>Scientific Reports</i> , 2016, 6, 31264.	3.3	35
121	Higher Plasma S100B Concentrations in Schizophrenia Patients and Dependently Associated with Inflammatory Markers. <i>Scientific Reports</i> , 2016, 6, 27584.	3.3	32
122	Demographic and clinical differences between early- and late-onset bipolar disorders in a multicenter study in China. <i>Psychiatry Research</i> , 2016, 246, 688-691.	3.3	8
123	Plasma levels of Th17-related cytokines and complement C3 correlated with aggressive behavior in patients with schizophrenia. <i>Psychiatry Research</i> , 2016, 246, 700-706.	3.3	59
124	Validating GWAS-Identified Risk Loci for Alzheimer's Disease in Han Chinese Populations. <i>Molecular Neurobiology</i> , 2016, 53, 379-390.	4.0	62
125	CFH Variants Affect Structural and Functional Brain Changes and Genetic Risk of Alzheimer's Disease. <i>Neuropsychopharmacology</i> , 2016, 41, 1034-1045.	5.4	58
126	Neprilysin Confers Genetic Susceptibility to Alzheimer's Disease in Han Chinese. <i>Molecular Neurobiology</i> , 2016, 53, 4883-4892.	4.0	21

#	ARTICLE	IF	CITATIONS
127	PLD3 in Alzheimer's Disease: a Modest Effect as Revealed by Updated Association and Expression Analyses. <i>Molecular Neurobiology</i> , 2016, 53, 4034-4045.	4.0	30
128	ZNF804A Genetic Variation Confers Risk to Bipolar Disorder. <i>Molecular Neurobiology</i> , 2016, 53, 2936-2943.	4.0	21
129	Important clinical features of atypical antipsychotics in acute bipolar depression that inform routine clinical care: a review of pivotal studies with number needed to treat. <i>Neuroscience Bulletin</i> , 2015, 31, 572-588.	2.9	17
130	Authors' reply. <i>British Journal of Psychiatry</i> , 2015, 206, 79-80.	2.8	0
131	IL-23 and TGF- β 1 levels as potential predictive biomarkers in treatment of bipolar I disorder with acute manic episode. <i>Journal of Affective Disorders</i> , 2015, 174, 361-366.	4.1	50
132	Atypical features and treatment choices in bipolar disorders: a result of the National Bipolar Mania Pathway Survey in China. <i>Neuroscience Bulletin</i> , 2015, 31, 22-30.	2.9	4
133	Mitochondrial DNA haplogroup B5 confers genetic susceptibility to Alzheimer's disease in Han Chinese. <i>Neurobiology of Aging</i> , 2015, 36, 1604.e7-1604.e16.	3.1	50
134	Down-regulation of PRKCB1 expression in Han Chinese patients with subsyndromal symptomatic depression. <i>Journal of Psychiatric Research</i> , 2015, 69, 1-6.	3.1	6
135	Guidelines concordance of maintenance treatment in euthymic patients with bipolar disorder: Data from the national bipolar mania pathway survey (BIPAS) in mainland China. <i>Journal of Affective Disorders</i> , 2015, 182, 101-105.	4.1	6
136	Dissociated large-scale functional connectivity networks of the precuneus in medication-naïve first-episode depression. <i>Psychiatry Research - Neuroimaging</i> , 2015, 232, 250-256.	1.8	65
137	Demographic and clinical differences between early- and late-onset major depressions in thirteen psychiatric institutions in China. <i>Journal of Affective Disorders</i> , 2015, 170, 266-269.	4.1	8
138	Validation of the Chinese Version of the Short TEMPS-A and its application in patients with mood disorders. <i>Journal of Affective Disorders</i> , 2015, 170, 178-184.	4.1	9
139	Surface Vulnerability of Cerebral Cortex to Major Depressive Disorder. <i>PLoS ONE</i> , 2015, 10, e0120704.	2.5	62
140	Risk Factors for Anxiety in Major Depressive Disorder Patients. <i>Clinical Psychopharmacology and Neuroscience</i> , 2015, 13, 263-268.	2.0	24
141	Evaluation of Mood Disorder Questionnaire (MDQ) in Patients with Mood Disorders: A Multicenter Trial across China. <i>PLoS ONE</i> , 2014, 9, e91895.	2.5	14
142	Nerve growth factor variations in patients with mood disorders: no changes in eight weeks of clinical treatment. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 835.	2.2	11
143	Alterations in effective connectivity anchored on the insula in major depressive disorder. <i>European Neuropsychopharmacology</i> , 2014, 24, 1784-1792.	0.7	58
144	Identification of ANKK1 rs1800497 variant in schizophrenia: New data and meta-analysis. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2014, 165, 564-571.	1.7	15

#	ARTICLE	IF	CITATIONS
145	The efficacy of plasma biomarkers in early diagnosis of Alzheimer's disease. <i>International Journal of Geriatric Psychiatry</i> , 2014, 29, 713-719.	2.7	49
146	Association between brain-derived neurotrophic factor genetic polymorphism Val66Met and susceptibility to bipolar disorder: a meta-analysis. <i>BMC Psychiatry</i> , 2014, 14, 366.	2.6	20
147	Suicide risk in major affective disorder: Results from a national survey in China. <i>Journal of Affective Disorders</i> , 2014, 155, 174-179.	4.1	27
148	Genetic modulation of working memory deficits by ankyrin 3 gene in schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2014, 50, 110-115.	4.8	33
149	No association between genetic variants of the LRRK2 gene and schizophrenia in Han Chinese. <i>Neuroscience Letters</i> , 2014, 566, 210-215.	2.1	6
150	Metabolic syndrome in patients taking clozapine: prevalence and influence of catechol-O-methyltransferase genotype. <i>Psychopharmacology</i> , 2014, 231, 2211-2218.	3.1	28
151	A study of N-methyl-D-aspartate receptor gene (GRIN2B) variants as predictors of treatment-resistant major depression. <i>Psychopharmacology</i> , 2014, 231, 685-693.	3.1	65
152	Glutamate receptor 1 phosphorylation at serine 845 contributes to the therapeutic effect of olanzapine on schizophrenia-like cognitive impairments. <i>Schizophrenia Research</i> , 2014, 159, 376-384.	2.0	16
153	Brain-derived neurotrophic factor levels and bipolar disorder in patients in their first depressive episode: 3-year prospective longitudinal study. <i>British Journal of Psychiatry</i> , 2014, 205, 29-35.	2.8	54
154	Altered brain network modules induce helplessness in major depressive disorder. <i>Journal of Affective Disorders</i> , 2014, 168, 21-29.	4.1	57
155	Decreased serum fibroblast growth factor - 2 levels in pre- and post-treatment patients with major depressive disorder. <i>Neuroscience Letters</i> , 2014, 579, 168-172.	2.1	39
156	Influence of BCL2 gene in major depression susceptibility and antidepressant treatment outcome. <i>Journal of Affective Disorders</i> , 2014, 155, 288-294.	4.1	27
157	Elevated serum levels of FGF-2, NGF and IGF-1 in patients with manic episode of bipolar disorder. <i>Psychiatry Research</i> , 2014, 218, 54-60.	3.3	58
158	MiRNA-206 and BDNF genes interacted in bipolar I disorder. <i>Journal of Affective Disorders</i> , 2014, 162, 116-119.	4.1	35
159	Authors' reply. <i>British Journal of Psychiatry</i> , 2014, 205, 410-411.	2.8	1
160	Guidelines Discordance in Acute Bipolar Depression: Data from the National Bipolar Mania Pathway Survey (BIPAS) in Mainland China. <i>PLoS ONE</i> , 2014, 9, e96096.	2.5	11
161	Comorbidity of depressive and anxiety disorders: challenges in diagnosis and assessment. <i>Shanghai Archives of Psychiatry</i> , 2014, 26, 227-31.	0.7	53
162	Evaluation of antidepressant polypharmacy and other interventions for treatment-resistant depression. <i>Shanghai Archives of Psychiatry</i> , 2014, 26, 365-7.	0.7	0

#	ARTICLE	IF	CITATIONS
163	Neurotrophic Tyrosine Kinase Receptor Type 2 (NTRK2) Gene Associated with Treatment Response to Mood Stabilizers in Patients with Bipolar I Disorder. <i>Journal of Molecular Neuroscience</i> , 2013, 50, 305-310.	2.3	40
164	Difference in remission in a Chinese population with anxious versus nonanxious treatment-resistant depression: A report of OPERATION study. <i>Journal of Affective Disorders</i> , 2013, 150, 834-839.	4.1	58
165	The role of BDNF, NTRK2 gene and their interaction in development of treatment-resistant depression: Data from multicenter, prospective, longitudinal clinic practice. <i>Journal of Psychiatric Research</i> , 2013, 47, 8-14.	3.1	56
166	Venlafaxine inhibits the upregulation of plasma tumor necrosis factor-alpha (TNF- α) in the Chinese patients with major depressive disorder: A prospective longitudinal study. <i>Psychoneuroendocrinology</i> , 2013, 38, 107-114.	2.7	58
167	Sociodemographic and clinical features of bipolar disorder patients misdiagnosed with major depressive disorder in China. <i>Bipolar Disorders</i> , 2013, 15, 199-205.	1.9	41
168	Are subsyndromal symptomatic depression and major depressive disorder distinct disorders?. <i>Shanghai Archives of Psychiatry</i> , 2012, 24, 286-7.	0.7	3
169	A Pilot Study of the Efficacy and Safety of Paroxetine Augmented With Risperidone, Valproate, Buspirone, Trazodone, or Thyroid Hormone in Adult Chinese Patients With Treatment-Resistant Major Depression. <i>Journal of Clinical Psychopharmacology</i> , 2011, 31, 638-642.	1.4	47
170	Comparisons of the Efficacy and Tolerability of Extended-Release Venlafaxine, Mirtazapine, and Paroxetine in Treatment-Resistant Depression. <i>Journal of Clinical Psychopharmacology</i> , 2010, 30, 357-364.	1.4	48
171	Biochemical and Endocrine Parameters for the Discrimination and Calibration of Bipolar Disorder or Major Depressive Disorder. <i>Frontiers in Psychiatry</i> , 0, 13, .	2.6	3