

Satoko Hayashi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6589309/publications.pdf>

Version: 2024-02-01

98
papers

2,232
citations

201674

27
h-index

254184

43
g-index

102
all docs

102
docs citations

102
times ranked

893
citing authors

#	ARTICLE	IF	CITATIONS
1	Intrinsic Dynamic and Static Nature of π - π Interactions in Fused Benzene-Type Helicenes and Dimers, Elucidated with QTAIM Dual Functional Analysis. <i>Nanomaterials</i> , 2022, 12, 321.	4.1	4
2	Synthesis and Characterization of Monomeric Hexacoordinated Chalcogenonium Salts Bearing 2-(2-Pyridyl)phenyl Ligands. <i>Bulletin of the Chemical Society of Japan</i> , 2021, 94, 1192-1200.	3.2	0
3	Intrinsic Dynamic and Static Nature of Halogen Bonding in Neutral Polybromine Clusters, with the Structural Feature Elucidated by QTAIM Dual-Functional Analysis and MO Calculations. <i>Molecules</i> , 2021, 26, 2936.	3.8	1
4	Linear Multiselenium Interactions in Dicationic Oligomers of 1,5-(Diseleno)canes: Behavior of Se _{mc} \int (m) Tj ETQq 0 0 rgBT /Overlock	1.9	0
5	Linear Multiselenium Interactions in Dicationic Oligomers of 1,5-(Diseleno)canes: Behavior of Se _{mc} \int (m) Tj ETQq 1 1 0.784314 rgBT	1.9	0
6	Dynamic and static nature of activated interactions in transition states as elucidated by quantum theory of atoms-in-molecules dual functional analysis: A case of ligand exchange at the N of sulfonylimino-3-bromanes. <i>International Journal of Quantum Chemistry</i> , 2020, 120, e26073.	2.0	2
7	Dynamic and Static Nature of Br ₄ and Se ₂ Br ₅ in the Selenanthrene System and Related Species Elucidated by QTAIM Dual Functional Analysis with QC Calculations. <i>Bioinorganic Chemistry and Applications</i> , 2020, 2020, 1-14.	4.1	1
8	Intrinsic dynamic and static nature of each HB in the multi-HBs between nucleobase pairs and its behavior, elucidated with QTAIM dual functional analysis and QC calculations. <i>RSC Advances</i> , 2020, 10, 24730-24742.	3.6	2
9	Nature of the E π -E π^2 interactions (E, E π^2 = O, S, Se, and Te) at naphthalene 1,8-positions with fine details of the structures: experimental and theoretical investigations. <i>New Journal of Chemistry</i> , 2019, 43, 14224-14237.	2.8	9
10	Effects from Basis Sets and Levels of Calculations on the Nature of Interactions Predicted by QTAIM Dual Functional Analysis with QTAIM Functions. <i>ChemistrySelect</i> , 2019, 4, 6198-6208.	1.5	5
11	Nature of intramolecular O π -H π interactions as elucidated by QTAIM dual functional analysis with QC calculations. <i>RSC Advances</i> , 2019, 9, 15521-15530.	3.6	1
12	The nature of G π -E π (3 π -4e) in Me _n GCH ₂ C ₆ H ₄ EY (Me _n G =) Tj ETQq 0 0,0 rgBT /Oylock 10 and compliance constants in noncovalent G π -E interactions. <i>RSC Advances</i> , 2019, 9, 39435-39446.	3.6	0
13	Behavior of Multi-HBs in Acetic Acid Dimer and Related Species: QTAIM Dual Functional Analysis Employing Perturbed Structures Generated Using Coordinates from Compliance Force Constants. <i>Bulletin of the Chemical Society of Japan</i> , 2019, 92, 87-96.	3.2	4
14	High-resolution X-ray diffraction determination of the electron density of 1-(8-PhSC ₁₀ H ₆)SS(C ₁₀ H ₆)SPh-8 π^2 -1 π^2 with the QTAIM approach: evidence for S ₄ \int (4 π -6e) at the naphthalene π -peri-positions. <i>RSC Advances</i> , 2018, 8, 9651-9660.	3.6	12
15	Perturbed structures generated using coordinates derived from compliance constants in internal vibrations for QTAIM dual functional analysis: Intrinsic dynamic nature of interactions. <i>International Journal of Quantum Chemistry</i> , 2018, 118, e25590.	2.0	17
16	Behaviour of the XH π and YX π interactions (X, Y = F, Cl, Br and I) in the coronene π -system, as elucidated by QTAIM dual functional analysis with QC calculations. <i>RSC Advances</i> , 2018, 8, 16349-16361.	3.6	3
17	Behavior of I ₄ \int (4 π -6e) in tellurolane system and related species, elucidated by QTAIM dual functional analysis with QC calculations. <i>Heteroatom Chemistry</i> , 2018, 29, .	0.7	2
18	Intrinsic Dynamic Nature of Neutral Hydrogen Bonds Elucidated with QTAIM Dual Functional Analysis: Role of the Compliance Force Constants and QTAIM DFA Parameters in Stability. <i>ChemistryOpen</i> , 2018, 7, 565-575.	1.9	5

#	ARTICLE	IF	CITATIONS
19	Behavior of the Eâ€™Eâ€™™ Bonds (E, Eâ€™™ = S and Se) in Glutathione Disulfide and Derivatives Elucidated by Quantum Chemical Calculations with the Quantum Theory of Atoms-in-Molecules Approach. <i>Molecules</i> , 2018, 23, 443.	3.8	5
20	Dynamic and Static Behavior of Intramolecular ĩ€-ĩ€ Interactions in [2.2]- and [3.3]Cyclophanes, Elucidated by QTAIM Dual Functional Analysis with QC Calculations. <i>ChemistrySelect</i> , 2017, 2, 1774-1782.	1.5	2
21	Behavior of interactions between hydrogen chalcogenides and an anthracene ĩ€-system elucidated by QTAIM dual functional analysis with QC calculations. <i>RSC Advances</i> , 2017, 7, 31858-31865.	3.6	4
22	Behavior of Intramolecular ĩ€â€™ĩ€ Interactions with Doubly Degenerated Bond Paths Between Carbon Atoms in Opposite Benzene Rings of Diethenodihydronaphthalenes by QTAIM Approach. <i>ChemistrySelect</i> , 2017, 2, 90-100.	1.5	4
23	Linear Four-Chalcogen Interactions in Radical Cationic and Dicationic Dimers of 1,5-(Dichalcogena)canes: Nature of the Interactions Elucidated by QTAIM Dual Functional Analysis with QC Calculations. <i>Journal of Physical Chemistry A</i> , 2017, 121, 2482-2496.	2.5	5
24	Relativistic Effect on 1 J (M,C) in Me ₄ M, Me ₃ Mâ™™, Ph ₄ M, and Ph ₃ Mâ™™ (M=Pb, Sn, Ge, Si, and/or C): Role of s-Type Lone Pair Orbitals in the Distinct Effect for the Anionic Species. <i>ChemPhysChem</i> , 2017, 18, 2466-2474.	2.1	2
25	Dichlorosilylene Transfer to aP-Fluorophosphaalkene: The Route to aC-Dichlorofluorosilyl-Functionalized Dialkyldiphosphene. <i>European Journal of Inorganic Chemistry</i> , 2017, 2017, 1526-1536.	2.0	4
26	Nature of E2X2ĭf(4câ€™6e) of the X---Eâ€™”E---X type at naphthalene 1,8-positions and model, elucidated by X-ray crystallographic analysis and QC calculations with the QTAIM approach. <i>Acta Crystallographica Section B: Structural Science, Crystal Engineering and Materials</i> , 2017, 73, 265-275.	1.1	4
27	Behavior of Halogen Bonds of the Yâ™”Xâ€™...â€™...ĩ€ Type (X, Y=F, Cl, Br, I) in the Benzene ĩ€ System, Elucidated by Using a Quantum Theory of Atoms in Molecules Dualâ€™Functional Analysis. <i>ChemPhysChem</i> , 2016, 17, 2579-2589.	2.1	12
28	Nature of S₂Se₂ ĭf(4câ€™6e) at naphthalene 1,8-positions and models, elucidated by QTAIM dual functional analysis. <i>RSC Advances</i> , 2016, 6, 93195-93204.	3.6	12
29	Mechanistic Studies on the Generation and Properties of Superelectrophilic Singlet Carbenes from Bis(perfluoroalkanesulfonyl)bromonium Ylides. <i>Journal of Organic Chemistry</i> , 2016, 81, 3188-3198.	3.2	16
30	Quantum chemical calculations with the AIM approach applied to the ĩ€-interactions between hydrogen chalcogenides and naphthalene. <i>RSC Advances</i> , 2016, 6, 49651-49660.	3.6	7
31	Intramolecular ĩ€â€™ĩ€ Interactions in Diethanodihydronaphthalene and Derivatives: Dynamic and Static Behavior of the Interactions Elucidated by QTAIM Dual Functional Analysis. <i>ChemistrySelect</i> , 2016, 1, 2344-2353.	1.5	3
32	Dynamic and static behavior of the Hâ€™ĩ€ and Eâ€™ĩ€ interactions in EH₂ adducts of benzene ĩ€-system (E = O, S, Se and Te), elucidated by QTAIM dual functional analysis. <i>Physical Chemistry Chemical Physics</i> , 2016, 18, 9948-9960.	2.8	14
33	Transannular EÂˆ-Âˆ-Eâ€™² Interactions in Neutral, Radical Cationic, and Dicationic Forms of <i>cyclo</i>-[E(CH₂CH₂CH₂)₂Eâ€™²] (E, Eâ€™² = S, Se, Te, and Q)	3.2	12
34	Dynamic and static behavior of the Eâ€™Eâ€™² bonds (E, Eâ€™² = S and Se) in cystine and derivatives, elucidated by AIM dual functional analysis. <i>RSC Advances</i> , 2015, 5, 11534-11540.	3.6	4
35	Dynamic and static behavior of hydrogen bonds of the Xâ€™Hâ€™ĩ€ type (X = F, Cl, Br, I, RO and RRâ€™²N; R, Râ€™² = H or Tj ETQq1 1 0.784	2.8	16
36	Photoinduced Regio- and Stereoselective Introduction of Phenylchalcogeno Moieties to Ethynylferrocene. <i>Bulletin of the Chemical Society of Japan</i> , 2014, 87, 550-552.	3.2	4

#	ARTICLE	IF	CITATIONS
37	Relativistic effects on the ^{125}Te and ^{33}S NMR chemical shifts of various tellurium and sulfur species, together with ^{77}Se of selenium congeners, in the framework of a zeroth-order regular approximation: applicability to tellurium compounds. <i>RSC Advances</i> , 2014, 4, 44795-44810.	3.6	9
38	Aromatic Selenoic, Selenothioic, and Diselenoic Acid Salts: Isolation, Characterization, and ^{77}Se NMR Spectra, Together with Theoretical Elucidation. <i>Bulletin of the Chemical Society of Japan</i> , 2014, 87, 677-692.	3.2	7
39	Dynamic and Static Behavior of E \cdots E' Bonds in Neutral and Charged Forms of HEE'H, MeEE'Me, and $\langle i \rangle \text{Cyclo} \langle i \rangle \text{â€}1,2 \text{â€} \text{E}'(\text{CH} \langle \text{sub} \rangle 2 \langle \text{sub} \rangle) \langle \text{sub} \rangle 3 \langle \text{sub} \rangle$ (E, E' = O, S, Se, and Te) Elucidated by AIM Dual Functional Analysis. <i>Heteroatom Chemistry</i> , 2014, 25, 449-472.	0.7	3
40	Role of $d \langle i \rangle G \langle i \rangle d \langle i \rangle w \langle i \rangle$ and $d \langle i \rangle V \langle i \rangle d \langle i \rangle w \langle i \rangle$ in AIM Analysis: An Approach to the Nature of Weak to Strong Interactions. <i>Journal of Physical Chemistry A</i> , 2013, 117, 1795-1803.	2.5	41
41	Dynamic Behavior of Hydrogen Bonds from $\langle i \rangle \text{Pure} \langle i \rangle$ Closed Shell to Shared Shell Interaction Regions Elucidated by AIM Dual Functional Analysis. <i>Journal of Physical Chemistry A</i> , 2013, 117, 1804-1816.	2.5	30
42	CHAPTER 12.3. Hypervalent Chalcogen Compounds. , 2013, , 335-372.		7
43	Experimental and Theoretical Advances in Functional Understanding of Flavonoids as Anti-Tumor Agents. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2013, 13, 307-332.	1.7	14
44	Applications of Normal Coordinates of Internal Vibrations to Generate Perturbed Structures: Dynamic Behavior of Weak to Strong Interactions Elucidated by Atoms-in-Molecules Dual Functional Analysis. <i>Bulletin of the Chemical Society of Japan</i> , 2012, 85, 1293-1305.	3.2	39
45	Unusual Saddle-like Structure of (2-MeOC $_6$ H $_4$ CS) $_2$ S: Theoretical Studies and Comparison with its Oxygen Isologues. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2012, 638, 2508-2520.	1.2	1
46	Role of $p(Z) \text{â€} \text{f}(\text{Ar}/\text{Nap})$ conjugation in structures of 1-(arylchalcogena)naphthalenes for Z = Te versus Se, S and O: experimental and theoretical investigations. <i>Dalton Transactions</i> , 2012, 41, 7485.	3.3	10
47	Relativistic Effect on $\langle \text{sup} \rangle 77 \langle \text{sup} \rangle \text{Se}$ NMR Chemical Shifts of Various Selenium Species in the Framework of Zeroth-Order Regular Approximation. <i>Journal of Physical Chemistry A</i> , 2011, 115, 8721-8730.	2.5	17
48	Dynamic and Static Behaviors of $N \text{â€} Z \text{â€} N \text{ f}(3c \text{â€} 4e)$ (Z = S, Se, and Te) Interactions: Atoms-in-Molecules Dual Functional Analysis with High-Resolution X-ray Diffraction Determination of Electron Densities for 2-(2-Pyridylimino)-2 $\langle i \rangle \text{H} \langle i \rangle$ -1,2,4-thiadiazolo[2,3- $\langle i \rangle a \langle i \rangle$]pyridine. <i>Journal of Physical Chemistry A</i> , 2011, 115, 11775-11787.	2.5	12
49	Atoms-in-Molecules Dual Functional Analysis of Weak to Strong Interactions. <i>Current Organic Chemistry</i> , 2010, 14, 181-197.	1.6	62
50	P(O, S, Se, and Te) $\text{â€} \text{f}(\text{Ar})$ Conjugations as Factors to Control Fine Structures of 1-(Chalcogena)naphthalenes. <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , 2010, 185, 1031-1045.	1.6	4
51	Dynamic Behaviors of Interactions: Application of Normal Coordinates of Internal Vibrations to AIM Dual Functional Analysis. <i>Journal of Physical Chemistry A</i> , 2010, 114, 7423-7430.	2.5	44
52	H/D Isotope Effect on ^{77}Se NMR Chemical Shifts in 8-Methyl-1-(arylselanyl)naphthalenes and Related Selenides: Nonbonded $\text{C} \text{â€} \text{H} \text{â€} \text{Se}$ Through-Space Versus Through-Bond Mechanisms. <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , 2009, 184, 1481-1495.	1.6	2
53	Fine Structures of 8-G-1-($\langle \text{mml} \rangle \text{math} \rangle \text{Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 112 Td} (\text{xmlns:mml}="http://www.w3.org/1998/Math$ Chemistry and Applications. 2009, 2009, 1-11.	4.1	1
54	Evidence for Effective $p(Z) \text{â€} \text{f}(\text{Ar})$ Conjugations (Z = S, Se, and Te, as Well as Z = O) in 9-(Arylchalcogenyl)tritycenes: Experimental and Theoretical Investigations. <i>Journal of Organic Chemistry</i> , 2009, 74, 4763-4771.	3.2	12

#	ARTICLE	IF	CITATIONS
55	Polar Coordinate Representation of $\langle i \rangle H \langle /i \rangle \langle sub \rangle b \langle /sub \rangle (\langle i \rangle r \langle /i \rangle \langle sub \rangle c \langle /sub \rangle)$ versus $(\hat{a}, \langle sup \rangle 2 \langle /sup \rangle / 8 \langle i \rangle m \langle /i \rangle) \hat{a}^{-1/2} \langle sup \rangle 2 \langle /sup \rangle \langle sub \rangle b \langle /sub \rangle (\langle i \rangle r \langle /i \rangle \langle sub \rangle c \langle /sub \rangle)$ at BCP in AIM Analysis: Classification and Evaluation of Weak to Strong Interactions. <i>Journal of Physical Chemistry A</i> , 2009, 113, 10050-7.	2.5	106
56	Structures and dynamic stereochemistry of 9-arylselanyltritycenes: X-ray crystallographic, spectroscopic and theoretical investigations. <i>New Journal of Chemistry</i> , 2009, 33, 1588.	2.8	1
57	Imido transfer of sulfonylimino- λ^3 -bromane makes possible the synthesis of sulfonylimino- λ^3 -iodanes. <i>Chemical Communications</i> , 2009, , 959.	4.1	18
58	How does non-covalent $Se \cdots Se \cdots O$ interaction stabilize selenoxides at naphthalene 1,8-positions: structural and theoretical investigations. <i>New Journal of Chemistry</i> , 2009, 33, 196-206.	2.8	15
59	How Are Non-Bonded $G \cdots A \cdots Z$ (Z = O, S, and Se) Distances at Benzene 1,2-, Naphthalene 1,8-, and Anthracene 1,8,9-Positions Controlled? An Approach to Causality in Weak Interactions. <i>Bulletin of the Chemical Society of Japan</i> , 2009, 82, 712-722.	3.2	8
60	Torsional Angular Dependence of $\langle sup \rangle 1 \langle /sup \rangle \langle i \rangle J \langle /i \rangle$ (Se,Se) and Fermi Contact Control of $\langle sup \rangle 4 \langle /sup \rangle \langle i \rangle J \langle /i \rangle$ (Se,Se): Analysis of $\langle sup \rangle \langle i \rangle n \langle /sup \rangle \langle i \rangle J \langle /i \rangle$ (Se,Se) ($\langle i \rangle n \langle /i \rangle = 1 \cdots 4$) Based on Molecular Orbital Theory. <i>Chemistry - A European Journal</i> , 2008, 14, 5645-5655.	3.3	20
61	Evaluation of Electron Population Terms for $\hat{a}^{\langle i \rangle r \langle /i \rangle \langle sub \rangle Se \langle /sub \rangle \langle sup \rangle \hat{a}^{\langle i \rangle 3 \langle /sup \rangle} \hat{a}^{\langle i \rangle 4 p \langle /sub \rangle}$, $\hat{a}^{\langle i \rangle r \langle /i \rangle \langle sub \rangle S \langle /sub \rangle \langle sup \rangle \hat{a}^{\langle i \rangle 3 \langle /sup \rangle} \hat{a}^{\langle i \rangle 3 p \langle /sub \rangle}$, and $\hat{a}^{\langle i \rangle r \langle /i \rangle \langle sub \rangle O \langle /sub \rangle \langle sup \rangle \hat{a}^{\langle i \rangle 3 \langle /sup \rangle} \hat{a}^{\langle i \rangle 2 p \langle /sub \rangle}$: How Do HOMO and LUMO Shrink or Expand Depending on Nuclear Charges?. <i>Chemistry - A European Journal</i> , 2008, 14, 7278-7284.	3.3	5
62	Contributions from Atomic p(Se), d(Se), and f(Se) Orbitals to Absolute Paramagnetic Shielding Tensors in Neutral and Charged $SeH \langle sub \rangle \langle i \rangle n \langle /sub \rangle \langle /sub \rangle$ and Some Oxides Including the Effect of Methyl and Halogen Substitutions on $f \langle sup \rangle p \langle /sup \rangle$ (Se). <i>Chemistry - A European Journal</i> , 2008, 14, 9647-9655.	3.3	5
63	Fine structures of 8-G-1-(arylethynylselanyl)naphthalenes (G = H, Cl, Br): Factors to control the linear alignment of five $G \cdots Se \cdots C \cdots C \cdots Ar$ atoms in crystals and the behavior in solution. <i>Polyhedron</i> , 2008, 27, 3557-3566.	2.2	12
64	Atoms-in-Molecules Dual Parameter Analysis of Weak to Strong Interactions: Behaviors of Electronic Energy Densities versus Laplacian of Electron Densities at Bond Critical Points. <i>Journal of Physical Chemistry A</i> , 2008, 112, 13593-13599.	2.5	152
65	Extended hypervalent $E \cdots \hat{a} \cdots E \cdots E \cdots 4c \cdots 6e$ (E, $E \cdots =$ Se, S) interactions: structure, stability and reactivity of 1-(8-Ph $E \cdots C_{10}H_6$) $EE(C_{10}H_6E \cdots Ph-8 \hat{a} \cdots)-1 \hat{a} \cdots$. <i>New Journal of Chemistry</i> , 2008, 32, 1881.	2.8	26
66	$\langle sup \rangle 77 \langle /sup \rangle$ Se NMR Chemical Shifts of 9-(Arylselanyl)tritycenes: New Standard for Planar Structures of ArSeR and Applications to Determine the Structures in Solutions. <i>Journal of Organic Chemistry</i> , 2008, 73, 9259-9269.	3.2	11
67	Origin of $\langle sup \rangle 77 \langle /sup \rangle$ Se NMR Chemical Shifts Revealed for Pre- \hat{I}_{\pm} , \hat{I}_{\pm}^2 , and \hat{I}^3 Effects. <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , 2008, 183, 1067-1071.	1.6	3
68	Noncovalent $Z \cdots A \cdots Z$ (Z=O, S, Se, and Te) Interactions: How Do They Operate to Control Fine Structures of 1,8-Dichalcogene-Substituted Naphthalenes?. <i>Bulletin of the Chemical Society of Japan</i> , 2008, 81, 1605-1615.	3.2	37
69	A New Type of Imido Group Donor: $\hat{a}^{\langle i \rangle 3 \langle /sup \rangle}$ Synthesis and Characterization of Sulfonylimino- λ^3 -bromane that Acts as a Nitrenoid in the Aziridination of Olefins at Room Temperature under Metal-Free Conditions. <i>Journal of the American Chemical Society</i> , 2007, 129, 12938-12939.	13.7	70
70	Orientational Effect of Aryl Groups in Aryl Selenides: How Can 1H and ^{13}C NMR Chemical Shifts Clarify the Effect?. <i>Journal of Organic Chemistry</i> , 2007, 72, 7587-7596.	3.2	22
71	Atoms-in-Molecules Analysis of Extended Hypervalent Five-Center, Six-Electron (5c-6e) $C_{22}O$ Interactions at the 1,8,9-Positions of Anthraquinone and 9-Methoxyanthracene Systems. <i>Chemistry - A European Journal</i> , 2007, 13, 255-268.	3.3	77
72	How ^{77}Se NMR Chemical Shifts Originate from Pre- \hat{I}_{\pm} , \hat{I}_{\pm}^2 , and \hat{I}^3 Effects: Interpretation Based on Molecular Orbital Theory. <i>Chemistry - A European Journal</i> , 2007, 13, 5282-5293.	3.3	13

#	ARTICLE	IF	CITATIONS
73	Structures of 1-(Arylseleninyl)naphthalenes: λ O, G, and Y Dependences in 8-G-1-[p-YC ₆ H ₄ Se(O)]C ₁₀ H ₆ . <i>Journal of Organic Chemistry</i> , 2006, 71, 5574-5585.	3.2	48
74	Proposal for Sets of ⁷⁷ Se NMR Chemical Shifts in Planar and Perpendicular Orientations of Aryl Group and the Applications. <i>Bioinorganic Chemistry and Applications</i> , 2006, 2006, 1-13.	4.1	8
75	Orientalional Effect of Aryl Groups on ⁷⁷ Se NMR Chemical Shifts: Experimental and Theoretical Investigations. <i>Chemistry - A European Journal</i> , 2006, 12, 3829-3846.	3.3	38
76	The Structure of 1-(Arylthio)naphthalenes, Together with the Selenium and Oxygen Derivatives in Crystals and Solutions. <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , 2005, 180, 1431-1432.	1.6	1
77	How Do δ -Se Nonbonded and γ -Se Through π -Bond Interactions Determine the Structures of 8-G-1-(λ) Tj ETQ ₀ 1 1.0.784314 rgB...	1.6	0
78	Extended Hypervalent 5c \sim 6e Interactions: Linear Alignment of Five C \sim Z \sim O \sim Z \sim C (Z = S, Se) Atoms in Anthraquinone and Anthracene Systems. <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , 2005, 180, 1351-1355.	1.6	8
79	First Br ₄ four centre \sim six electron and Se ₂ Br ₅ seven centre \sim ten electron bonds in nonionic bromine adducts of selenanthrene. <i>Chemical Communications</i> , 2004, , 140-141.	4.1	24
80	Extended Hypervalent 5c \sim 6e Interactions: A Linear Alignment of Five C \sim Se \sim O \sim Se \sim C Atoms in Anthraquinone and 9-Methoxyanthracene Bearing Arylselanyl Groups at the 1,8-Positions. <i>Journal of Organic Chemistry</i> , 2004, 69, 1676-1684.	3.2	35
81	First linear alignment of five C \sim Se \sim O \sim Se \sim C atoms in anthraquinone and 9-(methoxy)anthracene bearing phenylselanyl groups at 1,8-positions. <i>Chemical Communications</i> , 2003, , 124-125.	4.1	48
82	Nonbonded P \sim A \sim P and P \sim A \sim Se Interactions in Naphthalene 1,8-Positions: Role of Lone-Pair Orbitals. <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , 2002, 177, 1833-1837.	1.6	27
83	Structure of 1-(Arylselanyl)naphthalenes. 2. G Dependence in 8-G-1-(p-YC ₆ H ₄ Se)C ₁₀ H ₆ . <i>Journal of Organic Chemistry</i> , 2002, 67, 38-48.	3.2	79
84	Linear alignment of four sulfur atoms in bis[(8-phenylthio)naphthyl] disulfide: contribution of linear S ₄ hypervalent four-centre six-electron bond to the structure. <i>Chemical Communications</i> , 2002, , 2416-2417.	4.1	49
85	Structural studies on diaryl selenide dihalides in solution: molecular complex formation of substituted diphenyl selenides with bromine. <i>Perkin Transactions II RSC</i> , 2002, , 262-270.	1.1	13
86	On the equilibrium between molecular complexes and trigonal bipyramidal adducts of diaryl selenide dibromides in solution. <i>Heteroatom Chemistry</i> , 2001, 12, 369-379.	0.7	10
87	Structure of 1-(Arylselanyl)naphthalenes λ Y Dependence in 1-(p-YC ₆ H ₄ Se)C ₁₀ H ₇ . <i>European Journal of Organic Chemistry</i> , 2001, 2001, 3933-3943.	2.4	49
88	Successive Change in Conformation Caused by p-Y Groups in 1-(MeSe)-8-(p-YC ₆ H ₄ Se)C ₁₀ H ₆ : A Role of Linear Se \sim A \sim Se \sim C Three-Center \sim Four-Electron versus n(Se) \sim A \sim n(Se) Two-Center \sim Four-Electron Nonbonded Interactions. <i>Journal of Physical Chemistry A</i> , 1999, 103, 9906-9912.	2.5	64
89	Novel Substituent Effect on ⁷⁷ Se NMR Chemical Shifts Caused by 4c-6e versus 2c-4e and 3c-4e in Naphthalene Peri Positions: A Spectroscopic and Theoretical Study. <i>Journal of Organic Chemistry</i> , 1999, 64, 6688-6696.	3.2	102
90	Structural Study of Aryl Selenides in Solution Based on ⁷⁷ Se NMR Chemical Shifts: A Application of the GIAO Magnetic Shielding Tensor of the ⁷⁷ Se Nucleus. <i>Journal of Physical Chemistry A</i> , 1999, 103, 6074-6081.	2.5	52

#	ARTICLE	IF	CITATIONS
91	Four-Center Six-Electron Interaction versus Lone Pair-Lone Pair Interaction between Selenium Atoms in Naphthalene Peri Positions. <i>Journal of Organic Chemistry</i> , 1998, 63, 8790-8800.	3.2	79
92	Attractive Interaction Caused by the Linear F-Å-Å-Se-C Alignment in Naphthalene Peri Positions. <i>Journal of the American Chemical Society</i> , 1998, 120, 3635-3640.	13.7	104
93	On the Factors to Determine ⁷⁷ Se NMR Chemical Shifts of Organic Selenium Compounds: Application of GIAO Magnetic Shielding Tensor to the ⁷⁷ Se NMR Spectroscopy. <i>Chemistry Letters</i> , 1998, 27, 523-524.	1.3	24
94	Structure of bis[8-(phenylselanyl)naphthyl] diselenide: first linear alignment of four Se atoms as a four-centre six-electron bond. <i>Chemical Communications</i> , 1996, , 371.	4.1	78
95	Inverse Substituent Effect on ⁷⁷ Se NMR Chemical Shifts in Naphthalene Systems with Linear 4c-6e Se ₄ Bond: 1-[8-(p- ⁶ H ₄ Se)C ₁₀ H ₆]SeSe[C ₁₀ H ₆ (SeC ₆ H ₄ -p)-8]-1 vs. 1-(MeSe)-8-(p- ⁶ H ₄ Se)C ₁₀ H ₆ . <i>Chemistry Letters</i> , 1996, 25, 947-948.		59
96	Molecular Complex Formation of Diphenyl Selenides with Bromine: Electronic and Steric Effects. Phosphorus, Sulfur and Silicon and the Related Elements, 1992, 67, 79-82.	1.6	9
97	Structural studies of halogen adducts of diorganyl chalcogenides in solutions by ¹ H, ¹³ C, ⁷⁷ Se and ¹²⁵ Te NMR. <i>Journal of Physical Organic Chemistry</i> , 1990, 3, 358-368.	1.9	62
98	Structural studies of halogen adducts of some cyclic selenides and tellurides by ¹ H, ¹³ C, ⁷⁷ Se and ¹²⁵ Te NMR. Evidence for the formation of molecular complexes of selenoxanthone and selenanthrene with bromine. <i>Journal of Physical Organic Chemistry</i> , 1990, 3, 369-374.	1.9	44