

Yoshitaka Hashimoto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6603755/publications.pdf>

Version: 2024-02-01

169
papers

6,402
citations

147801

31
h-index

76900

74
g-index

178
all docs

178
docs citations

178
times ranked

7518
citing authors

#	ARTICLE	IF	CITATIONS
1	The Metabolic Syndrome as a Predictor of Nonalcoholic Fatty Liver Disease. <i>Annals of Internal Medicine</i> , 2005, 143, 722.	3.9	911
2	The Severity of Ultrasonographic Findings in Nonalcoholic Fatty Liver Disease Reflects the Metabolic Syndrome and Visceral Fat Accumulation. <i>American Journal of Gastroenterology</i> , 2007, 102, 2708-2715.	0.4	688
3	Association of Non-alcoholic Fatty Liver Disease with Chronic Kidney Disease: A Systematic Review and Meta-analysis. <i>PLoS Medicine</i> , 2014, 11, e1001680.	8.4	507
4	Nonalcoholic fatty liver disease is a novel predictor of cardiovascular disease. <i>World Journal of Gastroenterology</i> , 2007, 13, 1579.	3.3	469
5	Asiaâ€‘Pacific Working Party on Nonâ€‘alcoholic Fatty Liver Disease guidelines 2017â€‘Part 1: Definition, risk factors and assessment. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018, 33, 70-85.	2.8	358
6	Soy protein isolate and its hydrolysate reduce body fat of dietary obese rats and genetically obese mice (yellow KK). <i>Nutrition</i> , 2000, 16, 349-354.	2.4	180
7	Ectopic fat obesity presents the greatest risk for incident type 2 diabetes: a population-based longitudinal study. <i>International Journal of Obesity</i> , 2019, 43, 139-148.	3.4	164
8	Metabolically Healthy Obesity and Risk of Incident CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2015, 10, 578-583.	4.5	129
9	The impact of nonâ€‘alcoholic fatty liver disease on incident type 2 diabetes mellitus in nonâ€‘overweight individuals. <i>Liver International</i> , 2016, 36, 275-283.	3.9	125
10	The Asiaâ€‘Pacific Working Party on Nonâ€‘alcoholic Fatty Liver Disease guidelines 2017â€‘Part 2: Management and special groups. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018, 33, 86-98.	2.8	117
11	Ageing is a risk factor of nonalcoholic fatty liver disease in premenopausal women. <i>World Journal of Gastroenterology</i> , 2012, 18, 237.	3.3	114
12	Decreased the creatinine to cystatin C ratio is a surrogate marker of sarcopenia in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018, 139, 52-58.	2.8	108
13	The modest alcohol consumption reduces the incidence of fatty liver in men: a populationâ€‘based largeâ€‘scale cohort study. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2015, 30, 546-552.	2.8	102
14	Nonoverweight nonalcoholic fatty liver disease and incident cardiovascular disease. <i>Medicine (United States)</i> , 2017, 96, e6712.	1.0	86
15	Identification of individuals with non-alcoholic fatty liver disease by the diagnostic criteria for the metabolic syndrome. <i>World Journal of Gastroenterology</i> , 2012, 18, 1508.	3.3	84
16	Reduction by Phytate-reduced Soybean Î²-Conglycinin of Plasma Triglyceride Level of Young and Adult Rats. <i>Bioscience, Biotechnology and Biochemistry</i> , 2001, 65, 1071-1075.	1.3	75
17	Triglycerides to highâ€‘density lipoprotein cholesterol ratio is an independent predictor of incident fatty liver; a populationâ€‘based cohort study. <i>Liver International</i> , 2016, 36, 713-720.	3.9	75
18	Late-night-dinner is associated with poor glycemic control in people with type 2 diabetes: The KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2018, 65, 395-402.	1.6	73

#	ARTICLE	IF	CITATIONS
19	Impact of low-carbohydrate diet on body composition: meta-analysis of randomized controlled studies. <i>Obesity Reviews</i> , 2016, 17, 499-509.	6.5	72
20	Shortage of energy intake rather than protein intake is associated with sarcopenia in elderly patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort. <i>Journal of Diabetes</i> , 2019, 11, 477-483.	1.8	61
21	Effect of coronavirus disease 2019 pandemic on the lifestyle and glycemic control in patients with type 2 diabetes: a cross-section and retrospective cohort study. <i>Endocrine Journal</i> , 2021, 68, 201-210.	1.6	59
22	The relationship between hepatic steatosis and skeletal muscle mass index in men with type 2 diabetes. <i>Endocrine Journal</i> , 2016, 63, 877-884.	1.6	57
23	The Triglyceride and Glucose Index Is a Predictor of Incident Nonalcoholic Fatty Liver Disease: A Population-Based Cohort Study. <i>Canadian Journal of Gastroenterology and Hepatology</i> , 2019, 2019, 1-7.	1.9	55
24	Lower vegetable protein intake and higher dietary acid load associated with lower carbohydrate intake are risk factors for metabolic syndrome in patients with type 2 diabetes: a cross-sectional study. <i>Journal of Diabetes Investigation</i> , 2015, 6, 465-472.	2.4	40
25	Fatty liver as a risk factor for progression from metabolically healthy to metabolically abnormal in non-overweight individuals. <i>Endocrine</i> , 2017, 57, 89-97.	2.3	39
26	Hemoglobin concentration and incident metabolic syndrome: a population-based large-scale cohort study. <i>Endocrine</i> , 2015, 50, 390-396.	2.3	38
27	Sarcopenia is associated with blood pressure variability in older patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort study. <i>Geriatrics and Gerontology International</i> , 2018, 18, 1345-1349.	1.5	36
28	Sarcopenia is associated with tongue pressure in older patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort study. <i>Geriatrics and Gerontology International</i> , 2019, 19, 153-158.	1.5	36
29	Intake of sucrose affects gut dysbiosis in patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1623-1634.	2.4	35
30	Transient remission of nonalcoholic fatty liver disease decreases the risk of incident type 2 diabetes mellitus in Japanese men. <i>European Journal of Gastroenterology and Hepatology</i> , 2016, 28, 1443-1449.	1.6	34
31	Relationship between nonalcoholic fatty liver disease and muscle quality as well as quantity evaluated by computed tomography. <i>Liver International</i> , 2020, 40, 120-130.	3.9	34
32	Impact of low-carbohydrate diet on renal function: a meta-analysis of over 1000 individuals from nine randomised controlled trials. <i>British Journal of Nutrition</i> , 2016, 116, 632-638.	2.3	33
33	Short Sleep Duration is a Risk of Incident Nonalcoholic Fatty Liver Disease: A Population-based Longitudinal Study. <i>Journal of Gastrointestinal and Liver Diseases</i> , 2019, 28, 73-81.	0.9	32
34	Sarcopenia Is Associated With a Risk of Mortality in People With Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , 2021, 12, 783363.	3.5	32
35	Metabolically healthy obesity without fatty liver and risk of incident type 2 diabetes: A meta-analysis of prospective cohort studies. <i>Obesity Research and Clinical Practice</i> , 2018, 12, 4-15.	1.8	30
36	Non-alcoholic fatty liver disease with obesity as an independent predictor for incident gastric and colorectal cancer: a population-based longitudinal study. <i>BMJ Open Gastroenterology</i> , 2019, 6, e000295.	2.7	29

#	ARTICLE	IF	CITATIONS
37	The effect of COVID-19 pandemic on the lifestyle and glycemic control in patients with type 1 diabetes: a retrospective cohort study. <i>Diabetology International</i> , 2022, 13, 85-90.	1.4	29
38	Triglyceride-glucose index is a predictor of incident chronic kidney disease: a population-based longitudinal study. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 948-955.	1.6	27
39	Combined effect of body mass index and waist-height ratio on incident diabetes; a population based cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2017, 61, 118-122.	1.4	26
40	Divided consumption of late-night-dinner improves glycemic excursions in patients with type 2 diabetes: A randomized cross-over clinical trial. <i>Diabetes Research and Clinical Practice</i> , 2017, 129, 206-212.	2.8	25
41	Group 3 Innate Lymphoid Cells Protect Steatohepatitis From High-Fat Diet Induced Toxicity. <i>Frontiers in Immunology</i> , 2021, 12, 648754.	4.8	25
42	Reduced dietary omega-3 fatty acids intake is associated with sarcopenia in elderly patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020, 66, 233-237.	1.4	24
43	Relationship between metabolic syndrome and trunk muscle quality as well as quantity evaluated by computed tomography. <i>Clinical Nutrition</i> , 2020, 39, 1818-1825.	5.0	23
44	Triglyceride-glucose index (TyG index) is a predictor of incident colorectal cancer: a population-based longitudinal study. <i>BMC Endocrine Disorders</i> , 2020, 20, 113.	2.2	23
45	The Visceral Adiposity Index Is a Predictor of Incident Chronic Kidney Disease: A Population-Based Longitudinal Study. <i>Kidney and Blood Pressure Research</i> , 2020, 45, 407-418.	2.0	23
46	Erythritol Ameliorates Small Intestinal Inflammation Induced by High-Fat Diets and Improves Glucose Tolerance. <i>International Journal of Molecular Sciences</i> , 2021, 22, 5558.	4.1	23
47	Short energy intake is associated with muscle mass loss in older patients with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Clinical Nutrition</i> , 2021, 40, 1613-1620.	5.0	22
48	Trans Fatty Acid Intake Induces Intestinal Inflammation and Impaired Glucose Tolerance. <i>Frontiers in Immunology</i> , 2021, 12, 669672.	4.8	22
49	U-shaped association between the triglyceride-glucose index and the risk of incident diabetes in people with normal glycemic level: A population-base longitudinal cohort study. <i>Clinical Nutrition</i> , 2021, 40, 1555-1561.	5.0	22
50	Metabolic associated fatty liver disease is a risk factor for chronic kidney disease. <i>Journal of Diabetes Investigation</i> , 2022, 13, 308-316.	2.4	22
51	BMI history and risk of incident fatty liver. <i>European Journal of Gastroenterology and Hepatology</i> , 2016, 28, 1188-1193.	1.6	21
52	Urinary pH is a predictor of diabetes in men; a population based large scale cohort study. <i>Diabetes Research and Clinical Practice</i> , 2017, 130, 9-14.	2.8	21
53	Protein Intake, Especially Vegetable Protein Intake, Is Associated with Higher Skeletal Muscle Mass in Elderly Patients with Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2017, 2017, 1-7.	2.3	21
54	The sodium-glucose cotransporter 2 inhibitor luseogliflozin can suppress muscle atrophy in Db/Db mice by suppressing the expression of <i>foxo1</i> . <i>Journal of Clinical Biochemistry and Nutrition</i> , 2019, 65, 23-28.	1.4	21

#	ARTICLE	IF	CITATIONS
55	The Effects of Metformin on the Gut Microbiota of Patients with Type 2 Diabetes: A Two-Center, Quasi-Experimental Study. <i>Life</i> , 2020, 10, 195.	2.4	20
56	High brain natriuretic peptide is associated with sarcopenia in patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2019, 66, 369-377.	1.6	19
57	Weight gain since age of 20 as risk of metabolic syndrome even in non-overweight individuals. <i>Endocrine</i> , 2017, 58, 253-261.	2.3	18
58	Reduction of Fat to Muscle Mass Ratio Is Associated with Improvement of Liver Stiffness in Diabetic Patients with Non-Alcoholic Fatty Liver Disease. <i>Journal of Clinical Medicine</i> , 2019, 8, 2175.	2.4	18
59	Skipping breakfast is associated with glycemic variability in patients with type 2 diabetes. <i>Nutrition</i> , 2020, 71, 110639.	2.4	18
60	Immune modulating effects of additional supplementation of estradiol combined with testosterone in murine testosterone-deficient NAFLD model. <i>American Journal of Physiology - Renal Physiology</i> , 2020, 318, G989-G999.	3.4	18
61	Impact of metabolically healthy obesity on the risk of incident gastric cancer: a population-based cohort study. <i>BMC Endocrine Disorders</i> , 2020, 20, 11.	2.2	18
62	Association between Geriatric Nutrition Risk Index and The Presence of Sarcopenia in People with Type 2 Diabetes Mellitus: A Cross-Sectional Study. <i>Nutrients</i> , 2021, 13, 3729.	4.1	18
63	Urinary pH reflects dietary acid load in patients with type 2 diabetes. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2017, 61, 74-77.	1.4	17
64	Intake of Carbohydrate to Fiber Ratio Is a Useful Marker for Metabolic Syndrome in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Annals of Nutrition and Metabolism</i> , 2018, 72, 329-335.	1.9	17
65	Low-attenuation muscle is a predictor of diabetes mellitus: A population-based cohort study. <i>Nutrition</i> , 2020, 74, 110752.	2.4	17
66	ILC2s Improve Glucose Metabolism Through the Control of Saturated Fatty Acid Absorption Within Visceral Fat. <i>Frontiers in Immunology</i> , 2021, 12, 669629.	4.8	17
67	Sodium-chloride Difference and Metabolic Syndrome: A Population-based Large-scale Cohort Study. <i>Internal Medicine</i> , 2016, 55, 3085-3090.	0.7	16
68	Creatinine-to-bodyweight ratio is a predictor of incident non-alcoholic fatty liver disease: A population-based longitudinal study. <i>Hepatology Research</i> , 2020, 50, 57-66.	3.4	16
69	Association between sleep disorder and quality of life in patients with type 2 diabetes: a cross-sectional study. <i>BMC Endocrine Disorders</i> , 2020, 20, 98.	2.2	16
70	Creatinine/(cystatin C \times body weight) ratio is associated with skeletal muscle mass index. <i>Endocrine Journal</i> , 2020, 67, 733-740.	1.6	16
71	Habitual Dietary Intake Affects the Altered Pattern of Gut Microbiome by Acarbose in Patients with Type 2 Diabetes. <i>Nutrients</i> , 2021, 13, 2107.	4.1	16
72	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with a Low Prevalence of Sarcopenia in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021, 13, 72.	4.1	16

#	ARTICLE	IF	CITATIONS
73	Impact of different timing of consuming sweet snack on postprandial glucose excursions in healthy women. <i>Diabetes and Metabolism</i> , 2019, 45, 369-374.	2.9	15
74	Effect of Exercise Habit on Skeletal Muscle Mass Varies with Protein Intake in Elderly Patients with Type 2 Diabetes: A Retrospective Cohort Study. <i>Nutrients</i> , 2020, 12, 3220.	4.1	15
75	Vitamin Intake and Loss of Muscle Mass in Older People with Type 2 Diabetes: A Prospective Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , 2021, 13, 2335.	4.1	15
76	Platelet to lymphocyte ratio correlates with diabetic foot risk and foot ulcer in patients with type 2 diabetes. <i>Endocrine Journal</i> , 2019, 66, 905-913.	1.6	14
77	Frequent Usage of Convenience Stores is Associated with Low Diet Quality. <i>Nutrients</i> , 2019, 11, 1212.	4.1	14
78	Relationship between limited joint mobility of the hand and diabetic foot risk in patients with type 2 diabetes. <i>Journal of Diabetes</i> , 2017, 9, 628-633.	1.8	13
79	Relationship between skeletal muscle mass and hepatic fibrosis in patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2017, 43, 184-186.	2.9	13
80	Divided consumption of late-night-dinner improves glucose excursions in young healthy women: A randomized cross-over clinical trial. <i>Diabetes Research and Clinical Practice</i> , 2018, 136, 78-84.	2.8	13
81	Consuming snacks mid-afternoon compared with just after lunch improves mean amplitude of glycaemic excursions in patients with type 2 diabetes: A randomized crossover clinical trial. <i>Diabetes and Metabolism</i> , 2018, 44, 482-487.	2.9	12
82	Potential impact of the joint association of total bilirubin and gamma-glutamyltransferase with metabolic syndrome. <i>Diabetology and Metabolic Syndrome</i> , 2019, 11, 12.	2.7	12
83	Effect of alcohol consumption and the presence of fatty liver on the risk for incident type 2 diabetes: a population-based longitudinal study. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001629.	2.8	12
84	Eating Fast Has a Significant Impact on Glycemic Excursion in Healthy Women: Randomized Controlled Cross-Over Trial. <i>Nutrients</i> , 2020, 12, 2767.	4.1	12
85	Impact of fatty liver disease and metabolic syndrome on incident type 2 diabetes; a population based cohort study. <i>Endocrine Journal</i> , 2017, 64, 1105-1114.	1.6	11
86	Impact of extracellular to intracellular fluid volume ratio on albuminuria in patients with type 2 diabetes: A cross-sectional and longitudinal cohort study. <i>Journal of Diabetes Investigation</i> , 2021, 12, 1202-1211.	2.4	11
87	Japanese radio calisthenics prevents the reduction of skeletal muscle mass volume in people with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001027.	2.8	11
88	Association between Sleep Duration and Incident Chronic Kidney Disease: A Population-Based Cohort Analysis of the NAGALA Study. <i>Kidney and Blood Pressure Research</i> , 2020, 45, 339-349.	2.0	11
89	Effect of COVID-19 Pandemic on the Change in Skeletal Muscle Mass in Older Patients with Type 2 Diabetes: A Retrospective Cohort Study. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 4188.	2.6	11
90	Low urine pH is a risk for non-alcoholic fatty liver disease: A population-based longitudinal study. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2018, 42, 570-576.	1.5	10

#	ARTICLE	IF	CITATIONS
91	Eating Fast Is Associated with Nonalcoholic Fatty Liver Disease in Men But Not in Women with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020, 12, 2174.	4.1	10
92	Sarcopenic obesity is associated with macroalbuminuria in patients with type 2 diabetes: a cross-sectional study. <i>Endocrine Journal</i> , 2021, 68, 781-789.	1.6	10
93	The visceral adiposity index is a predictor of incident nonalcoholic fatty liver disease: A population-based longitudinal study. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2020, 44, 375-383.	1.5	10
94	High-sensitivity cardiac troponin T is associated with coronary artery calcification. <i>Journal of Cardiovascular Computed Tomography</i> , 2015, 9, 209-214.	1.3	9
95	Heart rate-corrected QT interval is a novel risk marker for the progression of albuminuria in people with Type 2 diabetes. <i>Diabetic Medicine</i> , 2015, 32, 1221-1226.	2.3	9
96	Malnutrition assessed by controlling nutritional status is correlated to carotid atherosclerosis in patients with type 2 diabetes. <i>Endocrine Journal</i> , 2019, 66, 1073-1082.	1.6	9
97	Distinct associations of intraperitoneal and retroperitoneal visceral adipose tissues with metabolic syndrome and its components. <i>Clinical Nutrition</i> , 2021, 40, 3479-3484.	5.0	9
98	miR-23b-3p acts as a counter-response against skeletal muscle atrophy. <i>Journal of Endocrinology</i> , 2020, 244, 535-547.	2.6	9
99	Partially Hydrolyzed Guar Gum Suppresses the Development of Sarcopenic Obesity. <i>Nutrients</i> , 2022, 14, 1157.	4.1	9
100	Relationship between limited joint mobility of hand and carotid atherosclerosis in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2017, 132, 79-84.	2.8	8
101	Visceral Adiposity Index is a predictor of incident colorectal cancer: a population-based longitudinal study. <i>BMJ Open Gastroenterology</i> , 2020, 7, e000400.	2.7	8
102	Low circulating dihomo-gamma-linolenic acid is associated with diabetic retinopathy: a cross sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2021, 68, 421-428.	1.6	8
103	Trunk muscle quality and quantity predict the development of metabolic syndrome and the increase in the number of its components in individuals without metabolic syndrome. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020, 30, 1161-1168.	2.6	8
104	Effects of probiotic <i>Bifidobacterium bifidum</i> on the gastrointestinal symptoms of patients with type 2 diabetes mellitus treated with metformin: An open-label, single-arm, exploratory research trial. <i>Journal of Diabetes Investigation</i> , 2022, 13, 489-500.	2.4	8
105	Intraperitoneal, but not retroperitoneal, visceral adipose tissue is associated with diabetes mellitus: a cross-sectional, retrospective pilot analysis. <i>Diabetology and Metabolic Syndrome</i> , 2020, 12, 103.	2.7	7
106	Changes in metabolic complications in patients with alcoholic fatty liver disease monitored over two decades: NAGALA study. <i>BMJ Open Gastroenterology</i> , 2020, 7, e000359.	2.7	7
107	Unique Habitual Food Intakes in the Gut Microbiota Cluster Associated with Type 2 Diabetes Mellitus. <i>Nutrients</i> , 2021, 13, 3816.	4.1	7
108	Nutritional Status Assessed with Objective Data Assessment Correlates with a High-Risk Foot in Patients with Type 2 Diabetes. <i>Journal of Clinical Medicine</i> , 2022, 11, 1314.	2.4	7

#	ARTICLE	IF	CITATIONS
109	The Association Between Taste Impairment and Serum Zinc Concentration in Adult Patients With Type 2 Diabetes. <i>Canadian Journal of Diabetes</i> , 2018, 42, 520-524.	0.8	6
110	Metabolically healthy obesity and risk of leukoaraiosis; a population based cross-sectional study. <i>Endocrine Journal</i> , 2018, 65, 669-675.	1.6	6
111	Understanding of antidiabetic medication is associated with blood glucose in patients with type 2 diabetes: At baseline date of the KAMOGAWA-DM cohort study. <i>Journal of Diabetes Investigation</i> , 2019, 10, 458-465.	2.4	6
112	Effects of dietary salt restriction on home blood pressure in diabetic patients with excessive salt intake: a pilot study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2019, 65, 252-257.	1.4	6
113	Trigger finger is associated with risk of incident cardiovascular disease in individuals with type 2 diabetes: a retrospective cohort study. <i>BMJ Open Diabetes Research and Care</i> , 2021, 9, e002070.	2.8	6
114	The Risk Factors for Development of Type 2 Diabetes: Panasonic Cohort Study 4. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 571.	2.6	6
115	Which Measurement of Blood Pressure Is More Associated With Albuminuria in Patients With Type 2 Diabetes: Central Blood Pressure or Peripheral Blood Pressure?. <i>Journal of Clinical Hypertension</i> , 2016, 18, 790-795.	2.0	5
116	Caffeine intake enhances the benefits of sodium glucose transporter 2 inhibitor. <i>Diabetes/Metabolism Research and Reviews</i> , 2016, 32, 694-699.	4.0	5
117	Impact of respiratory function on the progression from metabolically healthy non-overweight to metabolically abnormal phenotype. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018, 28, 922-928.	2.6	5
118	Neutrophil-lymphocyte ratio correlates with limited joint mobility of hand in patients with type 2 diabetes. <i>Endocrine Journal</i> , 2018, 65, 1011-1017.	1.6	5
119	Combined effect of hemoglobin and mean corpuscular volume levels on incident metabolic syndrome: A population-based cohort study. <i>Clinical Nutrition ESPEN</i> , 2020, 40, 314-319.	1.2	5
120	Handgrip measurement as a useful benchmark for locomotive syndrome in patients with type 2 diabetes mellitus: A KAMOGAWA-DM cohort study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1602-1611.	2.4	5
121	Microbe-associated metabolites as targets for incident type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2021, 12, 476-478.	2.4	5
122	Serum N-terminal Pro-brain Natriuretic Peptide Level is Associated with the Development of Chronic Kidney Diseases in Patients with Type 2 Diabetes. <i>Endocrine, Metabolic and Immune Disorders - Drug Targets</i> , 2018, 18, 590-595.	1.2	5
123	Eating Speed Is Associated with the Presence of Sarcopenia in Older Patients with Type 2 Diabetes: A Cross-Sectional Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , 2022, 14, 759.	4.1	5
124	Relationship between serum creatinine to cystatin C ratio and subclinical atherosclerosis in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2022, 10, e002910.	2.8	5
125	Serum levels of mac-2 binding protein are associated with diabetic microangiopathy and macroangiopathy in people with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001189.	2.8	4
126	Liver Stiffness Is Associated With Progression of Albuminuria in Adults With Type 2 Diabetes: Nonalcoholic Fatty Disease Cohort Study. <i>Canadian Journal of Diabetes</i> , 2020, 44, 428-433.	0.8	4

#	ARTICLE	IF	CITATIONS
127	Creatinine to Body Weight Ratio Is Associated with Incident Diabetes: Population-Based Cohort Study. <i>Journal of Clinical Medicine</i> , 2020, 9, 227.	2.4	4
128	Living alone is associated with visit-to-visit HbA1c variability in men but not in women in people with type 2 diabetes: KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2020, 67, 419-426.	1.6	4
129	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with Glycemic Variability in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021, 13, 1488.	4.1	4
130	Clinical characteristics and longitudinal changes of patients with non-alcoholic fatty liver disease in 2Âdecades: the NAGALA study. <i>BMC Gastroenterology</i> , 2021, 21, 223.	2.0	4
131	Obesity and metabolic abnormalities as risks of alcoholic fatty liver in men: NAGALA study. <i>BMC Gastroenterology</i> , 2021, 21, 321.	2.0	4
132	Asymptomatic postprandial hypotension in patients with diabetes: The KAMOGAWAâ€HBP study. <i>Journal of Diabetes Investigation</i> , 2021, 12, 837-844.	2.4	4
133	Visceral adipose tissue quality was associated with nonalcoholic fatty liver disease, independent of its quantity. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2022, 32, 973-980.	2.6	4
134	Let-7e-5p Regulates IGF2BP2, and Induces Muscle Atrophy. <i>Frontiers in Endocrinology</i> , 2021, 12, 791363.	3.5	4
135	Protein intake is not associated with progression of diabetic kidney disease in patients without macroalbuminuria. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3150.	4.0	3
136	Association of mean corpuscular volume with sarcopenia and visceral obesity in individuals without anemia. <i>Journal of Diabetes Investigation</i> , 2020, 12, 1287-1292.	2.4	3
137	Effect of probiotics, <i>Bifidobacterium bifidum</i>, G9-1, on gastrointestinal symptoms in patients with type 2 diabetes mellitus: study protocol for open-label, single-arm, exploratory research trial (Big STAR study). <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020, 67, 223-227.	1.4	3
138	Effect of Teriparatide on Bone Mineral Density and Trabecular Bone Score in Type 2 Diabetic Patients with Osteoporosis: A Retrospective Cohort Study. <i>Medicina (Lithuania)</i> , 2022, 58, 481.	2.0	3
139	Late-night-dinner deteriorates postprandial glucose and insulin whereas consuming dinner dividedly ameliorates them in patients with type 2 diabetes: A randomized crossover clinical trial. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2020, 29, 68-76.	0.4	3
140	Tomato juice preload has a significant impact on postprandial glucose concentration in healthy women: A randomized cross-over trial. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2020, 29, 491-497.	0.4	3
141	Impact of Dietitian-Led Nutrition Therapy of Food Order on 5-Year Glycemic Control in Outpatients with Type 2 Diabetes at Primary Care Clinic: Retrospective Cohort Study. <i>Nutrients</i> , 2022, 14, 2865.	4.1	3
142	A case of diabetic ketoacidosis complicated by fatal acute abdominal aortic thrombosis. <i>Diabetology International</i> , 2013, 4, 201-204.	1.4	2
143	Evaluation of the efficacy of simplified nutritional instructions from physicians on dietary salt restriction for patients with type 2 diabetes mellitus consuming excessive salt: protocol for a randomized controlled trial. <i>Trials</i> , 2019, 20, 761.	1.6	2
144	<p>Usefulness of Exercise for Home Blood Pressure Control in People with Diabetes: A Study Protocol for a Crossover Randomized Controlled Trial</p>. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020, Volume 13, 4747-4753.	2.4	2

#	ARTICLE	IF	CITATIONS
145	Limited joint mobility of the hand correlates incident hospitalisation with infection in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2020, 161, 108049.	2.8	2
146	A Pilot Study on the Effect of Anti-Thrombopoietin Antibody on Platelet Count in Patients with Type 2 Diabetes. <i>Molecules</i> , 2020, 25, 1667.	3.8	2
147	Status of online diet management program users in Japan during the 2020 Coronavirus disease 2019 pandemic. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021, 69, 305-310.	1.4	2
148	Low circulating arachidonic acid is associated with macroalbuminuria in diabetic patients: a cross-sectional examination of the KAMOGAWA-DM cohort study. <i>BMC Nephrology</i> , 2021, 22, 68.	1.8	2
149	Relationship between eosinophils counts and muscle mass decline in older people with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Experimental Gerontology</i> , 2022, 159, 111671.	2.8	2
150	Relative low muscle mass and muscle strength is associated with the prevalence of metabolic syndrome in patients with type 2 diabetes. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2022, 71, 136-142.	1.4	2
151	A survey on consciousness towards the proper use of metformin and medical cost in Japanese patients with type 2 diabetes. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021, 69, 286-293.	1.4	1
152	Changes in the Size of a Ruptured Pheochromocytoma after Transcatheter Arterial Embolization. <i>Case Reports in Medicine</i> , 2021, 2021, 1-5.	0.7	1
153	Correlation between Liver Stiffness by Two-Dimensional Shear Wave Elastography and Waist Circumference in Japanese Local Citizens with Abdominal Obesity. <i>Journal of Clinical Medicine</i> , 2021, 10, 1971.	2.4	1
154	Randomized Controlled Trial of Simple Salt Reduction Instructions by Physician for Patients with Type 2 Diabetes Consuming Excessive Salt. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 6913.	2.6	1
155	Association of Estimated Salt and Miso Intake with the Prevalence of Obesity in People with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021, 13, 3014.	4.1	1
156	752-P: Japanese Radio Calisthenics Prevents Reduction of Skeletal Muscle Volume in Patients with Type 2 Diabetes. <i>Diabetes</i> , 2019, 68, .	0.6	1
157	774-P: Not Carbohydrate but Sucrose Intake Is Associated with Gut Dysbiosis in Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2019, 68, .	0.6	1
158	Characterization of Peripheral Blood TCR in Patients with Type 1 Diabetes Mellitus by BD Rhapsody™ VDJ CDR3 Assay. <i>Cells</i> , 2022, 11, 1623.	4.1	1
159	Association between the frequency of toothbrushing and lifestyle in people with type 2 diabetes mellitus: at the baseline date of the Kamogawa-DM cohort study. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021, 69, 294-298.	1.4	0
160	Home-Measured Blood Pressure Is Associated with Handgrip Strength in Patients with Type 2 Diabetes: The KAMOGAWA-HBP Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 1913.	2.4	0
161	1182-P: Trans Fatty Acid Intake Induces Impaired Glucose Tolerance via Dysbiosis. <i>Diabetes</i> , 2021, 70, 1182-P.	0.6	0
162	801-P: SGLT2 Inhibitor, Luseogliflozin, Prevent Sarcopenia by Improving Extracellular Lipidome. <i>Diabetes</i> , 2021, 70, .	0.6	0

#	ARTICLE	IF	CITATIONS
163	1937-P: miR-23b-3p Upregulates Glucose Uptake in C2C12 Myotube Cells by Regulating the Expression of PTEN. <i>Diabetes</i> , 2019, 68, 1937-P.	0.6	0
164	1943-P: The Sodium-Glucose Cotransporter 2 Inhibitor Luseogliflozin Can Suppress Muscle Atrophy in Db/Db Mice by Suppressing the Expression of Foxo1. <i>Diabetes</i> , 2019, 68, .	0.6	0
165	510-P: Serum Mac-2 Binding Protein Is Associated with Diabetic Micro- and Macroangiopathy in People with Type 2 Diabetes. <i>Diabetes</i> , 2020, 69, .	0.6	0
166	698-P: Moderate-to-Vigorous Physical Activity Associated with Microalbuminuria in Type 2 Diabetes: A Cross-Sectional Study. <i>Diabetes</i> , 2020, 69, 698-P.	0.6	0
167	463-P: Deterioration of Atherosclerosis via Dysbiosis in ApoE Null Mice Fed with High-Salt Diet. <i>Diabetes</i> , 2020, 69, 463-P.	0.6	0
168	Usefulness of Aerobic Exercise for Home Blood Pressure Control in Patients with Diabetes: Randomized Crossover Trial. <i>Journal of Clinical Medicine</i> , 2022, 11, 650.	2.4	0
169	Usefulness of Exercise for Home Blood Pressure Control in People with Diabetes: A Study Protocol for a Crossover Randomized Controlled Trial [Corrigendum]. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 0, Volume 15, 1525-1526.	2.4	0