List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6623706/publications.pdf Version: 2024-02-01



FLIZARETH C. FAKINI

#	Article	IF	CITATIONS
1	Adults' Sedentary Behavior. American Journal of Preventive Medicine, 2011, 41, 189-196.	3.0	691
2	Website-Delivered Physical Activity Interventions. American Journal of Preventive Medicine, 2007, 33, 54-64.	3.0	434
3	Quality of Life and Associated Characteristics in a Large National Sample of Adults With Diabetes. Diabetes Care, 1997, 20, 562-567.	8.6	377
4	Validation of a New Dyspnea Measure. Chest, 1998, 113, 619-624.	0.8	367
5	Systematic review of maintenance of behavior change following physical activity and dietary interventions Health Psychology, 2011, 30, 99-109.	1.6	332
6	Sit–Stand Workstations. American Journal of Preventive Medicine, 2012, 43, 298-303.	3.0	318
7	The Diabetes Network Internet-Based Physical Activity Intervention: A randomized pilot study. Diabetes Care, 2001, 24, 1328-1334.	8.6	311
8	Telephone Interventions for Physical Activity and Dietary Behavior Change. American Journal of Preventive Medicine, 2007, 32, 419-434.	3.0	309
9	Lymphedema after gynecological cancer treatment. Cancer, 2007, 109, 2607-2614.	4.1	303
10	Reducing sitting time in office workers: Short-term efficacy of a multicomponent intervention. Preventive Medicine, 2013, 57, 43-48.	3.4	286
11	Reducing occupational sedentary time: a systematic review and metaâ€analysis of evidence on activityâ€permissive workstations. Obesity Reviews, 2014, 15, 822-838.	6.5	254
12	Clinical Oncology Society of Australia position statement on exercise in cancer care. Medical Journal of Australia, 2018, 209, 184-187.	1.7	254
13	Physician advice and support for physical activity. American Journal of Preventive Medicine, 2001, 21, 189-196.	3.0	233
14	Telephone-Delivered Interventions for Physical Activity and Dietary Behavior Change. American Journal of Preventive Medicine, 2012, 42, 81-88.	3.0	225
15	A Cluster Randomized Controlled Trial to Reduce Office Workers' Sitting Time. Medicine and Science in Sports and Exercise, 2016, 48, 1787-1797.	0.4	219
16	Feasibility of Reducing Older Adults' Sedentary Time. American Journal of Preventive Medicine, 2011, 41, 174-177.	3.0	213
17	Awareness and barriers to use of cancer support and information resources by HMO patients with breast, prostate, or colon cancer: patient and provider perspectives. Psycho-Oncology, 2001, 10, 103-113.	2.3	198
18	Occupational Sitting Time and Overweight and Obesity in Australian Workers. American Journal of Preventive Medicine, 2005, 29, 91-97.	3.0	193

#	Article	IF	CITATIONS
19	Objectively measured physical activity and sedentary time of breast cancer survivors, and associations with adiposity: findings from NHANES (2003–2006). Cancer Causes and Control, 2010, 21, 283-288.	1.8	192
20	Workplace Sitting and Height-Adjustable Workstations. American Journal of Preventive Medicine, 2014, 46, 30-40.	3.0	187
21	A social-ecologic approach to assessing support for disease self-management: the Chronic Illness Resources Survey. Journal of Behavioral Medicine, 2000, 23, 559-583.	2.1	181
22	Implementation and scale up of population physical activity interventions for clinical and community settings: the PRACTIS guide. International Journal of Behavioral Nutrition and Physical Activity, 2018, 15, 51.	4.6	177
23	Unmet supportive care needs and interest in services among patients with a brain tumour and their carers. Patient Education and Counseling, 2008, 71, 251-258.	2.2	167
24	Health behaviors of cancer survivors: data from an Australian population-based survey. Cancer Causes and Control, 2007, 18, 881-894.	1.8	164
25	Population-based interventions for the prevention of fall-related injuries in older people. The Cochrane Library, 2005, , CD004441.	2.8	154
26	Quality of life among patients with a brain tumor and their carers. Journal of Psychosomatic Research, 2007, 63, 617-623.	2.6	153
27	Exercise for health: a randomized, controlled trial evaluating the impact of a pragmatic, translational exercise intervention on the quality of life, function and treatment-related side effects following breast cancer. Breast Cancer Research and Treatment, 2013, 137, 175-186.	2.5	150
28	Measuring Older Adults' Sedentary Time. Medicine and Science in Sports and Exercise, 2011, 43, 2127-2133.	0.4	143
29	Telephone, print, and Web-based interventions for physical activity, diet, and weight control among cancer survivors: a systematic review. Journal of Cancer Survivorship, 2015, 9, 660-682.	2.9	143
30	Associations Between Television Viewing Time and Overall Sitting Time with the Metabolic Syndrome in Older Men and Women: The Australian Diabetes Obesity and Lifestyle Study. Journal of the American Geriatrics Society, 2011, 59, 788-796.	2.6	142
31	Health behavior changes after a cancer diagnosis: What do we know and where do we go from here?. Annals of Behavioral Medicine, 2000, 22, 38-52.	2.9	141
32	Treatment of Dyspnea in COPD. Chest, 1995, 107, 724-729.	0.8	140
33	Unmet needs of gynaecological cancer survivors: implications for developing community support services. Psycho-Oncology, 2008, 17, 392-400.	2.3	139
34	Supportive care needs of people with brain tumours and their carers. Supportive Care in Cancer, 2006, 14, 1094-1103.	2.2	131
35	Weight loss intervention trials in women with breast cancer: a systematic review. Obesity Reviews, 2014, 15, 749-768.	6.5	131
36	Physical activity and/or dietary interventions in breast cancer survivors: a systematic review of the maintenance of outcomes. Journal of Cancer Survivorship, 2013, 7, 74-82.	2.9	123

#	Article	IF	CITATIONS
37	Telephone Counseling for Physical Activity and Diet in Primary Care Patients. American Journal of Preventive Medicine, 2009, 36, 142-149.	3.0	119
38	Older Persons' Perception of Risk of Falling: Implications for Fall-Prevention Campaigns. American Journal of Public Health, 2008, 98, 351-357.	2.7	116
39	Gender differences in prevalence of the metabolic syndrome in Gulf Cooperation Council Countries: a systematic review. Diabetic Medicine, 2010, 27, 593-597.	2.3	115
40	Reaching those most in need: a review of diabetes self-management interventions in disadvantaged populations. Diabetes/Metabolism Research and Reviews, 2002, 18, 26-35.	4.0	112
41	Reducing office workers' sitting time: rationale and study design for the Stand Up Victoria cluster randomized trial. BMC Public Health, 2013, 13, 1057.	2.9	111
42	Weight management and physical activity throughout the cancer care continuum. Ca-A Cancer Journal for Clinicians, 2018, 68, 64-89.	329.8	109
43	10,000 Steps Rockhampton: Evaluation of a Whole Community Approach to Improving Population Levels of Physical Activity. Journal of Physical Activity and Health, 2006, 3, 1-14.	2.0	104
44	Evidence of physical activity participation among men and women in the countries of the Gulf Cooperation Council: a review. Obesity Reviews, 2010, 11, 457-464.	6.5	104
45	Associations of objectively assessed physical activity and sedentary time with biomarkers of breast cancer risk in postmenopausal women: findings from NHANES (2003–2006). Breast Cancer Research and Treatment, 2011, 130, 183-194.	2.5	103
46	Sedentary Behavior and Public Health: Integrating the Evidence and Identifying Potential Solutions. Annual Review of Public Health, 2020, 41, 265-287.	17.4	103
47	A Cluster RCT to Reduce Workers' Sitting Time. Medicine and Science in Sports and Exercise, 2017, 49, 2032-2039.	0.4	101
48	Non-Cancer Mortality among People Diagnosed with Cancer (Australia). Cancer Causes and Control, 2006, 17, 287-297.	1.8	94
49	Accelerometer-Derived Sedentary and Physical Activity Time in Overweight/Obese Adults with Type 2 Diabetes: Cross-Sectional Associations with Cardiometabolic Biomarkers. PLoS ONE, 2015, 10, e0119140.	2.5	94
50	Self-efficacy expectations predict survival for patients with chronic obstructive pulmonary disease Health Psychology, 1994, 13, 366-368.	1.6	90
51	Who Participates in Physical Activity Intervention Trials?. Journal of Physical Activity and Health, 2011, 8, 85-103.	2.0	88
52	Iterative development of Stand Up Australia: a multi-component intervention to reduce workplace sitting. International Journal of Behavioral Nutrition and Physical Activity, 2014, 11, 21.	4.6	87
53	Recruitment and retention of Latinos in a primary care-based physical activity and diet trial: The Resources for Health study. Health Education Research, 2006, 22, 361-371.	1.9	85
54	A Randomized Trial of a Telephone-Delivered Exercise Intervention for Non-urban Dwelling Women Newly Diagnosed with Breast Cancer: Exercise for Health. Annals of Behavioral Medicine, 2012, 43, 229-238.	2.9	84

#	Article	IF	CITATIONS
55	Office workers' objectively assessed total and prolonged sitting time: Individual-level correlates and worksite variations. Preventive Medicine Reports, 2016, 4, 184-191.	1.8	84
56	Health Status of Long-term Cancer Survivors: Results from an Australian Population-Based Sample. Cancer Epidemiology Biomarkers and Prevention, 2006, 15, 1969-1976.	2.5	77
57	Effectiveness of Australia's Get Healthy Information and Coaching Service®: Translational research with population wide impact. Preventive Medicine, 2012, 55, 292-298.	3.4	76
58	Reliability and validity of dyspnea measures in patients with obstructive lung disease. International Journal of Behavioral Medicine, 1995, 2, 118-134.	1.7	74
59	Efficacy of a Text Message-Delivered Extended Contact Intervention on Maintenance of Weight Loss, Physical Activity, and Dietary Behavior Change. JMIR MHealth and UHealth, 2015, 3, e88.	3.7	73
60	Cost-Effectiveness of a Telephone-Delivered Intervention for Physical Activity and Diet. PLoS ONE, 2009, 4, e7135.	2.5	72
61	Randomized trial of a neighborhood environment-focused physical activity website intervention. Preventive Medicine, 2009, 48, 144-150.	3.4	71
62	Objectively assessed physical activity, sedentary time and waist circumference among prostate cancer survivors: findings from the National Health and Nutrition Examination Survey (2003-2006). European Journal of Cancer Care, 2011, 20, 514-519.	1.5	67
63	Living Well With Diabetes: 24-Month Outcomes From a Randomized Trial of Telephone-Delivered Weight Loss and Physical Activity Intervention to Improve Glycemic Control. Diabetes Care, 2014, 37, 2177-2185.	8.6	67
64	Objectively Measured Activity Patterns among Adults in Residential Aged Care. International Journal of Environmental Research and Public Health, 2013, 10, 6783-6798.	2.6	65
65	Gynecological cancer survivors' health behaviors and their associations with quality of life. Cancer Causes and Control, 2008, 19, 775-782.	1.8	64
66	Control Group Improvements in Physical Activity Intervention Trials and Possible Explanatory Factors: A Systematic Review. Journal of Physical Activity and Health, 2012, 9, 884-895.	2.0	64
67	Effectiveness of lifestyleâ€based weight loss interventions for adults with type 2 diabetes: a systematic review and metaâ€analysis. Diabetes, Obesity and Metabolism, 2015, 17, 371-378.	4.4	64
68	Resources for health: A primary-care-based diet and physical activity intervention targeting urban Latinos with multiple chronic conditions Health Psychology, 2007, 26, 392-400.	1.6	60
69	Cardiometabolic Impact of Changing Sitting, Standing, and Stepping in the Workplace. Medicine and Science in Sports and Exercise, 2018, 50, 516-524.	0.4	60
70	The Logan Healthy Living Program: A cluster randomized trial of a telephone-delivered physical activity and dietary behavior intervention for primary care patients with type 2 diabetes or hypertension from a socially disadvantaged community — Rationale, design and recruitment. Contemporary Clinical Trials, 2008, 29, 439-454.	1.8	56
71	Sensitivity to Change of Objectively-Derived Measures of Sedentary Behavior. Measurement in Physical Education and Exercise Science, 2015, 19, 138-147.	1.8	56
72	Evaluating the Population Health Impact of Physical Activity Interventions in Primary Care—Are We Asking the Right Questions?. Journal of Physical Activity and Health, 2005, 2, 197-215.	2.0	55

#	Article	IF	CITATIONS
73	Safety, feasibility and effects of an individualised walking intervention for women undergoing chemotherapy for ovarian cancer: a pilot study. BMC Cancer, 2011, 11, 389.	2.6	53
74	Measurement of dyspnoea in chronic obstructive pulmonary disease. Quality of Life Research, 1993, 2, 181-191.	3.1	48
75	10,000 Steps Rockhampton: Establishing a multi-strategy physical activity promotion project in a community. Health Promotion Journal of Australia, 2003, 14, 95-100.	1.2	48
76	Reducing occupational sitting: Workers' perspectives on participation in a multi-component intervention. International Journal of Behavioral Nutrition and Physical Activity, 2017, 14, 73.	4.6	48
77	Physical activity promotion in primary care. American Journal of Preventive Medicine, 2004, 27, 297-303.	3.0	47
78	Sun exposure and sun protection behaviours among young adult sport competitors. Australian and New Zealand Journal of Public Health, 2007, 31, 230-234.	1.8	46
79	Measuring Physical Activity Change in Broad-Reach Intervention Trials. Journal of Physical Activity and Health, 2010, 7, 194-202.	2.0	46
80	Living Well with Diabetes: a randomized controlled trial of a telephone-delivered intervention for maintenance of weight loss, physical activity and glycaemic control in adults with type 2 diabetes. BMC Public Health, 2010, 10, 452.	2.9	46
81	Follow-up care after breast cancer treatment: experiences and perceptions of service provision and provider interactions in rural Australian women. Supportive Care in Cancer, 2011, 19, 1975-1982.	2.2	45
82	General Practitioner Advice on Physical Activity—Who Gets it?. American Journal of Health Promotion, 2007, 21, 225-228.	1.7	41
83	Active adults recall their physical activity differently to less active adults: test–retest reliability and validity of a physical activity survey. Health Promotion Journal of Australia, 2013, 24, 26-31.	1.2	41
84	Intervening to reduce workplace sitting time: how and when do changes to sitting time occur?. British Journal of Sports Medicine, 2014, 48, 1037-1042.	6.7	41
85	How Generalizable are the Results of Diabetes Self-Management Research? The Impact of Participation and Attrition. The Diabetes Educator, 1996, 22, 573-585.	2.5	40
86	Concern about weight gain associated with quitting smoking: Prevalence and association with outcome in a sample of young female smokers Journal of Consulting and Clinical Psychology, 1999, 67, 1009-1011.	2.0	39
87	Healthy Living after Cancer: a dissemination and implementation study evaluating a telephone-delivered healthy lifestyle program for cancer survivors. BMC Cancer, 2015, 15, 992.	2.6	39
88	The Living Well after Breast Cancerâ,,¢ Pilot Trial: a weight loss intervention for women following treatment for breast cancer. Asia-Pacific Journal of Clinical Oncology, 2017, 13, 125-136.	1.1	39
89	Medical Office-based Interventions. , 0, , 141-168.		38
90	The Queensland Cancer Risk Study: behavioural risk factor results. Australian and New Zealand Journal of Public Health, 2006, 30, 375-382.	1.8	37

#	Article	IF	CITATIONS
91	Six-Month Outcomes from Living Well with Diabetes: A Randomized Trial of a Telephone-Delivered Weight Loss and Physical Activity Intervention to Improve Glycemic Control. Annals of Behavioral Medicine, 2013, 46, 193-203.	2.9	37
92	The inside scoop on the stuff called snuff: An interview study of 94 adult male smokeless tobacco users. Journal of Substance Abuse, 1990, 2, 77-85.	1.1	36
93	Evaluating the Maintenance of Lifestyle Changes in a Randomized Controlled Trial of the â€~Get Healthy, Stay Healthy' Program. JMIR MHealth and UHealth, 2016, 4, e42.	3.7	36
94	Maintenance of physical activity and dietary change following a telephone-delivered intervention Health Psychology, 2010, 29, 566-573.	1.6	34
95	Prevalence and correlates of multiple cancer risk behaviors in an Australian population-based survey: results from the Queensland Cancer Risk Study. Cancer Causes and Control, 2008, 19, 1339-1347.	1.8	33
96	Design and implementation of the Exercise for Health trial — A pragmatic exercise intervention for women with breast cancer. Contemporary Clinical Trials, 2011, 32, 577-585.	1.8	32
97	Associations of Physical Activity and Sitting Time With the Metabolic Syndrome Among Omani Adults. Obesity, 2012, 20, 2290-2295.	3.0	32
98	Effectiveness of Australia's Get Healthy Information and Coaching Service®: maintenance of self-reported anthropometric and behavioural changes after program completion. BMC Public Health, 2013, 13, 175.	2.9	32
99	Feasibility, acceptability and efficacy of a text message-enhanced clinical exercise rehabilitation intervention for increasing â€ĩwhole-of-day' activity in people living with and beyond cancer. BMC Public Health, 2019, 19, 542.	2.9	32
100	Cost-effectiveness of a pragmatic exercise intervention for women with breast cancer: results from a randomized controlled trial. Psycho-Oncology, 2017, 26, 649-655.	2.3	31
101	Resources for Health: A Social-Ecological Intervention for Supporting Self-management of Chronic Conditions. Journal of Health Psychology, 2001, 6, 693-705.	2.3	30
102	Evaluation Framework for Translational Research. Health Promotion Practice, 2013, 14, 380-389.	1.6	30
103	Correlates of Omani adults' physical inactivity and sitting time. Public Health Nutrition, 2013, 16, 65-72.	2.2	30
104	Economic evaluation of a randomized controlled trial of an intervention to reduce office workers' sitting time: the "Stand Up Victoria" trial. Scandinavian Journal of Work, Environment and Health, 2018, 44, 503-511.	3.4	30
105	Solaria use in Queensland, Australia. Australian and New Zealand Journal of Public Health, 2006, 30, 479-482.	1.8	29
106	Intervening to reduce workplace sitting: mediating role of social-cognitive constructs during a cluster randomised controlled trial. International Journal of Behavioral Nutrition and Physical Activity, 2017, 14, 27.	4.6	29
107	Program sustainability of a community-based intervention to prevent falls among older Australians. Health Promotion International, 2004, 19, 281-288.	1.8	28
108	Associations of Monitor-Assessed Activity with Performance-Based Physical Function. PLoS ONE, 2016, 11, e0153398.	2.5	28

#	Article	IF	CITATIONS
109	Evaluation of the efficacy of 3D total-body photography with sequential digital dermoscopy in a high-risk melanoma cohort: protocol for a randomised controlled trial. BMJ Open, 2019, 9, e032969.	1.9	27
110	A Telephone-Delivered Physical Activity and Dietary Intervention for Type 2 Diabetes and Hypertension: Does Intervention Dose Influence Outcomes?. American Journal of Health Promotion, 2011, 25, 257-263.	1.7	26
111	Addressing physical inactivity in Omani adults: perceptions of public health managers. Public Health Nutrition, 2014, 17, 674-681.	2.2	26
112	Multi-level support for physical activity and healthy eating. Journal of Advanced Nursing, 2006, 54, 585-593.	3.3	25
113	The BeUpstanding Program TM : Scaling up the Stand Up Australia Workplace Intervention for Translation into Practice. AIMS Public Health, 2016, 3, 341-347.	2.6	24
114	Measuring physical activity change in broad-reach intervention trials. Journal of Physical Activity and Health, 2010, 7, 194-202.	2.0	24
115	Gynecological cancer survivors and community support services: referral, awareness, utilization and satisfaction. Psycho-Oncology, 2010, 19, 54-61.	2.3	23
116	Fat and fibre behaviour questionnaire: Reliability, relative validity and responsiveness to change in A ustralian adults with type 2 diabetes and/or hypertension. Nutrition and Dietetics, 2015, 72, 368-376.	1.8	23
117	Relationship between Intervention Dose and Outcomes in Living Well with Diabetes—A Randomized Trial of a Telephone-Delivered Lifestyle-Based Weight Loss Intervention. American Journal of Health Promotion, 2015, 30, 120-129.	1.7	23
118	Women's Perceptions of Participation in an Extended Contact Text Message–Based Weight Loss Intervention: An Explorative Study. JMIR MHealth and UHealth, 2017, 5, e21.	3.7	22
119	Exploring the feasibility and acceptability of using Internet technology to promote physical activity within a defined community. Health Promotion Journal of Australia, 2005, 16, 82-84.	1.2	21
120	Multiple Health Behavior Changes and Co-variation in a Telephone Counseling Trial. Annals of Behavioral Medicine, 2010, 39, 250-257.	2.9	21
121	Breast cancer survivors' experience of making weight, dietary and physical activity changes during participation in a weight loss intervention. Supportive Care in Cancer, 2017, 25, 1455-1463.	2.2	21
122	The Patients' Perspective on the Self-management of Chronic Obstructive Pulmonary Disease. Journal of Health Psychology, 1997, 2, 245-253.	2.3	20
123	Validation of the spanish-language version of the chronic Illness resources survey. International Journal of Behavioral Medicine, 2007, 14, 76-85.	1.7	19
124	Living well after breast cancer randomized controlled trial protocol: evaluating a telephone-delivered weight loss intervention versus usual care in women following treatment for breast cancer. BMC Cancer, 2016, 16, 830.	2.6	19
125	Use of tailored videos in primary care smoking cessation interventions. Health Education Research, 1998, 13, 519-527.	1.9	18
126	Promoting Physical Activity among Middle-Aged and Older Adults in Health Care Settings. Journal of Aging and Physical Activity, 2001, 9, S29-S37.	1.0	18

8

#	Article	IF	CITATIONS
127	Correlates of pedometer use: Results from a community-based physical activity intervention trial (10,000 Steps Rockhampton). International Journal of Behavioral Nutrition and Physical Activity, 2007, 4, 31.	4.6	18
128	A cluster randomized controlled trial to reduce office workers' sitting time: effect on productivity outcomes. Scandinavian Journal of Work, Environment and Health, 2019, 45, 483-492.	3.4	17
129	Responsiveness to Change of Self-Report and Device-Based Physical Activity Measures in the Living Well With Diabetes Trial. Journal of Physical Activity and Health, 2015, 12, 1082-1087.	2.0	16
130	Twelve-Year Television Viewing Time Trajectories and Physical Function in Older Adults. Medicine and Science in Sports and Exercise, 2017, 49, 1359-1365.	0.4	16
131	What strategies do desk-based workers choose to reduce sitting time and how well do they work? Findings from a cluster randomised controlled trial. International Journal of Behavioral Nutrition and Physical Activity, 2018, 15, 98.	4.6	16
132	Effect of a Remotely Delivered Weight Loss Intervention in Early-Stage Breast Cancer: Randomized Controlled Trial. Nutrients, 2021, 13, 4091.	4.1	16
133	Cost-effectiveness analyses and modelling the lifetime costs and benefits of health-behaviour interventions. Chronic Illness, 2006, 2, 97-107.	1.5	15
134	Translation from Research to Practice: Community Dissemination of a Telephone-Delivered Physical Activity and Dietary Behavior Change Intervention. American Journal of Health Promotion, 2012, 26, 253-259.	1.7	15
135	Supporting Workers to Sit Less and Move More Through the Web-Based BeUpstanding Program: Protocol for a Single-Arm, Repeated Measures Implementation Study. JMIR Research Protocols, 2020, 9, e15756.	1.0	15
136	Feasibility, effectiveness and cost-effectiveness of a telephone-based weight loss program delivered via a hospital outpatient setting. Translational Behavioral Medicine, 2016, 6, 386-395.	2.4	14
137	Results from the dissemination of an evidence-based telephone-delivered intervention for healthy lifestyle and weight loss: the Optimal Health Program. Translational Behavioral Medicine, 2013, 3, 340-350.	2.4	13
138	Moderators of health behavior initiation and maintenance in a randomized telephone counseling trial. Preventive Medicine, 2014, 61, 34-41.	3.4	13
139	Precision Public Health for Non-communicable Diseases: An Emerging Strategic Roadmap and Multinational Use Cases. Frontiers in Public Health, 2022, 10, 854525.	2.7	13
140	Physician's Role in Diabetes Self-Management. , 1996, 6, 186-195.		12
141	Patients' self-reports of dyspnea: An important and independent outcome in chronic obstructive pulmonary disease. Annals of Behavioral Medicine, 1996, 18, 87-90.	2.9	12
142	Randomised controlled trial of a supervised exercise rehabilitation program for colorectal cancer survivors immediately after chemotherapy: study protocol. BMC Cancer, 2007, 7, 154.	2.6	12
143	Temporal features of sitting, standing and stepping changes in a cluster-randomised controlled trial of a workplace sitting-reduction intervention. International Journal of Behavioral Nutrition and Physical Activity, 2019, 16, 111.	4.6	12
144	Recruitment of Managed Care Medicare Patients for a Physical Activity Study. American Journal of Health Promotion, 1997, 12, 98-101.	1.7	11

#	Article	IF	CITATIONS
145	Relationships of Sun-Protection Habit Strength with Sunscreen Use During Outdoor Sport and Physical Activity. International Journal of Environmental Research and Public Health, 2012, 9, 916-923.	2.6	11
146	Assessing the Feasibility and Pre-Post Impact Evaluation of the Beta (Test) Version of the BeUpstanding Champion Toolkit in Reducing Workplace Sitting: Pilot Study. JMIR Formative Research, 2018, 2, e17.	1.4	11
147	Characteristics of control group participants who increased their physical activity in a cluster-randomized lifestyle intervention trial. BMC Public Health, 2011, 11, 27.	2.9	10
148	Is Measurement Error Altered by Participation in a Physical Activity Intervention?. Medicine and Science in Sports and Exercise, 2013, 45, 1004-1011.	0.4	10
149	â€~Get Healthy, Stay Healthy': protocol for evaluation of a lifestyle intervention delivered by text-message following the Get Healthy Information and Coaching Service®. BMC Public Health, 2014, 14, 112.	2.9	10
150	Translating research into practice: outcomes from the Healthy Living after Cancer partnership project. BMC Cancer, 2020, 20, 963.	2.6	10
151	Reaching those most in need: Participation in a Planned Parenthood smoking cessation program. Annals of Behavioral Medicine, 1998, 20, 216-220.	2.9	9
152	Falls prevention in rural general practice: what stands the test of time and where to from here?. Australian and New Zealand Journal of Public Health, 2003, 27, 481-485.	1.8	9
153	Dissemination of an evidence-based telephone-delivered lifestyle intervention: factors associated with successful implementation and evaluation. Translational Behavioral Medicine, 2013, 3, 351-356.	2.4	9
154	A National Strategy for Promoting Physical Activity in Oman: A call for action. Sultan Qaboos University Medical Journal, 2014, 14, e170-5.	1.0	9
155	Design of a randomized controlled trial for multiple cancer risk behaviors among Spanish-speaking Mexican-origin smokers. BMC Public Health, 2013, 13, 237.	2.9	8
156	Common mental disorders and recent physical activity status: findings from a National Community Survey. Social Psychiatry and Psychiatric Epidemiology, 2017, 52, 795-802.	3.1	8
157	Exercise as part of routine cancer care. Lancet Oncology, The, 2018, 19, e432.	10.7	8
158	Get Healthy, Stay Healthy: Evaluation of the Maintenance of Lifestyle Changes Six Months After an Extended Contact Intervention. JMIR MHealth and UHealth, 2019, 7, e11070.	3.7	8
159	Physical activity is important, but can it be promoted in general practice?. Medical Journal of Australia, 2003, 179, 70-71.	1.7	7
160	What do cancer survivors and their health care providers want from a healthy living program? Results from the first round of a co-design project. Supportive Care in Cancer, 2021, 29, 4847-4858.	2.2	7
161	Using pedometers to increase physical activity in a family planning clinic: a feasibility study. Health Promotion Journal of Australia, 2003, 14, 165-170.	1.2	6
162	Prevalence and determinants of sunburn in Queensland. Health Promotion Journal of Australia, 2009, 20, 102-106.	1.2	6

#	Article	IF	CITATIONS
163	The impact of behavioural screening on intervention outcomes in a randomised, controlled multiple behaviour intervention trial. International Journal of Behavioral Nutrition and Physical Activity, 2011, 8, 24.	4.6	6
164	Adaptation of a Counseling Intervention to Address Multiple Cancer Risk Factors Among Overweight/Obese Latino Smokers. Health Education and Behavior, 2015, 42, 65-72.	2.5	6
165	Cost-Effectiveness Analysis from a Randomized Controlled Trial of Tailored Exercise Prescription for Women with Breast Cancer with 8-Year Follow-Up. International Journal of Environmental Research and Public Health, 2020, 17, 8608.	2.6	6
166	A Randomised, Comparative, Effectiveness Trial Evaluating Low- versus High-Level Supervision of an Exercise Intervention for Women with Breast Cancer: The SAFE Trial. Cancers, 2022, 14, 1528.	3.7	6
167	Designing for the Dissemination of Environmental and Policy Initiatives and Programs for High-Risk Groups. , 2012, , 114-127.		5
168	Dietary and Physical Activity Changes and Adherence to WCRF/AICR Cancer Prevention Recommendations following a Remotely Delivered Weight Loss Intervention for Female Breast Cancer Survivors: The Living Well after Breast Cancer Randomized Controlled Trial. Journal of the Academy of Nutrition and Dietetics, 2022, , .	0.8	5
169	Preventing Falls Among Older People—Current Practice and Attitudes Among Community Pharmacists. Journal of Pharmacy Practice and Research, 2003, 33, 51-54.	0.8	4
170	Telephoneâ€delivered weight management services in the hospital outpatient setting: Decisionâ€makers' perceptions of their use in routine practice. Nutrition and Dietetics, 2017, 74, 261-267.	1.8	4
171	Data sources for precision public health of obesity: a scoping review, evidence map and use case in Queensland, Australia. BMC Public Health, 2022, 22, 584.	2.9	4
172	Complexities and Context of Scaling Up: A Qualitative Study of Stakeholder Perspectives of Scaling Physical Activity and Nutrition Interventions in Australia. Frontiers in Public Health, 2022, 10, 771235.	2.7	4
173	Sitting less and moving more for improved metabolic and brain health in type 2 diabetes: â€ ⁻ OPTIMISE your health' trial protocol. BMC Public Health, 2022, 22, 929.	2.9	4
174	Implementation of the Living Well During Pregnancy Telecoaching Program for Women at High Risk of Excessive Gestational Weight Gain: Protocol for an Effectiveness-Implementation Hybrid Study. JMIR Research Protocols, 2021, 10, e27196.	1.0	3
175	Exercise Behavior, Motivation, and Maintenance Among Cancer Survivors. , 2013, , 215-231.		3
176	Evaluation of the Healthy Living after Cancer text message-delivered, extended contact intervention using the RE-AIM framework. BMC Cancer, 2021, 21, 1081.	2.6	3
177	Cost-effectiveness analyses and modelling the lifetime costs and benefits of health-behaviour interventions. Chronic Illness, 2006, 2, 97-107.	1.5	3
178	Outcomes from a hybrid implementation-effectiveness study of the living well during pregnancy Tele-coaching program for women at high risk of excessive gestational weight gain. BMC Health Services Research, 2022, 22, 589.	2.2	3
179	Correlates of Omani adults' physical inactivity and sitting time – Corrigendum. Public Health Nutrition, 2012, 15, 2164-2164.	2.2	2
180	Translating research into practice: The healthy living after cancer partnership project. Journal of Science and Medicine in Sport, 2014, 18, e13.	1.3	2

#	Article	IF	CITATIONS
181	Relative validity of a brief Fat and Fibre Behaviour Questionnaire in a population of overweight and obese breast cancer survivors: A note of caution. Nutrition and Dietetics, 2017, 74, 18-28.	1.8	2
182	Translating Research into Community Practice: The Healthy Living after Cancer Partnership Project. Obesity, 2017, 25, S31-S31.	3.0	2
183	Depressive symptoms and obesity: Assessing and addressing the black dog in the room. Nutrition and Dietetics, 2012, 69, 234-235.	1.8	1
184	Dose and engagement during an extended contact physical activity and dietary behavior change intervention delivered via tailored text messaging: exploring relationships with behavioral outcomes. International Journal of Behavioral Nutrition and Physical Activity, 2021, 18, 119.	4.6	1
185	Is Not the Treatment Program of Dyspnea Management Strategies Effective for COPD?-To the Editor. Chest, 1995, 108, 1774.	0.8	0