

Robin M Giblin-Davis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6636054/publications.pdf>

Version: 2024-02-01

138
papers

2,874
citations

201674
27
h-index

233421
45
g-index

139
all docs

139
docs citations

139
times ranked

1416
citing authors

#	ARTICLE	IF	CITATIONS
1	Evaluating high-throughput sequencing as a method for metagenomic analysis of nematode diversity. <i>Molecular Ecology Resources</i> , 2009, 9, 1439-1450.	4.8	264
2	Phylogenetic relationships among <i>Bursaphelenchus</i> species (Nematoda: Parasitaphelenchidae) inferred from nuclear ribosomal and mitochondrial DNA sequence data. <i>Molecular Phylogenetics and Evolution</i> , 2007, 43, 1185-1197.	2.7	260
3	Molecular phylogenetics and the evolution of host plant associations in the nematode genus <i>Fergusobia</i> (Tylenchida: Fergusobiinae). <i>Molecular Phylogenetics and Evolution</i> , 2007, 45, 123-141.	2.7	93
4	Reproducibility of read numbers in high-throughput sequencing analysis of nematode community composition and structure. <i>Molecular Ecology Resources</i> , 2010, 10, 666-676.	4.8	93
5	Ecometagenetics confirm high tropical rainforest nematode diversity. <i>Molecular Ecology</i> , 2010, 19, 5521-5530.	3.9	90
6	Large-scale diversification without genetic isolation in nematode symbionts of figs. <i>Science Advances</i> , 2016, 2, e1501031.	10.3	82
7	The biology and associations of <i>Fergusobia</i> (Nematoda) from the Melaleuca leucadendra-complex in eastern Australia. <i>Invertebrate Systematics</i> , 2004, 18, 291.	1.3	57
8	Morphological and molecular analysis of six aphelenchoidoids from Australian conifers and their relationship to <i>Bursaphelenchus</i> (Fuchs, 1937). <i>Nematology</i> , 2008, 10, 663-678.	0.6	47
9	A review of the taxonomy, phylogeny, distribution and co-evolution of <i>Schistonchus</i> Cobb, 1927 with proposal of <i>Ficophagus</i> n. gen. and <i>Martininema</i> n. gen. (Nematoda: Aphelenchoididae). <i>Nematology</i> , 2015, 17, 761-829.	0.6	46
10	Diversity and Plant Pathogenicity of <i>Bursaphelenchus</i> and Related Nematodes in Relation to Their Vector Bionomics. <i>Current Forestry Reports</i> , 2018, 4, 85-100.	7.4	44
11	Optimization of semiochemical-based trapping of <i>Metamasius hemipterus sericeus</i> (Olivier) (Coleoptera: Curculionidae). <i>Journal of Chemical Ecology</i> , 1996, 22, 1389-1410.	1.8	40
12	Linking operational clustered taxonomic units (OCTUs) from parallel ultra sequencing (PUS) to nematode species. <i>Zootaxa</i> , 2010, 2427, .	0.5	40
13	Morphology and systematics of <i>Bursaphelenchus gerberae</i> n. sp. (Nematoda: Parasitaphelenchidae), a rare associate of the palm weevil, <i>Rhynchophorus palmarum</i> in Trinidad. <i>Zootaxa</i> , 2006, 1189, 39-53.	0.5	39
14	Two new species of <i>Schistonchus</i> (Nematoda: Aphelenchoididae) associated with <i>Ficus hispida</i> in China. <i>Nematology</i> , 2007, 9, 169-187.	0.6	39
15	<i>Pseudaphelenchus yukiae</i> n. gen., n. sp. (Tylenchina: Aphelenchoididae) associated with <i>Cylindrotermes macrognathus</i> (Termitidae: Termitinae) in La Selva, Costa Rica. <i>Nematology</i> , 2009, 11, 869-881.	0.6	38
16	New Plant-Parasitic Nematode from the Mostly Mycophagous Genus <i>Bursaphelenchus</i> Discovered inside Figs in Japan. <i>PLoS ONE</i> , 2014, 9, e99241.	2.5	38
17	Identification of Chilean <I>Reticulitermes</I> (Isoptera: Rhinotermitidae) Inferred from Three Mitochondrial Gene DNA Sequences and Soldier Morphology. <i>Annals of the Entomological Society of America</i> , 2006, 99, 352-363.	2.5	37
18	<i>Schistonchus</i> (Aphelenchoididae) from <i>Ficus</i> (Moraceae) in Australia, with description of <i>S. aculeata</i> sp. n.. <i>Nematology</i> , 2010, 12, 935-958.	0.6	37

#	ARTICLE	IF	CITATIONS
19	Description of <i>Bursaphelenchus kiyoharai</i> n. sp. (Tylenchina: Aphelenchoididae) with remarks on the taxonomic framework of the Parasitaphelenchinae RÄ¼hm, 1956 and Aphelenchoidinae Fuchs, 1937. <i>Nematology</i> , 2011, 13, 787-804.	0.6	36
20	Vulnerability of Stressed Palms to Attack by <i>Rhynchophorus cruentatus</i> (Coleoptera: Curculionidae) and Insecticidal Control of the Pest. <i>Journal of Economic Entomology</i> , 1989, 82, 1185-1190.	1.8	35
21	Pheromone chirality of african palm weevil, <i>Rhynchophorus phoenicis</i> (F.) and palmetto weevil, <i>Rhynchophorus cruentatus</i> (F.) (Coleoptera: Curculionidae). <i>Journal of Chemical Ecology</i> , 1994, 20, 2653-2671.	1.8	35
22	The nematode genus <i>Fergusobia</i> (Nematoda: Neotylenchidae): molecular phylogeny, descriptions of clades and associated galls, host plants and Fergusonina fly larvae. <i>Zootaxa</i> , 2010, 2633, 1.	0.5	35
23	<i>Schistonchus virens</i> sp. n. (Aphelenchoididae) and <i>Parasitodiplogaster australis</i> sp. n. (Diplogastridae) from <i>Ficus virens</i> (Moraceae) in Australia. <i>Nematology</i> , 2009, 11, 583-601.	0.6	33
24	Stomatal Ultrastructure, Molecular Phylogeny, and Description of <i>Parasitodiplogaster laevigata</i> n. sp. (Nematoda: Diplogastridae), a Parasite of Fig Wasps. <i>Journal of Nematology</i> , 2006, 38, 137-49.	0.9	31
25	<i>Bursaphelenchus rufipennis</i> n. sp. (Nematoda: Parasitaphelenchinae) and redescription of <i>Ektaphelenchus obtusus</i> (Nematoda: Ektaphelenchinae), associates from nematangia on the hind wings of <i>Dendroctonus rufipennis</i> (Coleoptera: Scolytidae). <i>Nematology</i> , 2008, 10, 925-955.	0.6	30
26	Water Loss Dynamics and Humidity Preference of <i>Rhynchophorus cruentatus</i> (Coleoptera:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 462 To 1.4		
27	< i>Teratodiplogaster fignewmani</i> gen. nov., sp. nov. (Nematoda: Diplogastridae) from the Syconia of < i>Ficus racemose</i> in Australia. <i>Zoological Science</i> , 2009, 26, 569-578.	0.7	29
28	Description of two <i>Pseudaphelenchus</i> species (Tylenchomorpha: Aphelenchoididae) associated with Asian termites and proposal of <i>Tylaphelenchinae</i> n. subfam.. <i>Nematology</i> , 2014, 16, 963-978.	0.6	29
29	The fig wasp followers and colonists of a widely introduced fig tree, < i>Ficus microcarpa</i>. <i>Insect Conservation and Diversity</i> , 2015, 8, 322-336.	3.0	27
30	Description of <i>Ektaphelenchoides spondylis</i> n. sp. (Nematoda: Ektaphelenchinae) isolated from <i>Spondylis buprestoides</i> (Coleoptera: Cerambycidae) in Japan. <i>Nematology</i> , 2009, 11, 181-188.	0.6	26
31	Description of <i>Schistonchus microcarpus</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus microcarpa</i> in China. <i>Nematology</i> , 2011, 13, 221-233.	0.6	26
32	Molecular Characterization and Development of Real-Time PCR Assay for Pine-Wood Nematode <i>Bursaphelenchus xylophilus</i> (Nematoda: Parasitaphelenchidae). <i>PLoS ONE</i> , 2013, 8, e78804.	2.5	24
33	Resistance in <i>Stenotaphrum</i> to the Sting Nematode. <i>Crop Science</i> , 1993, 33, 1066-1070.	1.8	24
34	<i>Schistonchus hirtus</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus hirta</i> in China. <i>Nematology</i> , 2010, 12, 543-556.	0.6	23
35	Description of <i>Caenorhabditis angaria</i> n. sp. (Nematoda: Rhabditidae), an associate of sugarcane and palm weevils (Coleoptera: Curculionidae). <i>Nematology</i> , 2011, 13, 61-78.	0.6	23
36	Laboratory and field evidence for maleproduced aggregation pheromone in <i>Rhynchophorus cruentatus</i> (F.) (Coleoptera: Curculionidae). <i>Journal of Chemical Ecology</i> , 1993, 19, 1195-1203.	1.8	22

#	ARTICLE	IF	CITATIONS
37	Acrostichus rhynchophori n. sp. (Rhabditida: Diplogastridae): a phoretic associate of Rhynchophorus cruentatus Fabricius and R. palmarum L. (Coleoptera: Curculionidae) in the Americas. <i>Nematology</i> , 2009, 11, 669-688.	0.6	21
38	Three new species of Schistonchus (Aphelenchoididae) from the Ficus subgenus Sycomorus (Moraceae) in northern Australia. <i>Nematology</i> , 2013, 15, 347-362.	0.6	21
39	Ficotylus congestae gen. n., sp. n. (Anguinata), from <i>Ficus congesta</i> (Moraceae) sycones in Australia. <i>Nematology</i> , 2009, 11, 63-75.	0.6	20
40	Teratodiplogaster martini n. sp. and Parasitodiplogaster doliosstoma Åsp. (Nematoda: Diplogastridae) from the syconia of <i>Ficus</i> species from Africa. <i>Nematology</i> , 2012, 14, 529-546.	0.6	20
41	Nematodes from galls on Myrtaceae. I. Fergusobia/Fergusonina galls on <i>Corymbia</i> spp., with re-description of <i>F. magna</i> and notes on its phylogenetic relationships. <i>Zootaxa</i> , 2010, 2634, 25.	0.5	19
42	Teratorhabditis synpapillata Sudhaus, 1985 (Rhabditida: Rhabditidae) is an associate of the red palm weevil, <i>Rhynchophorus ferrugineus</i> (Coleoptera: Curculionidae). <i>Nematology</i> , 2008, 10, 207-218.	0.6	18
43	Pseudaphelenchus vindai n. sp. (Tylenchomorpha: Aphelenchoididae) associated with termites (Termitidae) in Barro Colorado Island, Panama. <i>Nematology</i> , 2010, 12, 905-914.	0.6	18
44	Description of Ektaphelenchus berbericus n. sp. (Rhabditida: Ektaphelenchinae) from eastern Iran. <i>Nematology</i> , 2016, 18, 1063-1077.	0.6	18
45	Parasitodiplogaster species associated with <i>Pharmacosycea</i> figs in Panama. <i>Nematology</i> , 2014, 16, 607-619.	0.6	17
46	Acrostichus puri n. sp. (Nematoda: Diplogastridae), a phoretic associate of <i>Augochlora pura mosieri</i> Cockerell (Hymenoptera: Halictidae). <i>Nematology</i> , 2010, 12, 49-64.	0.6	16
47	Schistonchus (Aphelenchoididae) from <i>Ficus benjamina</i> in Australia, with description of <i>S. benjamina</i> sp. n.. <i>Nematology</i> , 2012, 14, 1005-1013.	0.6	16
48	Nematodes from galls on Myrtaceae. II. Fergusobia/Fergusonina from small axillary bud ('stem') and leaf ('pea') galls in Australia, with descriptions of two new species. <i>Zootaxa</i> , 2012, 3415, .	0.5	16
49	An emerging example of tritrophic coevolution between flies (Diptera: Fergusoninidae) and nematodes (Nematoda: Neotylenchidae) on Myrtaceae host plants. <i>Biological Journal of the Linnean Society</i> , 2014, 111, 699-718.	1.6	16
50	Evaluation of an Entomopathogenic Nematode and Chemical Insecticides for Control of <i>Metamasius hemipterus sericeus</i> (Coleoptera: Curculionidae)2. <i>Journal of Entomological Science</i> , 1996, 31, 240-251.	0.3	16
51	Resource Opportunities from the Nest of Dying Subterranean Termite (Isoptera: Rhinotermitidae) Colonies: A Laboratory Case of Ecological Succession. <i>Annals of the Entomological Society of America</i> , 2013, 106, 771-778.	2.5	15
52	<p class="HeadingRunIn">Description of Schistonchus altissimus n. sp. (Nematoda: Aphelenchoididae), an associate of Ficus altissima in China</p>. <i>Zootaxa</i> , 2013, 3700, 561.	0.5	15
53	Feeding Dimorphism in a Mycophagous Nematode, <i>Bursaphelenchus sinensis</i> . <i>Scientific Reports</i> , 2019, 9, 13956.	3.3	15
54	Poikilolaimus floridensis n. sp. (Rhabditida: Rhabditidae) associated with termites (Kalotermitidae). <i>Nematology</i> , 2009, 11, 203-216.	0.6	14

#	ARTICLE	IF	CITATIONS
55	Acrostichus megaloptae n. sp. (Nematoda: Diplogastridae), a phoretic associate of Megalopta spp. (Hymenoptera: Halictidae) in Central America. <i>Nematology</i> , 2010, 12, 453-468.	0.6	14
56	Nematodes from galls on Myrtaceae. III. Fergusobia from flower bud and stigma galls on Eucalyptus, with descriptions of four new species. <i>Zootaxa</i> , 2012, 3532, 1.	0.5	14
57	Schistonchus fistulosus n. sp. (Nematoda: Aphelenchoididae) associated with <i>Ficus fistulosa</i> in China. <i>Nematology</i> , 2013, 15, 879-890.	0.6	14
58	Description of Schistonchus superbus n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus superba</i> in China. <i>Nematology</i> , 2013, 15, 771-781.	0.6	14
59	Description of Parasitodiplogaster pharmaconema n. sp. and redescription of <i>P. maxinema</i> from <i>Ficus maxima</i> Mill. (Moraceae). <i>Nematology</i> , 2013, 15, 957-974.	0.6	14
60	Molecular variability of <i>Schistonchus caprifici</i> (Nematoda: Aphelenchoididae) from <i>Ficus carica</i> in Turkey. <i>Nematology</i> , 2008, 10, 639-649.	0.6	13
61	Bursaphelenchus tokyoensis n. sp. (Nematoda: Parasitaphelenchinae) isolated from dead wood of the Japanese red pine, <i>Pinus densiflora</i> Sieb. & Zucc., in Japan. <i>Nematology</i> , 2009, 11, 171-180.	0.6	13
62	Ditylenchus halictus n. sp. (Nematoda: Anguinidae), an associate of the sweat bee, <i>Halictus sexcinctus</i> (Halictidae), from Germany. <i>Nematology</i> , 2010, 12, 891-904.	0.6	13
63	Schistonchus molochi sp. n. and <i>S. athertonensis</i> sp. n. (Nematoda: Aphelenchoididae) from <i>Ficus watkinsiana</i> (Moraceae) in Australia. <i>Nematology</i> , 2013, 15, 389-400.	0.6	13
64	Revision of the paraphyletic genus <i>Koerneria</i> Meyl, 1960 and resurrection of two other genera of Diplogastridae (Nematoda). <i>ZooKeys</i> , 2014, 442, 17-30.	1.1	13
65	Description of <i>Ruehmaphelenchus juliae</i> n. sp. (Tylenchina: Aphelenchoididae) isolated from an ambrosia beetle, <i>Xylosandrus crassiusculus</i> (Motschulsky), from South Florida. <i>Nematology</i> , 2015, 17, 639-653.	0.6	13
66	A molecular phylogenetic approach for unravelling the taxonomic status of <i>Basilaphelenchus persicus</i> n. gen., n. sp. (Aphelenchoididae: Tyaphelenchinae). <i>Nematology</i> , 2018, 20, 567-582.	0.6	13
67	A new species of <i>Basilaphelenchus</i> Pedram, Kanzaki, Giblin-Davis & Pourjam, 2018 (Aphelenchoidea: Tj ETQq1 1 0.784314 rgBT /Overloo	0.6	13
68	Fecundity and Fertility of <i>Rhynchophorus cruentatus</i> (Coleoptera: Curculionidae). <i>Florida Entomologist</i> , 1994, 77, 373.	0.5	12
69	Flight Behavior and Seasonal Trapping of <i>Rhynchophorus cruentatus</i> (Coleoptera: Curculionidae). <i>Annals of the Entomological Society of America</i> , 1994, 87, 641-647.	2.5	12
70	Diversity and distribution of nematodes associated with wild bees in Turkey. <i>Nematology</i> , 2010, 12, 65-80.	0.6	12
71	<p class="HeadingRunIn">Nematodes from galls on Myrtaceae. IV. Fergusobia from flat leaf galls on Eucalyptus and Corymbia, with descriptions of two new species</p>. <i>Zootaxa</i> , 2013, 3741, 151.	0.5	12
72	Molecular and morphological observations on <i>Parasitodiplogaster sycophilon</i> Poinar, 1979 (Nematoda: Diplogastrina) associated with <i>Ficus burkei</i> in Africa. <i>Nematology</i> , 2014, 16, 453-462.	0.6	12

#	ARTICLE	IF	CITATIONS
73	Redescription of two Panamanian nematodes, <i>Parasitodiplogaster citrinema</i> Poinar & Herre, 1991 and <i>P. popenema</i> Poinar & Herre, 1991 (Nematoda: Diplogastrina). <i>Nematology</i> , 2010, 12, 89-104.	0.6	11
74	Distribution of southern sting nematode, <i>Ibipora lolii</i> (Nematoda: Belonolaimidae), on turfgrass in Australia and its taxonomic relationship to other belonolaimids. <i>Nematology</i> , 2013, 15, 401-415.	0.6	11
75	<p class="HeadingRunIn">Nematodes from galls on Myrtaceae. V. Fergusobia from large multilocular shoot bud galls from Angophora and Eucalyptus in Australia, with descriptions of six new species</p>. <i>Zootaxa</i> , 2013, 3741, 101.	0.5	11
76	Nematodes from galls on Myrtaceae. VI. <i>Fergusobia</i> from galls on <i>Angophora</i> in Australia, with description of <i>F. colbrani</i> n. sp. and key. <i>Zootaxa</i> , 2014, 3856, 326-48.	0.5	11
77	Allodiplogaster josephi n. sp. and <i>A. seani</i> n. sp. (Nematoda: Diplogastridae), associates of soil-dwelling bees in the eastern USA. <i>Nematology</i> , 2015, 17, 831-863.	0.6	11
78	<i>Bursaphelenchus anatolius</i> n. sp. (Nematoda: Parasitaphelenchidae), an Associate of Bees in the Genus <i>Halictus</i> . <i>Journal of Nematology</i> , 2005, 37, 336-42.	0.9	11
79	Evaluation of Econemâ„¢, a formulated <i>Pasteuria</i> sp. Bionematicide, for management of <i>Belonolaimus longicaudatus</i> on golf course turf. <i>Journal of Nematology</i> , 2011, 43, 101-9.	0.9	11
80	Effects of formulation and host nematode density on the ability of in vitro-produced <i>Pasteuria</i> endospores to control its host <i>Belonolaimus longicaudatus</i> . <i>Journal of Nematology</i> , 2010, 42, 87-90.	0.9	11
81	<i>Poikilolaimus carsiops</i> n. sp. (Rhabditida: Rhabditidae) associated with <i>Neotermes koshunensis</i> (Kalotermitidae) in Kenting National Park, Taiwan. <i>Nematology</i> , 2011, 13, 155-164.	0.6	10
82	Nematodes from galls on Myrtaceae. VII. <i>Fergusobia</i> from leafy leaf bud galls in Australia, with re-description of <i>Fergusobia tumifaciens</i> (Currie 1937) Wachek 1955 and descriptions of <i>Fergusobia planchoniana</i> n. sp. and <i>Fergusobia viminalisae</i> n. sp.. <i>Zootaxa</i> , 2014, 3856, 529-54.	0.5	10
83	<i>Bursaphelenchus penai</i> n. sp. (Tylenchomorpha: Aphelenchoididae), a phoretic associate of ambrosia beetles (Coleoptera: Scolytinae) from avocado in Florida. <i>Nematology</i> , 2014, 16, 683-693.	0.6	10
84	<i>Bursaphelenchus acaloleptae</i> n. sp. sharing tree and beetle carrier hosts with <i>B. luxuriosae</i> Kanzaki & Futai, 2003 in Japan. <i>Nematology</i> , 2020, 22, 515-527.	0.6	10
85	Morphological and Molecular Characterization of <i>Labrys filiformis</i> n. sp. (Rhabditida: Tj ETQq1 1 0.784314 rgBT /Overlock 10 TF 5	0.9	10
86	Ultrastructure and life history of <i>Myolaimus byersi</i> n. sp. (Myolaimina: Myolaimidae), a phoretic associate of the crane fly, <i>Limonia schwarzi</i> (Alexander) (Limonidae), in Florida. <i>Nematology</i> , 2010, 12, 519-542.	0.6	9
87	Bermudagrass Cultivar Responses to Sting Nematodes. <i>Crop Science</i> , 2011, 51, 2199-2203.	1.8	9
88	<i>Diploscapter formicidae</i> sp. n. (Rhabditida: Diploscapteridae), from the ant <i>Prolasius advenus</i> (Hymenoptera: Formicidae) in New Zealand. <i>Nematology</i> , 2013, 15, 109-123.	0.6	9
89	Plant invasions from a belowground nematocentric perspective. <i>Soil Biology and Biochemistry</i> , 2014, 77, 213-220.	8.8	9
90	Morphology and Description of <i>Bursaphelenchus platzeri</i> n. sp. (Nematoda: Parasitaphelenchidae), an Associate of Nitidulid Beetles. <i>Journal of Nematology</i> , 2006, 38, 150-7.	0.9	9

#	ARTICLE	IF	CITATIONS
91	Diplogastrellus metamasius n. sp. (Rhabditida: Diplogastridae) isolated from <i>Metamasius hemipterus</i> (L.) (Coleoptera: Dryophthoridae) in southern Florida and Costa Rica. <i>Nematology</i> , 2008, 10, 853-868.	0.6	8
92	Ficotylus laselvae n. sp. (Tylenchomorpha: Anguinidae) associated with <i>Ficus colubrinae</i> in Costa Rica. <i>Nematology</i> , 2014, 16, 1139-1151.	0.6	8
93	Loop-mediated isothermal amplification (LAMP) for detection of the red ring nematode, <i>Bursaphelenchus cocophilus</i> . <i>Nematology</i> , 2017, 19, 559-565.	0.6	8
94	Nematodes associated with palm and sugarcane weevils in South Florida with a description of <i>Acrostichus floridensis</i> n. sp.. <i>Nematology</i> , 2017, 19, 515-531.	0.6	8
95	<i>Ficophagus auriculatae</i> n. sp. (Nematoda: Aphelenchoididae), an associate of the <i>Ficus auriculata</i> -complex in China. <i>Nematology</i> , 2019, 21, 497-508.	0.6	8
96	Description of <i>Basilaphelenchus brevicaudatus</i> n. sp. (Aphelenchoidea: Typhloelenchinae) from a dead forest tree in northern Iran. <i>Nematology</i> , 2019, 21, 147-158.	0.6	8
97	<i>Bradyneuma listronoti</i> n. sp. (Nematoda: Allantonematidae), a parasite of the carrot weevil <i>Listronotus oregonensis</i> (Coleoptera: Curculionidae) in Quebec, Canada. <i>Nematology</i> , 2007, 9, 608-622.	0.6	7
98	<i>Bursaphelenchus debrae</i> n. sp. (Nematoda: Parasitaphelenchidae), an associate of the bee <i>Halictus brunnescens</i> in Turkey. <i>Nematology</i> , 2007, 9, 777-789.	0.6	7
99	Redescription of four species of North American <i>Bursaphelenchus</i> Fuchs, 1937 (Nematoda: Tylaphelenchinae). <i>Typhloelenchia</i> 10, 7-15.	0.6	7
100	Responses of Seashore Paspalum Cultivars to Sting and Spiral Nematodes. <i>Crop Science</i> , 2011, 51, 2864-2867.	1.8	7
101	Sexually limited association of <i>Fergusobia ergusobia</i> (nematodes with female flies in a unique <i>Australasian mutualism</i> (Nematoda: Etylenchidae; Diptera: Fergusoniidae). <i>Australian Journal of Entomology</i> , 2013, 52, 125-128.	1.1	7
102	Nematodes from galls on Myrtaceae. VIII. <i>Fergusobia</i> from small galls on shoot buds, with descriptions of four new species. <i>Zootaxa</i> , 2014, 3857, 1-40.	0.5	7
103	Recharacterisation of three <i>Parasitodiplogaster</i> species based on morphological and molecular profiles. <i>Nematology</i> , 2016, 18, 417-437.	0.6	7
104	<i>Basilaphelenchus gorganensis</i> n. sp. (Aphelenchoidea, Typhloelenchinae) from wood from northern Iran. <i>Forest Pathology</i> , 2019, 49, e12490.	1.1	7
105	Field Responses of Bermudagrass and Seashore paspalum Cultivars to Sting and Spiral Nematodes. <i>Journal of Nematology</i> , 2011, 43, 201-8.	0.9	7
106	The nature and frequency of chimeras in eukaryotic metagenetic samples. <i>Journal of Nematology</i> , 2012, 44, 18-25.	0.9	7
107	Intraspecific Dueling in Palm Aphids, <i>Cerataphis brasiliensis</i> (Homoptera: Hormaphididae). <i>Florida Entomologist</i> , 1998, 81, 552.	0.5	6
108	The release and unsuccessful establishment of the <i>Melaleuca</i> biological control agent <i>Fergusonia turneri</i> and its mutualistic nematode <i>Fergusobia quinquenerviae</i> . <i>BioControl</i> , 2013, 58, 553-561.	2.0	6

#	ARTICLE	IF	CITATIONS
109	Nematodes from galls on Myrtaceae. X. <i>Fergusobia</i> from galls on narrow-leaved <i>Melaleuca</i> spp. in Australia, with descriptions of three new species. <i>Zootaxa</i> , 2014, 3889, 237-58.	0.5	6
110	Nematodes from galls on Myrtaceae. IX. <i>Fergusobia rosettae</i> n. sp. on <i>Melaleuca quinquenervia</i> and <i>F. tolgaensis</i> n. sp. on <i>Syzygium luehmannii</i> , from Queensland. <i>Zootaxa</i> , 2014, 3889, 214-36.	0.5	6
111	<i>Rhabditoides humicolus</i> n. sp. associated with arthropods from rotting palm tissue in Florida, USA. <i>Nematology</i> , 2015, 17, 487-498.	0.6	6
112	Two new species and three morphospecies of <i>Ficophagus</i> Davies & Bartholomaeus, 2015 (Nematoda: Tj ETQq0 O O rgBT /Overlock 2017, 19, 351-374.	0.6	6
113	Phoresy and Within-colony Transmission of Nematodes Associated with Alates of Subterranean Termites (Isoptera: Rhinotermitidae). <i>Environmental Entomology</i> , 2018, 47, 1107-1116.	1.4	6
114	<i>Ficophagus curtipes</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus curtipes</i> in China. <i>Nematology</i> , 2019, 21, 1091-1101.	0.6	6
115	<i>Parasitodiplogaster salicifoliae</i> n. sp. (Nematoda: Diplogastridae) from <i>Ficus ingens</i> and <i>F. salicifolia</i> in South Africa. <i>Nematology</i> , 2015, 17, 301-311.	0.6	5
116	<i>Acrostichus palmarum</i> n. sp., a cryptic species separated from <i>A. rhynchophori</i> by molecular sequences and hybridisation tests. <i>Nematology</i> , 2018, 20, 751-768.	0.6	5
117	<i>Ficophagus chaozhouensis</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus hirta</i> Vahl var. <i>roxburghii</i> (Miq.) King in China. <i>Nematology</i> , 2020, 22, 299-312.	0.6	5
118	The genus <i>Berntsenus</i> Massey, 1974 is a junior synonym of <i>Bursaphelenchus</i> Fuchs, 1937. <i>Nematology</i> , 2020, 22, 677-695.	0.6	5
119	Screening Bermudagrass Germplasm Accessions for Tolerance to Sting Nematodes. <i>Hortscience: A Publication of the American Society for Horticultural Science</i> , 2011, 46, 1503-1506.	1.0	5
120	<i>Ficophagus</i> (Nematoda: Aphelenchoididae) from sycones of <i>Ficus</i> subgenus <i>Urostigma</i> , sections <i>Malvanthera</i> and <i>Urostigma</i> , in eastern Australia. <i>Nematology</i> , 2020, 22, 627-653.	0.6	5
121	Effects of Infection by <i>Belonolaimus longicaudatus</i> on Rooting Dynamics among St. Augustinegrass and Bermudagrass Genotypes. <i>Journal of Nematology</i> , 2015, 47, 322-31.	0.9	5
122	Calling problems – the <i>Fergusobia</i> nematode/ <i>Fergusonina</i> fly mutualism on myrtaceous hosts. <i>Nematology</i> , 2016, 18, 629-649.	0.6	4
123	Recharacterisation and photo-documentation of five North American <i>Bursaphelenchus</i> species from C.L. Massey's type material. <i>Nematology</i> , 2016, 18, 311-336.	0.6	4
124	On the morphology and phylogeny of <i>Robustodorus</i> Andrassy, 2007 and two Aphelenchoides™ species (Nematoda: Aphelenchoidinae). <i>Nematology</i> , 2018, 20, 601-615.	0.6	4
125	Nematode epibionts on skin of the Florida manatee, <i>Trichechus manatus latirostris</i> . <i>Scientific Reports</i> , 2021, 11, 1211.	3.3	4
126	Phylogenetic Status and Morphological Characters of <i>Rhabditolaimus anoplophorae</i> (Rhabditida: Tj ETQq0 O O rgBT /Overlock 10 Tf 50 0.9		

#	ARTICLE	IF	CITATIONS
127	Integrated Pest Management of Nematodes on Bermudagrass Turf. Crop, Forage and Turfgrass Management, 2016, 2, 1-9.	0.6	3
128	Tylenchid entomoparasites isolated from <i>Spondylis buprestoides</i> (L.) and <i>Asemum striatum</i> (L.) (Coleoptera: Cerambycidae). Nematology, 2016, 18, 775-780.	0.6	3
129	<i>Sudhausia floridensis</i> n. sp. (Nematoda: Diplogastridae) isolated from <i>Onthophagus tuberculifrons</i> (Coleoptera: Scarabaeidae) from Florida, USA. Nematology, 2017, 19, 575-586.	0.6	3
130	<i>Martininema semicordatae</i> n. sp. (Nematoda: Aphelenchoididae) associated with <i>Ficus semicordata</i> in China. Nematology, 2020, 22, 759-770.	0.6	3
131	Morphological and molecular characteristics of <i>Parasitodiplogaster religiosae</i> n. sp. (Nematoda:) Tj ETQq1 1 0.784314 rgBT /Qverlock 10	2.5	1
132	<i>Schistonchus macrophylla</i> (Nematoda: Aphelenchoididae): an Åenigmatic taxon from sycones of <i>Ficus</i> section <i>Malvanthera</i> in Åeastern Australia, with description of a new subspecies. Nematology, 2020, 22, 999-1013.	0.6	2
133	Molecular diversity and relationships of fig associated nematodes from South Africa. PLoS ONE, 2021, 16, e0255451.	2.5	2
134	Effects of entomopathogenic nematodes and/or fungus on the Åred Åpalm weevil, <i>Rhynchophorus ferrugineus</i> (Curculionidae: Coleoptera). Nematology, 2020, 22, 1193-1207.	0.6	2
135	<i>Bursaphelenchus suri</i> n. sp.: A second <i>Bursaphelenchus</i> syconial parasite of figs supports adaptive radiation among section <i>Sycomorus</i> figs. PLoS ONE, 2022, 17, e0265339.	2.5	2
136	<p class="HeadingRunIn">YONGSAN ZENG, WEIMIN YE, JIANGHUA HUANG, CHANGHUI LI & ROBIN M. GIBLIN-DAVIS (2013) Description of Schistonchus altissimus n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus altissima</i> in China. Zootaxa, 3700 (4), 561â€“572.</p>. Zootaxa, 2013, 3717, 598.	0.5	1
137	Who has â€˜Dibslâ€™™ on a dead rhinoceros beetle?. Environmental Microbiology, 2017, 19, 2115-2116.	3.8	0
138	Nematodes from galls on Myrtaceae. XI. Descriptions of five new species of <i>Fergusobia</i> from Australia. Zootaxa, 2018, 4399, 1.	0.5	0