

Robin M Giblin-Davis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6636054/publications.pdf>

Version: 2024-02-01

138
papers

2,874
citations

201674

27
h-index

233421

45
g-index

139
all docs

139
docs citations

139
times ranked

1416
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>Bursaphelenchus suri</i> n. sp.: A second <i>Bursaphelenchus</i> syconial parasite of figs supports adaptive radiation among section <i>Sycomorus</i> figs. PLoS ONE, 2022, 17, e0265339.	2.5	2
2	Molecular diversity and relationships of fig associated nematodes from South Africa. PLoS ONE, 2021, 16, e0255451.	2.5	2
3	Nematode epibionts on skin of the Florida manatee, <i>Trichechus manatus latirostris</i> . Scientific Reports, 2021, 11, 1211.	3.3	4
4	<i>Ficophagus chaozhouensis</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus hirta</i> Vahl var. <i>roxburghii</i> (Miq.) King in China. Nematology, 2020, 22, 299-312.	0.6	5
5	<i>Martininema semicordatae</i> n. sp. (Nematoda: Aphelenchoididae) associated with <i>Ficus semicordata</i> in China. Nematology, 2020, 22, 759-770.	0.6	3
6	<i>Bursaphelenchus acaloleptae</i> n. sp. sharing tree and beetle carrier hosts with <i>B. luxuriosae</i> Kanzaki & Futai, 2003 in Japan. Nematology, 2020, 22, 515-527.	0.6	10
7	The genus <i>Berntsensus</i> Massey, 1974 is a junior synonym of <i>Bursaphelenchus</i> Fuchs, 1937. Nematology, 2020, 22, 677-695.	0.6	5
8	<i>Schistonchus macrophylla</i> (Nematoda: Aphelenchoididae): an enigmatic taxon from sycones of <i>Ficus</i> section <i>Malvanthera</i> in eastern Australia, with description of a new subspecies. Nematology, 2020, 22, 999-1013.	0.6	2
9	A new species of <i>Basilaphelenchus</i> Pedram, Kanzaki, Giblin-Davis & Pourjam, 2018 (Aphelenchoidea: Tj ETQq1 1 0.784314 rgBT / Over	0.6	13
10	Effects of entomopathogenic nematodes and/or fungus on the red palm weevil, <i>Rhynchophorus ferrugineus</i> (Curculionidae: Coleoptera). Nematology, 2020, 22, 1193-1207.	0.6	2
11	<i>Ficophagus</i> (Nematoda: Aphelenchoididae) from sycones of <i>Ficus</i> subgenus <i>Urostigma</i> , sections <i>Malvanthera</i> and <i>Urostigma</i> , in eastern Australia. Nematology, 2020, 22, 627-653.	0.6	5
12	Feeding Dimorphism in a Mycophagous Nematode, <i>Bursaphelenchus sinensis</i> . Scientific Reports, 2019, 9, 13956.	3.3	15
13	<i>Ficophagus auriculatae</i> n. sp. (Nematoda: Aphelenchoididae), an associate of the <i>Ficus auriculata</i> -complex in China. Nematology, 2019, 21, 497-508.	0.6	8
14	<i>Ficophagus curtipes</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus curtipes</i> in China. Nematology, 2019, 21, 1091-1101.	0.6	6
15	Description of <i>Basilaphelenchus brevicaudatus</i> n. sp. (Aphelenchoidea: Tylaphelenchinae) from a dead forest tree in northern Iran. Nematology, 2019, 21, 147-158.	0.6	8
16	<i>Basilaphelenchus gorganensis</i> n. sp. (Aphelenchoidea, Tylaphelenchinae) from wood from northern Iran. Forest Pathology, 2019, 49, e12490.	1.1	7
17	On the morphology and phylogeny of <i>Robustodorus</i> Andr�ssy, 2007 and two "Aphelenchoidea" species (Nematoda: Aphelenchoidinae). Nematology, 2018, 20, 601-615.	0.6	4
18	Diversity and Plant Pathogenicity of <i>Bursaphelenchus</i> and Related Nematodes in Relation to Their Vector Bionomics. Current Forestry Reports, 2018, 4, 85-100.	7.4	44

#	ARTICLE	IF	CITATIONS
19	Acrostichus palmarum n. sp., a cryptic species separated from A.Ârhynchophori by molecular sequences and hybridisation tests. Nematology, 2018, 20, 751-768.	0.6	5
20	A molecular phylogenetic approach for unravelling the taxonomic status of Basilaphelenchus persicus n. gen., n. sp. (Aphelenchoididae: Tylaphelenchinae). Nematology, 2018, 20, 567-582.	0.6	13
21	Nematodes from galls on Myrtaceae. XI. Descriptions of five new species of Fergusobia from Australia. Zootaxa, 2018, 4399, 1.	0.5	0
22	Phoresy and Within-colony Transmission of Nematodes Associated with Alates of Subterranean Termites (Isoptera: Rhinotermitidae). Environmental Entomology, 2018, 47, 1107-1116.	1.4	6
23	Morphological and molecular characteristics of Parasitodiplogaster religiosae n. sp. (Nematoda: Tj ETQq1 1 0.784314 rgBT /Qverlock 2.5 2	2.5	2
24	Morphological and Molecular Characterization of <i>Labrys filiformis</i> n. sp. (Rhabditida: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542 T	0.9	10
25	Two new species and three morphospecies of Ficophagus Davies& Bartholomaeus, 2015 (Nematoda: Tj ETQq1 1 0.784314 rgBT 2017, 19, 351-374.	0.6	6
26	Loop-mediated isothermal amplification (LAMP) for detection of the red ring nematode, Bursaphelenchus cocophilus. Nematology, 2017, 19, 559-565.	0.6	8
27	Sudhausia floridensis n. sp. (Nematoda: Diplogastridae) isolated from Onthophagus tuberculifrons (Coleoptera: Scarabaeidae) from Florida, USA. Nematology, 2017, 19, 575-586.	0.6	3
28	Who has â€Dibslâ€™™ on a dead rhinoceros beetle?. Environmental Microbiology, 2017, 19, 2115-2116.	3.8	0
29	Nematodes associated with palm and sugarcane weevils in South Florida with a description of Acrostichus floridensis n. sp.. Nematology, 2017, 19, 515-531.	0.6	8
30	Integrated Pest Management of Nematodes on Bermudagrass Turf. Crop, Forage and Turfgrass Management, 2016, 2, 1-9.	0.6	3
31	Tylenchid entomoparasites isolated from Spondylis buprestoides (L.) and Asemum striatum (L.) (Coleoptera: Cerambycidae). Nematology, 2016, 18, 775-780.	0.6	3
32	Recharacterisation of three Parasitodiplogaster species based on morphological and molecular profiles. Nematology, 2016, 18, 417-437.	0.6	7
33	Description of Ektaphelenchus berbericus n. sp. (Rhabditida: Ektaphelenchinae) from eastern Iran. Nematology, 2016, 18, 1063-1077.	0.6	18
34	Galling problemsÂ€“ the Fergusobia nematode/Fergusonina fly mutualism on myrtaceous hosts. Nematology, 2016, 18, 629-649.	0.6	4
35	Large-scale diversification without genetic isolation in nematode symbionts of figs. Science Advances, 2016, 2, e1501031.	10.3	82
36	Recharacterisation and photo-documentation of five NorthÂAmerican Bursaphelenchus species fromÂC.L.ÂMasseyâ€™sÂtypeÂmaterial. Nematology, 2016, 18, 311-336.	0.6	4

#	ARTICLE	IF	CITATIONS
37	Description of <i>Ruehmaphelenchus juliae</i> n. sp. (Tylenchina: Aphelenchoididae) isolated from an ambrosia beetle, <i>Xylosandrus crassiusculus</i> (Motschulsky), from South Florida. <i>Nematology</i> , 2015, 17, 639-653.	0.6	13
38	A review of the taxonomy, phylogeny, distribution and co-evolution of <i>Schistonchus</i> Cobb, 1927 with proposal of <i>Ficophagus</i> n. gen. and <i>Martininema</i> n. gen. (Nematoda: Aphelenchoididae). <i>Nematology</i> , 2015, 17, 761-829.	0.6	46
39	<i>Allodiplogaster josephi</i> n. sp. and <i>A. seani</i> n. sp. (Nematoda: Diplogastridae), associates of soil-dwelling bees in the eastern USA. <i>Nematology</i> , 2015, 17, 831-863.	0.6	11
40	The fig wasp followers and colonists of a widely introduced fig tree, <i>Ficus microcarpa</i> . <i>Insect Conservation and Diversity</i> , 2015, 8, 322-336.	3.0	27
41	<i>Parasitodiplogaster salicifoliae</i> n. sp. (Nematoda: Diplogastridae) from <i>Ficus ingens</i> and <i>F. salicifolia</i> in South Africa. <i>Nematology</i> , 2015, 17, 301-311.	0.6	5
42	<i>Rhabditoides humicolus</i> n. sp. associated with arthropods from rotting palm tissue in Florida, USA. <i>Nematology</i> , 2015, 17, 487-498.	0.6	6
43	Effects of Infection by <i>Belonolaimus longicaudatus</i> on Rooting Dynamics among St. Augustinegrass and Bermudagrass Genotypes. <i>Journal of Nematology</i> , 2015, 47, 322-31.	0.9	5
44	Nematodes from galls on Myrtaceae. VII. <i>Fergusobia</i> from "leafy" leaf bud galls in Australia, with re-description of <i>Fergusobia tumifaciens</i> (Currie 1937) Wachek 1955 and descriptions of <i>Fergusobia planchoniana</i> n. sp. and <i>Fergusobia viminalisae</i> n. sp.. <i>Zootaxa</i> , 2014, 3856, 529-54.	0.5	10
45	Nematodes from galls on Myrtaceae. X. <i>Fergusobia</i> from galls on narrow-leaved <i>Melaleuca</i> spp. in Australia, with descriptions of three new species. <i>Zootaxa</i> , 2014, 3889, 237-58.	0.5	6
46	Nematodes from galls on Myrtaceae. VI. <i>Fergusobia</i> from galls on <i>Angophora</i> in Australia, with description of <i>F. colbrani</i> n. sp. and key. <i>Zootaxa</i> , 2014, 3856, 326-48.	0.5	11
47	Nematodes from galls on Myrtaceae. VIII. <i>Fergusobia</i> from small galls on shoot buds, with descriptions of four new species. <i>Zootaxa</i> , 2014, 3857, 1-40.	0.5	7
48	Nematodes from galls on Myrtaceae. IX. <i>Fergusobia rosettae</i> n. sp. on <i>Melaleuca quinquenervia</i> and <i>F. tolgaensis</i> n. sp. on <i>Syzygium luehmannii</i> , from Queensland. <i>Zootaxa</i> , 2014, 3889, 214-36.	0.5	6
49	<i>Bursaphelenchus penai</i> n. sp. (Tylenchomorpha: Aphelenchoididae), a phoretic associate of ambrosia beetles (Coleoptera: Scolytinae) from avocado in Florida. <i>Nematology</i> , 2014, 16, 683-693.	0.6	10
50	Description of two <i>Pseudaphelenchus</i> species (Tylenchomorpha: Aphelenchoididae) associated with Asian termites and proposal of <i>Tylaphelenchinae</i> n. subfam.. <i>Nematology</i> , 2014, 16, 963-978.	0.6	29
51	Revision of the paraphyletic genus <i>Koerneria</i> Meyl, 1960 and resurrection of two other genera of Diplogastridae (Nematoda). <i>ZooKeys</i> , 2014, 442, 17-30.	1.1	13
52	Molecular and morphological observations on <i>Parasitodiplogaster sycophilon</i> Poinar, 1979 (Nematoda: Diplogastrina) associated with <i>Ficus burkei</i> in Africa. <i>Nematology</i> , 2014, 16, 453-462.	0.6	12
53	<i>Ficotylus laselvae</i> n. sp. (Tylenchomorpha: Anguinidae) associated with <i>Ficus colubrinae</i> in Costa Rica. <i>Nematology</i> , 2014, 16, 1139-1151.	0.6	8
54	<i>Parasitodiplogaster</i> species associated with <i>Pharmacosycea</i> figs in Panama. <i>Nematology</i> , 2014, 16, 607-619.	0.6	17

#	ARTICLE	IF	CITATIONS
55	An emerging example of tritrophic coevolution between flies (Diptera: Fergusoninidae) and nematodes (Nematoda: Neotylenchidae) on Myrtaceae host plants. <i>Biological Journal of the Linnean Society</i> , 2014, 111, 699-718.	1.6	16
56	Plant invasions from a belowground nematocentric perspective. <i>Soil Biology and Biochemistry</i> , 2014, 77, 213-220.	8.8	9
57	New Plant-Parasitic Nematode from the Mostly Mycophagous Genus <i>Bursaphelenchus</i> Discovered inside Figs in Japan. <i>PLoS ONE</i> , 2014, 9, e99241.	2.5	38
58	Phylogenetic Status and Morphological Characters of <i>Rhabditolaimus anoplophorae</i> (Rhabditida: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	0.9	4
59	The release and unsuccessful establishment of the <i>Melaleuca</i> biological control agent <i>Fergusonina turneri</i> and its mutualistic nematode <i>Fergusobia quinquenerviae</i> . <i>BioControl</i> , 2013, 58, 553-561.	2.0	6
60	<i>Schistonchus molochi</i> sp. n. and <i>S. athertonensis</i> sp. n. (Nematoda: Aphelenchoididae) from <i>Ficus watkinsiana</i> (Moraceae) in Australia. <i>Nematology</i> , 2013, 15, 389-400.	0.6	13
61	Distribution of southern sting nematode, <i>Ibipora lolii</i> (Nematoda: Belonolaimidae), on turfgrass in Australia and its taxonomic relationship to other belonolaimids. <i>Nematology</i> , 2013, 15, 401-415.	0.6	11
62	<i>Schistonchus fistulosus</i> n. sp. (Nematoda: Aphelenchoididae) associated with <i>Ficus fistulosa</i> in China. <i>Nematology</i> , 2013, 15, 879-890.	0.6	14
63	Description of <i>Schistonchus superbus</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus superba</i> in China. <i>Nematology</i> , 2013, 15, 771-781.	0.6	14
64	Three new species of <i>Schistonchus</i> (Aphelenchoididae) from the <i>Ficus</i> subgenus <i>Sycomorus</i> (Moraceae) in northern Australia. <i>Nematology</i> , 2013, 15, 347-362.	0.6	21
65	Description of <i>Parasitodiplogaster pharmaconema</i> n. sp. and redescription of <i>P. maxinema</i> from <i>Ficus maxima</i> Mill. (Moraceae). <i>Nematology</i> , 2013, 15, 957-974.	0.6	14
66	<i>Diploscapter formicidae</i> sp. n. (Rhabditida: Diploscapteridae), from the ant <i>Prolasius advenus</i> (Hymenoptera: Formicidae) in New Zealand. <i>Nematology</i> , 2013, 15, 109-123.	0.6	9
67	Sex-limited association of <i>Fergusobia</i> nematodes with female <i>Fergusonina</i> flies in a unique Australasian mutualism (Nematoda: Neotylenchidae; Diptera: Fergusoninidae). <i>Australian Journal of Entomology</i> , 2013, 52, 125-128.	1.1	7
68	Resource Opportunities from the Nest of Dying Subterranean Termite (Isoptera: Rhinotermitidae) Colonies: A Laboratory Case of Ecological Succession. <i>Annals of the Entomological Society of America</i> , 2013, 106, 771-778.	2.5	15
69	Molecular Characterization and Development of Real-Time PCR Assay for Pine-Wood Nematode <i>Bursaphelenchus xylophilus</i> (Nematoda: Parasitaphelenchidae). <i>PLoS ONE</i> , 2013, 8, e78804.	2.5	24
70	Description of <i>Schistonchus altissimus</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus altissima</i> in China. <i>Zootaxa</i> , 2013, 3700, 561.	0.5	15
71	Nematodes from galls on Myrtaceae. <i>V. Fergusobia</i> from large multilocular shoot bud galls from <i>Angophora</i> and <i>Eucalyptus</i> in Australia, with descriptions of six new species. <i>Zootaxa</i> , 2013, 3741, 101.	0.5	11
72	Nematodes from galls on Myrtaceae. IV. <i>Fergusobia</i> from flat leaf galls on <i>Eucalyptus</i> and <i>Corymbia</i> , with descriptions of two new species. <i>Zootaxa</i> , 2013, 3741, 151.	0.5	12

#	ARTICLE	IF	CITATIONS
73	<p><p class="HeadingRunIn">YONGSAN ZENG, WEIMIN YE, JIANGHUA HUANG, CHANGHUI LI & ROBIN M. GIBLIN-DAVIS (2013) Description of <i>Schistonchus altissimus</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus altissima</i> in China. <i>Zootaxa</i>, 3700 (4), 561-572.</p></p>	0.5	1
74	<p>Schistonchus (Aphelenchoididae) from <i>Ficus benjamina</i> in Australia, with description of <i>S. benjamina</i> sp. n.. <i>Nematology</i>, 2012, 14, 1005-1013.</p>	0.6	16
75	<p><i>Teratodiplogaster martini</i> n. sp. and <i>Parasitodiplogaster doliostroma</i> sp. (Nematoda: Diplogastridae) from the syconia of <i>Ficus</i> species from Africa. <i>Nematology</i>, 2012, 14, 529-546.</p>	0.6	20
76	<p>Nematodes from galls on Myrtaceae. II. <i>Fergusobia</i>/<i>Fergusonina</i> from small axillary bud ('stem') and leaf ('pea') galls in Australia, with descriptions of two new species. <i>Zootaxa</i>, 2012, 3415, .</p>	0.5	16
77	<p>Nematodes from galls on Myrtaceae. III. <i>Fergusobia</i> from flower bud and stigma galls on <i>Eucalyptus</i>, with descriptions of four new species. <i>Zootaxa</i>, 2012, 3532, 1.</p>	0.5	14
78	<p>The nature and frequency of chimeras in eukaryotic metagenetic samples. <i>Journal of Nematology</i>, 2012, 44, 18-25.</p>	0.9	7
79	<p>Responses of Seashore <i>Paspalum</i> Cultivars to Sting and Spiral Nematodes. <i>Crop Science</i>, 2011, 51, 2864-2867.</p>	1.8	7
80	<p>Description of <i>Bursaphelenchus kiyoharai</i> n. sp. (Tylenchina: Aphelenchoididae) with remarks on the taxonomic framework of the <i>Parasitaphelenchinae</i> R&A, 1956 and <i>Aphelenchoidinae</i> Fuchs, 1937. <i>Nematology</i>, 2011, 13, 787-804.</p>	0.6	36
81	<p><i>Poikilolaimus carsiops</i> n. sp. (Rhabditida: Rhabditidae) associated with <i>Neotermes koshunensis</i> (Kalotermitidae) in Kenting National Park, Taiwan. <i>Nematology</i>, 2011, 13, 155-164.</p>	0.6	10
82	<p>Description of <i>Schistonchus microcarpus</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus microcarpa</i> in China. <i>Nematology</i>, 2011, 13, 221-233.</p>	0.6	26
83	<p>Description of <i>Caenorhabditis angaria</i> n. sp. (Nematoda: Rhabditidae), an associate of sugarcane and palm weevils (Coleoptera: Curculionidae). <i>Nematology</i>, 2011, 13, 61-78.</p>	0.6	23
84	<p>Bermudagrass Cultivar Responses to Sting Nematodes. <i>Crop Science</i>, 2011, 51, 2199-2203.</p>	1.8	9
85	<p>Screening Bermudagrass Germplasm Accessions for Tolerance to Sting Nematodes. <i>Hortscience: A Publication of the American Society for Horticultural Science</i>, 2011, 46, 1503-1506.</p>	1.0	5
86	<p>Evaluation of Econem[®], a formulated <i>Pasteuria</i> sp. Bionematicide, for management of <i>Belonolaimus longicaudatus</i> on golf course turf. <i>Journal of Nematology</i>, 2011, 43, 101-9.</p>	0.9	11
87	<p>Field Responses of Bermudagrass and Seashore <i>Paspalum</i> Cultivars to Sting and Spiral Nematodes. <i>Journal of Nematology</i>, 2011, 43, 201-8.</p>	0.9	7
88	<p>Ecometagenetics confirm high tropical rainforest nematode diversity. <i>Molecular Ecology</i>, 2010, 19, 5521-5530.</p>	3.9	90
89	<p>The nematode genus <i>Fergusobia</i> (Nematoda: Neotylenchidae): molecular phylogeny, descriptions of clades and associated galls, host plants and <i>Fergusonina</i> fly larvae. <i>Zootaxa</i>, 2010, 2633, 1.</p>	0.5	35
90	<p>Linking operational clustered taxonomic units (OCTUs) from parallel ultra sequencing (PUS) to nematode species. <i>Zootaxa</i>, 2010, 2427, .</p>	0.5	40

#	ARTICLE	IF	CITATIONS
91	<i>Ditylenchus halictus</i> n. sp. (Nematoda: Anguinidae), an associate of the sweat bee, <i>Halictus sexcinctus</i> (Halictidae), from Germany. <i>Nematology</i> , 2010, 12, 891-904.	0.6	13
92	Redescription of two Panamanian nematodes, <i>Parasitodiplogaster citrinema</i> Poinar & Herre, 1991 and <i>P. popenema</i> Poinar & Herre, 1991 (Nematoda: Diplogastrina). <i>Nematology</i> , 2010, 12, 89-104.	0.6	11
93	<i>Schistonchus</i> (Aphelenchoididae) from <i>Ficus</i> (Moraceae) in Australia, with description of <i>S. aculeata</i> sp. n.. <i>Nematology</i> , 2010, 12, 935-958.	0.6	37
94	Ultrastructure and life history of <i>Myolaimus byersi</i> n. sp. (Myolaimina: Myolaimidae), a phoretic associate of the crane fly, <i>Limonia schwarzi</i> (Alexander) (Limoniidae), in Florida. <i>Nematology</i> , 2010, 12, 519-542.	0.6	9
95	<i>Acrostichus puri</i> n. sp. (Nematoda: Diplogastridae), a phoretic associate of <i>Augochlora pura mosieri</i> Cockerell (Hymenoptera: Halictidae). <i>Nematology</i> , 2010, 12, 49-64.	0.6	16
96	<i>Acrostichus megaloptae</i> n. sp. (Nematoda: Diplogastridae), a phoretic associate of <i>Megalopta</i> spp. (Hymenoptera: Halictidae) in Central America. <i>Nematology</i> , 2010, 12, 453-468.	0.6	14
97	<i>Schistonchus hirtus</i> n. sp. (Nematoda: Aphelenchoididae), an associate of <i>Ficus hirta</i> in China. <i>Nematology</i> , 2010, 12, 543-556.	0.6	23
98	Diversity and distribution of nematodes associated with wild bees in Turkey. <i>Nematology</i> , 2010, 12, 65-80.	0.6	12
99	<i>Pseudaphelenchus vindai</i> n. sp. (Tylenchomorpha: Aphelenchoididae) associated with termites (Termitidae) in Barro Colorado Island, Panama. <i>Nematology</i> , 2010, 12, 905-914.	0.6	18
100	Reproducibility of read numbers in high-throughput sequencing analysis of nematode community composition and structure. <i>Molecular Ecology Resources</i> , 2010, 10, 666-676.	4.8	93
101	Nematodes from galls on Myrtaceae. I. <i>Fergusobia</i> / <i>Fergusonina</i> galls on <i>Corymbia</i> spp., with re-description of <i>F. magna</i> and notes on its phylogenetic relationships. <i>Zootaxa</i> , 2010, 2634, 25.	0.5	19
102	Effects of formulation and host nematode density on the ability of in vitro-produced pasteurized endospores to control its host <i>Belonolaimus longicaudatus</i> . <i>Journal of Nematology</i> , 2010, 42, 87-90.	0.9	11
103	Redescription of four species of North American <i>Bursaphelenchus</i> Fuchs, 1937 (Nematoda: Tylenchomorpha: Bursaphelenchidae). <i>Nematology</i> , 2010, 12, 959-967.	0.784314	107
104	<i>Poikilolaimus floridensis</i> n. sp. (Rhabditida: Rhabditidae) associated with termites (Kalotermitidae). <i>Nematology</i> , 2009, 11, 203-216.	0.6	14
105	<i>Schistonchus virens</i> sp. n. (Aphelenchoididae) and <i>Parasitodiplogaster australis</i> sp. n. (Diplogastridae) from <i>Ficus virens</i> (Moraceae) in Australia. <i>Nematology</i> , 2009, 11, 583-601.	0.6	33
106	<i>Bursaphelenchus tokyoensis</i> n. sp. (Nematoda: Parasitaphelenchinae) isolated from dead wood of the Japanese red pine, <i>Pinus densiflora</i> Sieb. & Zucc., in Japan. <i>Nematology</i> , 2009, 11, 171-180.	0.6	13
107	<i>Acrostichus rhynchophori</i> n. sp. (Rhabditida: Diplogastridae): a phoretic associate of <i>Rhynchophorus cruentatus</i> Fabricius and <i>R. palmarum</i> L. (Coleoptera: Curculionidae) in the Americas. <i>Nematology</i> , 2009, 11, 669-688.	0.6	21
108	<i>Ficotylus congestae</i> gen. n., sp. n. (Anguinata), from <i>Ficus congesta</i> (Moraceae) sycones in Australia. <i>Nematology</i> , 2009, 11, 63-75.	0.6	20

#	ARTICLE	IF	CITATIONS
109	<i>Pseudaphelenchus yukiae</i> n. gen., n. sp. (Tylenchina: Aphelenchoididae) associated with <i>Cylindrotermes macrognathus</i> (Termitidae: Termitinae) in La Selva, Costa Rica. <i>Nematology</i> , 2009, 11, 869-881.	0.6	38
110	<i>Teratodiplogaster fignewmani</i> gen. nov., sp. nov. (Nematoda: Diplogastridae) from the Syconia of <i>Ficus racemosa</i> in Australia. <i>Zoological Science</i> , 2009, 26, 569-578.	0.7	29
111	Evaluating high-throughput sequencing as a method for metagenomic analysis of nematode diversity. <i>Molecular Ecology Resources</i> , 2009, 9, 1439-1450.	4.8	264
112	Description of <i>Ektaphelenchoides spondylis</i> n. sp. (Nematoda: Ektaphelenchinae) isolated from <i>Spondylis buprestoides</i> (Coleoptera: Cerambycidae) in Japan. <i>Nematology</i> , 2009, 11, 181-188.	0.6	26
113	<i>Bursaphelenchus rufipennis</i> n. sp. (Nematoda: Parasitaphelenchinae) and redescription of <i>Ektaphelenchus obtusus</i> (Nematoda: Ektaphelenchinae), associates from nematangia on the hind wings of <i>Dendroctonus rufipennis</i> (Coleoptera: Scolytidae). <i>Nematology</i> , 2008, 10, 925-955.	0.6	30
114	Morphological and molecular analysis of six aphelenchoidids from Australian conifers and their relationship to <i>Bursaphelenchus</i> (Fuchs, 1937). <i>Nematology</i> , 2008, 10, 663-678.	0.6	47
115	<i>Teratorhabditis synpapillata</i> Sudhaus, 1985 (Rhabditida: Rhabditidae) is an associate of the red palm weevil, <i>Rhynchophorus ferrugineus</i> (Coleoptera: Curculionidae). <i>Nematology</i> , 2008, 10, 207-218.	0.6	18
116	Molecular variability of <i>Schistonchus caprifici</i> (Nematoda: Aphelenchoididae) from <i>Ficus carica</i> in Turkey. <i>Nematology</i> , 2008, 10, 639-649.	0.6	13
117	<i>Diplogastrellus metamasius</i> n. sp. (Rhabditida: Diplogastridae) isolated from <i>Metamasius hemipterus</i> (L.) (Coleoptera: Dryophthoridae) in southern Florida and Costa Rica. <i>Nematology</i> , 2008, 10, 853-868.	0.6	8
118	Two new species of <i>Schistonchus</i> (Nematoda: Aphelenchoididae) associated with <i>Ficus hispida</i> in China. <i>Nematology</i> , 2007, 9, 169-187.	0.6	39
119	<i>Bradynema listronoti</i> n. sp. (Nematoda: Allantonematidae), a parasite of the carrot weevil <i>Listronotus oregonensis</i> (Coleoptera: Curculionidae) in Quebec, Canada. <i>Nematology</i> , 2007, 9, 608-622.	0.6	7
120	<i>Bursaphelenchus debrae</i> n. sp. (Nematoda: Parasitaphelenchidae), an associate of the bee <i>Halictus brunnescens</i> in Turkey. <i>Nematology</i> , 2007, 9, 777-789.	0.6	7
121	Phylogenetic relationships among <i>Bursaphelenchus</i> species (Nematoda: Parasitaphelenchidae) inferred from nuclear ribosomal and mitochondrial DNA sequence data. <i>Molecular Phylogenetics and Evolution</i> , 2007, 43, 1185-1197.	2.7	260
122	Molecular phylogenetics and the evolution of host plant associations in the nematode genus <i>Fergusobia</i> (Tylenchida: Fergusobiinae). <i>Molecular Phylogenetics and Evolution</i> , 2007, 45, 123-141.	2.7	93
123	Morphology and systematics of <i>Bursaphelenchus gerberae</i> n. sp. (Nematoda: Parasitaphelenchidae), a rare associate of the palm weevil, <i>Rhynchophorus palmarum</i> in Trinidad. <i>Zootaxa</i> , 2006, 1189, 39-53.	0.5	39
124	Identification of Chilean <i>Reticulitermes</i> (Isoptera: Rhinotermitidae) Inferred from Three Mitochondrial Gene DNA Sequences and Soldier Morphology. <i>Annals of the Entomological Society of America</i> , 2006, 99, 352-363.	2.5	37
125	Morphology and Description of <i>Bursaphelenchus platzeri</i> n. sp. (Nematoda: Parasitaphelenchidae), an Associate of Nitidulid Beetles. <i>Journal of Nematology</i> , 2006, 38, 150-7.	0.9	9
126	Stomatal Ultrastructure, Molecular Phylogeny, and Description of <i>Parasitodiplogaster laevigata</i> n. sp. (Nematoda: Diplogastridae), a Parasite of Fig Wasps. <i>Journal of Nematology</i> , 2006, 38, 137-49.	0.9	31

#	ARTICLE	IF	CITATIONS
127	Bursaphelenchus anatolius n. sp. (Nematoda: Parasitaphelenchidae), an Associate of Bees in the Genus Halictus. Journal of Nematology, 2005, 37, 336-42.	0.9	11
128	The biology and associations of Fergusobia (Nematoda) from the Melaleuca leucadendra-complex in eastern Australia. Invertebrate Systematics, 2004, 18, 291.	1.3	57
129	Intraspecific Dueling in Palm Aphids, Cerataphis brasiliensis (Homoptera: Hormaphididae). Florida Entomologist, 1998, 81, 552.	0.5	6
130	Optimization of semiochemical-based trapping of Metamasius hemipterus sericeus (Olivier) (Coleoptera: Curculionidae). Journal of Chemical Ecology, 1996, 22, 1389-1410.	1.8	40
131	Evaluation of an Entomopathogenic Nematode and Chemical Insecticides for Control of Metamasius hemipterus sericeus (Coleoptera: Curculionidae). Journal of Entomological Science, 1996, 31, 240-251.	0.3	16
132	Pheromone chirality of african palm weevil, Rhynchophorus phoenicis (F.) and palmetto weevil, Rhynchophorus cruentatus (F.) (Coleoptera: Curculionidae). Journal of Chemical Ecology, 1994, 20, 2653-2671.	1.8	35
133	Fecundity and Fertility of Rhynchophorus cruentatus (Coleoptera: Curculionidae). Florida Entomologist, 1994, 77, 373.	0.5	12
134	Flight Behavior and Seasonal Trapping of Rhynchophorus cruentatus (Coleoptera: Curculionidae). Annals of the Entomological Society of America, 1994, 87, 641-647.	2.5	12
135	Laboratory and field evidence for male produced aggregation pheromone in Rhynchophorus cruentatus (F.) (Coleoptera: Curculionidae). Journal of Chemical Ecology, 1993, 19, 1195-1203.	1.8	22
136	Water Loss Dynamics and Humidity Preference of Rhynchophorus cruentatus (Coleoptera: Curculionidae). Journal of Chemical Ecology, 1993, 19, 1195-1203.	1.4	29
137	Resistance in Stenotaphrum to the Sting Nematode. Crop Science, 1993, 33, 1066-1070.	1.8	24
138	Vulnerability of Stressed Palms to Attack by Rhynchophorus cruentatus (Coleoptera: Curculionidae) and Insecticidal Control of the Pest. Journal of Economic Entomology, 1989, 82, 1185-1190.	1.8	35