## Burcin Ã-zdirik

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6647424/publications.pdf

Version: 2024-02-01

17 papers	181 citations	7 h-index	1125743 13 g-index
17	17	17	267
all docs	docs citations	times ranked	citing authors

#	Article	IF	Citations
1	Low Serum Levels of Soluble Receptor Activator of Nuclear Factor κ B Ligand (sRANKL) Are Associated with Metabolic Dysregulation and Predict Long-Term Mortality in Critically III Patients. Diagnostics, 2022, 12, 62.	2.6	1
2	Soluble Urokinase Plasminogen Activator Receptor Levels Are Associated with Severity of Fibrosis in Patients with Primary Sclerosing Cholangitis. Journal of Clinical Medicine, 2022, 11, 2479.	2.4	2
3	Models of Gastroenteropancreatic Neuroendocrine Neoplasms: Current Status and Future Directions. Neuroendocrinology, 2021, 111, 217-236.	2.5	17
4	From Liver Cirrhosis to Cancer: The Role of Micro-RNAs in Hepatocarcinogenesis. International Journal of Molecular Sciences, 2021, 22, 1492.	4.1	16
5	Lessons From Immune Checkpoint Inhibitor Trials in Hepatocellular Carcinoma. Frontiers in Immunology, 2021, 12, 652172.	4.8	21
6	The Role of Microbiota in Primary Sclerosing Cholangitis and Related Biliary Malignancies. International Journal of Molecular Sciences, 2021, 22, 6975.	4.1	22
7	Atezolizumab and bevacizumab-induced encephalitis in advanced hepatocellular carcinoma. Medicine (United States), 2021, 100, e26377.	1.0	10
8	The Role of miRNA in the Pathophysiology of Neuroendocrine Tumors. International Journal of Molecular Sciences, 2021, 22, 8569.	4.1	8
9	PD-L1 – inhibitors in neuroendocrine neoplasia. Medicine (United States), 2021, 100, e23835.	1.0	7
10	In Vivo Models for Cholangiocarcinoma—What Can We Learn for Human Disease?. International Journal of Molecular Sciences, 2020, 21, 4993.	4.1	8
11	Analysis of miR-29 Serum Levels in Patients with Neuroendocrine Tumors—Results from an Exploratory Study. Journal of Clinical Medicine, 2020, 9, 2881.	2.4	8
12	A case report of an excellent response to interferon- $\hat{l}_{\pm}$ in a patient with functional metastasized neuroendocrine tumor refractory to other treatments. Medicine (United States), 2020, 99, e20820.	1.0	1
13	Soluble Urokinase Plasminogen Activator Receptor (suPAR) Concentrations Are Elevated in Patients with Neuroendocrine Malignancies. Journal of Clinical Medicine, 2020, 9, 1647.	2.4	6
14	Systemic treatment of hepatocellular carcinoma: from sorafenib to combination therapies. Hepatic Oncology, 2020, 7, HEP20.	4.2	30
15	Primary Neuroendocrine Neoplasms of the Breast: Case Series and Literature Review. Cancers, 2020, 12, 733.	3.7	17
16	Serum levels of miR-223 but not miR-21 are decreased in patients with neuroendocrine tumors. PLoS ONE, 2020, 15, e0244504.	2.5	3
17	A rare case of a patient with a high grade neuroendocrine tumor developing neutropenic sepsis after receiving PRRT combined with Capecitabine or Temozolomide: A case report. Molecular and Clinical Oncology, 2020, 14, 20.	1.0	4