

# Gorm Boje Jensen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6666605/publications.pdf>

Version: 2024-02-01

35  
papers

786  
citations

567281

15  
h-index

526287

27  
g-index

35  
all docs

35  
docs citations

35  
times ranked

1036  
citing authors

#	ARTICLE	IF	CITATIONS
1	QT Dispersion Has No Prognostic Information for Patients With Advanced Congestive Heart Failure and Reduced Left Ventricular Systolic Function. <i>Circulation</i> , 2001, 103, 831-835.	1.6	105
2	Elevated remnant cholesterol increases the risk of peripheral artery disease, myocardial infarction, and ischaemic stroke: a cohort-based study. <i>European Heart Journal</i> , 2022, 43, 3258-3269.	2.2	82
3	QTc Interval as a Guide to Select Those Patients With Congestive Heart Failure and Reduced Left Ventricular Systolic Function Who Will Benefit From Antiarrhythmic Treatment With Dofetilide. <i>Circulation</i> , 2001, 103, 1422-1427.	1.6	74
4	Clarithromycin for stable coronary heart disease increases all-cause and cardiovascular mortality and cerebrovascular morbidity over 10years in the CLARICOR randomised, blinded clinical trial. <i>International Journal of Cardiology</i> , 2015, 182, 459-465.	1.7	67
5	Normal values and reference ranges for left atrial strain by speckle-tracking echocardiography: the Copenhagen City Heart Study. <i>European Heart Journal Cardiovascular Imaging</i> , 2021, 23, 42-51.	1.2	47
6	Various Leisure-Time Physical Activities Associated With Widely Divergent Life Expectancies: The Copenhagen City Heart Study. <i>Mayo Clinic Proceedings</i> , 2018, 93, 1775-1785.	3.0	42
7	Impact of persistence and non-persistence in leisure time physical activity on coronary heart disease and all-cause mortality: The Copenhagen City Heart Study. <i>European Journal of Preventive Cardiology</i> , 2017, 24, 1615-1623.	1.8	41
8	Recovery of cardiac function following COVID-19: A prospective longitudinal cohort study. <i>European Journal of Heart Failure</i> , 2021, 23, 1903-1912.	7.1	40
9	10-Year Associations Between Tumor Necrosis Factor Receptors 1 and 2 and Cardiovascular Events in Patients With Stable Coronary Heart Disease: A CLARICOR (Effect of Clarithromycin on Mortality and) Association. 2018, 7, .	3.7	33
10	Usefulness of left atrial strain for predicting incident atrial fibrillation and ischaemic stroke in the general population. <i>European Heart Journal Cardiovascular Imaging</i> , 2022, 23, 363-371.	1.2	28
11	Increasing population height and risk of incident atrial fibrillation: the Copenhagen City Heart Study. <i>European Heart Journal</i> , 2018, 39, 4012-4019.	2.2	25
12	Risk stratification in stable coronary artery disease is possible at cardiac troponin levels below conventional detection and is improved by use of N-terminal pro-B-type natriuretic peptide. <i>European Journal of Preventive Cardiology</i> , 2014, 21, 1275-1284.	1.8	22
13	Cathepsin B and S as markers for cardiovascular risk and all-cause mortality in patients with stable coronary heart disease during 10 years: a CLARICOR trial sub-study. <i>Atherosclerosis</i> , 2018, 278, 97-102.	0.8	22
14	U-Shaped Association Between Duration of Sports Activities and Mortality: Copenhagen City Heart Study. <i>Mayo Clinic Proceedings</i> , 2021, 96, 3012-3020.	3.0	21
15	Normal Values for Myocardial Work Indices Derived From Pressure-Strain Loop Analyses: From the CCHS. <i>Circulation: Cardiovascular Imaging</i> , 2022, 15, 101161CIRCIMAGING121013712.	2.6	16
16	Prognostic Impact of Mild Hypokalemia in Terms of Death and Stroke in the General Population: A Prospective Population Study. <i>American Journal of Medicine</i> , 2018, 131, 318.e9-318.e19.	1.5	13
17	Predictors for major cardiovascular outcomes in stable ischaemic heart disease (PREMAC): statistical analysis plan for data originating from the CLARICOR (clarithromycin for patients with stable) Association. 2018, 7, .	3.7	33
18	Circulating endostatin as a risk factor for cardiovascular events in patients with stable coronary heart disease: A CLARICOR trial sub-study. <i>Atherosclerosis</i> , 2019, 284, 202-208.	0.8	11

#	ARTICLE	IF	CITATIONS
19	Associations between body mass index trajectories in childhood and cardiovascular risk factors in adulthood. <i>Atherosclerosis</i> , 2020, 314, 10-17.	0.8	11
20	Changes in left atrial structure and function over a decade in the general population. <i>European Heart Journal Cardiovascular Imaging</i> , 2021, 23, 124-136.	1.2	10
21	Association between Isometric and Allometric Height-Indexed Left Atrial Size and Atrial Fibrillation. <i>Journal of the American Society of Echocardiography</i> , 2022, 35, 141-150.e4.	2.8	10
22	AHRR (cg5575921) methylation safely improves specificity of lung cancer screening eligibility criteria: A cohort study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2022, , cebp.1059.2021.	2.5	10
23	Serum osteoprotegerin as a long-term predictor for patients with stable coronary artery disease and its association with diabetes and statin treatment: A CLARICOR trial 10-year follow-up substudy. <i>Atherosclerosis</i> , 2020, 301, 8-14.	0.8	9
24	Pregnancy Associated Plasma Protein-A as a Cardiovascular Risk Marker in Patients with Stable Coronary Heart Disease During 10 Years Follow-Upâ€”A CLARICOR Trial Sub-Study. <i>Journal of Clinical Medicine</i> , 2020, 9, 265.	2.4	7
25	Measures of left atrial function predict incident heart failure in a lowâ€”risk general population: the Copenhagen City Heart Study. <i>European Journal of Heart Failure</i> , 2021, , .	7.1	6
26	Carotid intima media thickness and ankle brachial index are inversely associated in subjects with and without diabetes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018, 78, 464-469.	1.2	5
27	Global and regional wall motion abnormalities and incident heart failure in the general population. <i>International Journal of Cardiology</i> , 2022, 357, 146-151.	1.7	5
28	The cardiac isovolumetric contraction time is an independent predictor of incident atrial fibrillation and adverse outcomes following first atrial fibrillation event in the general population. <i>European Heart Journal Cardiovascular Imaging</i> , 2020, 21, 49-57.	1.2	4
29	Is abdominal obesity at baseline influencing weight changes in observational studies and during weight loss interventions?. <i>American Journal of Clinical Nutrition</i> , 2018, 108, 913-921.	4.7	2
30	Progression in risk factors during 36 years of follow-up and prediction of carotid intima-media thickness in a large cohort of adults with and without diabetes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2020, 80, 491-499.	1.2	2
31	Prognostic value of 12 novel cardiological biomarkers in stable coronary artery disease. A 10-year follow-up of the placebo group of the Copenhagen CLARICOR trial. <i>BMJ Open</i> , 2020, 10, e033720.	1.9	2
32	Atrial cardiomyopathy in patients with ischaemic stroke: a cross-sectional and prospective cohort studyâ€”the COAST study. <i>BMJ Open</i> , 2022, 12, e061018.	1.9	2
33	0171â€”...Sedentary work and risk of venous thromboembolism. , 2017, , .		0
34	The Reply. <i>American Journal of Medicine</i> , 2018, 131, e169.	1.5	0
35	A screening method to spot biomarkers that may warn of serious events in a chronic diseaseâ€”illustrated by cardiological CLARICOR trial data. <i>Clinical Chemistry and Laboratory Medicine</i> , 2021, 59, 1852-1860.	2.3	0