

Berend Terluin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6675692/publications.pdf>

Version: 2024-02-01

97
papers

3,505
citations

172457

29
h-index

161849

54
g-index

113
all docs

113
docs citations

113
times ranked

3699
citing authors

#	ARTICLE	IF	CITATIONS
1	A confirmatory factor analysis approach was found to accurately estimate the reliability of transition ratings. <i>Journal of Clinical Epidemiology</i> , 2022, 141, 36-45.	5.0	16
2	Barriers and Facilitators for Return to Work from the Perspective of Workers with Common Mental Disorders with Short, Medium and Long-Term Sickness Absence: A Longitudinal Qualitative Study. <i>Journal of Occupational Rehabilitation</i> , 2022, 32, 272-283.	2.2	18
3	Estimating power for clinical trials with Patient Reported Outcomes - using Item Response Theory. <i>Journal of Clinical Epidemiology</i> , 2022, 141, 141-148.	5.0	1
4	A systematic survey identified methodological issues in studies estimating anchor-based minimal important differences in patient-reported outcomes. <i>Journal of Clinical Epidemiology</i> , 2022, 142, 144-151.	5.0	8
5	Determining the minimal important change of the 6-minute walking test in Multiple Sclerosis patients using a predictive modelling anchor-based method. <i>Multiple Sclerosis and Related Disorders</i> , 2022, 57, 103438.	2.0	4
6	Present state bias in transition ratings was accurately estimated in simulated and real data. <i>Journal of Clinical Epidemiology</i> , 2022, 143, 128-136.	5.0	7
7	Commentary on Riddle and Dumenci: LCA is no viable alternative to the MCID. <i>Osteoarthritis and Cartilage</i> , 2022, , .	1.3	0
8	Improved adjusted minimal important change took reliability of transition ratings into account. <i>Journal of Clinical Epidemiology</i> , 2022, 148, 48-53.	5.0	10
9	Assessing measurement equivalence of the Danish and Dutch Four-Dimensional Symptom Questionnaire using differential item and test functioning analysis. <i>Scandinavian Journal of Public Health</i> , 2021, 49, 479-486.	2.3	1
10	Which Oxford Knee Score level represents a satisfactory symptom state after undergoing a total knee replacement?. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2021, 92, 85-90.	3.3	23
11	Four-Dimensional Symptom Questionnaire (4DSQ)., 2021, , 1-3.		1
12	Validation of the Four-Dimensional Symptom Questionnaire (4DSQ) in a mental health setting. <i>Psychology, Health and Medicine</i> , 2021, 26, 1-19.	2.4	11
13	Assessing baseline dependency of anchor-based minimal important change (MIC): donâ€™t stratify on the baseline score!. <i>Quality of Life Research</i> , 2021, 30, 2773-2782.	3.1	11
14	Cutoff Values to Interpret Short-term Treatment Outcomes After Arthroscopic Meniscal Surgery, Measured With the Knee Injury and Osteoarthritis Outcome Score. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2021, 51, 281-288.	3.5	12
15	Minimal important change (MIC): a conceptual clarification and systematic review of MIC estimates of PROMIS measures. <i>Quality of Life Research</i> , 2021, 30, 2729-2754.	3.1	153
16	Tests for central sensitization in general practice: a Delphi study. <i>BMC Family Practice</i> , 2021, 22, 206.	2.9	8
17	Impact of symptom focusing and somatosensory amplification on persistent physical symptoms: A three-year follow-up study. <i>Journal of Psychosomatic Research</i> , 2020, 135, 110131.	2.6	10
18	Unlike ROC analysis, a new IRT method identified clinical thresholds unbiased by disease prevalence. <i>Journal of Clinical Epidemiology</i> , 2020, 124, 118-125.	5.0	12

#	ARTICLE	IF	CITATIONS
19	Measurement equivalence of the Four-Dimensional Symptom Questionnaire (4DSQ) in adolescents and emerging adults. <i>PLoS ONE</i> , 2019, 14, e0221904.	2.5	1
20	Psychometric properties of the Dutch-Flemish Patient-Reported Outcomes Measurement Information System Pain Behavior item bank in patients with musculoskeletal complaints. <i>Journal of Pain</i> , 2019, 20, 1328-1337.	1.4	6
21	Effectiveness of a tailored implementation strategy to improve adherence to a guideline on mental health problems in occupational health care. <i>BMC Health Services Research</i> , 2019, 19, 281.	2.2	7
22	Searching for the optimal number of response alternatives for the distress scale of the four-dimensional symptom questionnaire. <i>BMC Psychiatry</i> , 2019, 19, 103.	2.6	2
23	“Can I mean what is depression?” A qualitative exploration of UK general practitioners’ perceptions of distinctions between emotional distress and depressive disorder. <i>BMJ Open</i> , 2019, 9, e032644.	1.9	7
24	Validation of the Dutch-Flemish PROMIS Pain Interference Item Bank in Patients With Musculoskeletal Complaints. <i>Spine</i> , 2019, 44, 411-419.	2.0	6
25	Central sensitization in chronic pain and medically unexplained symptom research: A systematic review of definitions, operationalizations and measurement instruments. <i>Journal of Psychosomatic Research</i> , 2019, 117, 32-40.	2.6	100
26	Measurement Properties of the Barthel Index in Geriatric Rehabilitation. <i>Journal of the American Medical Directors Association</i> , 2019, 20, 420-425.e1.	2.5	115
27	Meaningful Change Scores in the Knee Injury and Osteoarthritis Outcome Score in Patients Undergoing Anterior Cruciate Ligament Reconstruction. <i>American Journal of Sports Medicine</i> , 2018, 46, 1120-1128.	4.2	72
28	Assessing the equivalence of Web-based and paper-and-pencil questionnaires using differential item and test functioning (DIF and DTF) analysis: a case of the Four-Dimensional Symptom Questionnaire (4DSQ). <i>Quality of Life Research</i> , 2018, 27, 1191-1200.	3.1	9
29	Use of a mental health guideline by occupational physicians and associations with return to work in workers sick-listed due to common mental disorders: a retrospective cohort study. <i>Disability and Rehabilitation</i> , 2018, 40, 2623-2631.	1.8	7
30	From non-adaptive depression to general distress. <i>Brain, Behavior, and Immunity</i> , 2018, 69, 629.	4.1	1
31	Cross-cultural validation of the German version of the Four-Dimensional Symptom Questionnaire (4DSQ) in multimorbid elderly people. <i>Quality of Life Research</i> , 2018, 27, 2691-2697.	3.1	7
32	Factors contributing to the recognition of anxiety and depression in general practice. <i>BMC Family Practice</i> , 2018, 19, 99.	2.9	14
33	Minimal important change values for the Oxford Knee Score and the Forgotten Joint Score at 1 year after total knee replacement. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018, 89, 541-547.	3.3	59
34	Longitudinal associations of multiple physical symptoms with recurrence of depressive and anxiety disorders. <i>Journal of Psychosomatic Research</i> , 2017, 97, 96-101.	2.6	12
35	The anchor-based minimal important change, based on receiver operating characteristic analysis or predictive modeling, may need to be adjusted for the proportion of improved patients. <i>Journal of Clinical Epidemiology</i> , 2017, 83, 90-100.	5.0	81
36	Systematic review of measurement properties of questionnaires measuring somatization in primary care patients. <i>Journal of Psychosomatic Research</i> , 2017, 103, 42-62.	2.6	29

#	ARTICLE	IF	CITATIONS
37	Effectiveness of an Intervention to Enhance Occupational Physiciansâ€™ Guideline Adherence on Sickness Absence Duration in Workers with Common Mental Disorders: A Cluster-Randomized Controlled Trial. <i>Journal of Occupational Rehabilitation</i> , 2017, 27, 559-567.	2.2	15
38	De 4DKL bevordert juist persoonsgerichte zorg. <i>Huisarts En Wetenschap</i> , 2017, 60, 115-115.	0.0	0
39	Identifying levels of general distress in first line mental health services: can GP- and eHealth clientsâ€™ scores be meaningfully compared?. <i>BMC Psychiatry</i> , 2017, 17, 382.	2.6	4
40	Applying Computerized Adaptive Testing to the Four-Dimensional Symptom Questionnaire (4DSQ): A Simulation Study. <i>JMIR Mental Health</i> , 2017, 4, e7.	3.3	3
41	The Four-Dimensional Symptom Questionnaire (4DSQ) in the general population: scale structure, reliability, measurement invariance and normative data: a cross-sectional survey. <i>Health and Quality of Life Outcomes</i> , 2016, 14, 130.	2.4	31
42	Validation of the four-dimensional symptom questionnaire (4DSQ) and prevalence of psychological symptoms in orthopaedic shoulder patients. <i>Journal of Orthopaedic Research</i> , 2016, 34, 683-691.	2.3	9
43	Measuring maternal mental health using the Dutch Four-Dimensional Symptom Questionnaire (4DSQ): Pregnancy-related item bias across the perinatal period. <i>Midwifery</i> , 2016, 40, 192-199.	2.3	1
44	Occupational physiciansâ€™ perceived barriers and suggested solutions to improve adherence to a guideline on mental health problems: analysis of a peer group training. <i>BMC Health Services Research</i> , 2016, 16, 271.	2.2	24
45	Medically unexplained physical symptoms and work functioning over 2 years: their association and the influence of depressive and anxiety disorders and job characteristics. <i>BMC Family Practice</i> , 2016, 17, 46.	2.9	16
46	Cross-cultural validation of the Turkish Four-Dimensional Symptom Questionnaire (4DSQ) using differential item and test functioning (DIF and DTF) analysis. <i>BMC Family Practice</i> , 2016, 17, 53.	2.9	9
47	The association between medically unexplained physical symptoms and health care use over two years and the influence of depressive and anxiety disorders and personality traits: a longitudinal study. <i>BMC Health Services Research</i> , 2016, 16, 100.	2.2	28
48	Differences in Connection Strength between Mental Symptoms Might Be Explained by Differences in Variance: Reanalysis of Network Data Did Not Confirm Staging. <i>PLoS ONE</i> , 2016, 11, e0155205.	2.5	108
49	Meetinstrumenten. , 2016, , 225-241.		0
50	Lessons learnt from a cluster-randomised trial evaluating the effectiveness of Self-Management Support (SMS) delivered by practice nurses in routine diabetes care. <i>BMJ Open</i> , 2015, 5, e007014.	1.9	17
51	Distinguishing between emotional distress and psychiatric disorder in primary care attenders: A cross sectional study of the four-dimensional symptom questionnaire (4DSQ). <i>Journal of Affective Disorders</i> , 2015, 184, 198-204.	4.1	16
52	Evaluating the validity of the French version of the Four-Dimensional Symptom Questionnaire with differential item functioning analysis. <i>Family Practice</i> , 2015, 32, cmv025.	1.9	6
53	Effectiveness of a tailored implementation programme to improve recognition, diagnosis and treatment of anxiety and depression in general practice: a cluster randomised controlled trial. <i>Implementation Science</i> , 2015, 10, 33.	6.9	32
54	Improving occupational physiciansâ€™ adherence to a practice guideline: feasibility and impact of a tailored implementation strategy. <i>BMC Medical Education</i> , 2015, 15, 82.	2.4	37

#	ARTICLE	IF	CITATIONS
55	Effect of an intervention to enhance guideline adherence of occupational physicians on return-to-work self-efficacy in workers sick-listed with common mental disorders. <i>BMC Public Health</i> , 2015, 15, 796.	2.9	33
56	Minimal important change (MIC) based on a predictive modeling approach was more precise than MIC based on ROC analysis. <i>Journal of Clinical Epidemiology</i> , 2015, 68, 1388-1396.	5.0	99
57	An international comparison of occupational health guidelines for the management of mental disorders and stress-related psychological symptoms. <i>Occupational and Environmental Medicine</i> , 2015, 72, 313-322.	2.8	31
58	Impact of Verb Tense on Response to the Four-Dimensional Symptom Questionnaire (4DSQ). <i>European Journal of Psychological Assessment</i> , 2015, 31, 202-210.	3.0	3
59	The English version of the four-dimensional symptom questionnaire (4DSQ) measures the same as the original Dutch questionnaire: A validation study. <i>European Journal of General Practice</i> , 2014, 20, 320-326.	2.0	23
60	Non-pharmacological interventions for somatoform disorders and medically unexplained physical symptoms (MUPS) in adults. <i>The Cochrane Library</i> , 2014, 2014, CD011142.	2.8	132
61	Evaluation of the measurement properties of the Manchester foot pain and disability index. <i>BMC Musculoskeletal Disorders</i> , 2014, 15, 276.	1.9	10
62	To what extent does the anxiety scale of the Four-Dimensional Symptom Questionnaire (4DSQ) detect specific types of anxiety disorder in primary care? A psychometric study. <i>BMC Psychiatry</i> , 2014, 14, 121.	2.6	28
63	Systematic tailoring for the implementation of guideline recommendations for anxiety and depressive disorders in general practice: perceived usefulness of tailored interventions. <i>BMC Family Practice</i> , 2013, 14, 94.	2.9	12
64	Integrating nurse-led Self-Management Support (SMS) in routine primary care: design of a hybrid effectiveness-implementation study among type 2 diabetes patients with problems of daily functioning and emotional distress: a study protocol. <i>BMC Family Practice</i> , 2013, 14, 77.	2.9	6
65	Effectiveness of guideline-based care by occupational physicians on the return-to-work of workers with common mental disorders: design of a cluster-randomised controlled trial. <i>BMC Public Health</i> , 2013, 13, 193.	2.9	20
66	Assessing psychological health in midwifery practice: A validation study of the Four-Dimensional Symptom Questionnaire (4DSQ), a Dutch primary care instrument. <i>Midwifery</i> , 2013, 29, 608-615.	2.3	13
67	Collaborative stepped care <i>vs</i> care as usual for common mental disorders: 8-month, cluster randomised controlled trial. <i>British Journal of Psychiatry</i> , 2013, 203, 132-139.	2.8	48
68	Evaluating the cross-cultural validity of the Polish version of the Four-Dimensional Symptom Questionnaire (4DSQ) using differential item functioning (DIF) analysis. <i>Family Practice</i> , 2012, 29, 609-615.	1.9	15
69	Mathematical coupling does not account for the association between baseline severity and minimally important change values. <i>Journal of Clinical Epidemiology</i> , 2012, 65, 355-357.	5.0	3
70	Recognition of anxiety disorders by family physicians after rigorous medical record case extraction. <i>General Hospital Psychiatry</i> , 2012, 34, 460-467.	2.4	26
71	Prevention of anxiety disorders in primary care: A feasibility study. <i>BMC Psychiatry</i> , 2012, 12, 206.	2.6	10
72	A validation study of the Four-Dimensional Symptom Questionnaire (4DSQ) in insurance medicine. <i>Work</i> , 2012, 43, 369-380.	1.1	8

#	ARTICLE	IF	CITATIONS
73	Randomised controlled trial of tailored interventions to improve the management of anxiety and depressive disorders in primary care. <i>Implementation Science</i> , 2011, 6, 75.	6.9	20
74	Psychological symptoms and subsequent sickness absence. <i>International Archives of Occupational and Environmental Health</i> , 2011, 84, 825-837.	2.3	23
75	Training GP's to use a minimal intervention for stress-related mental disorders with sick leave (MISS): Effects on performance. <i>Patient Education and Counseling</i> , 2010, 78, 206-211.	2.2	18
76	Cost-effectiveness of a minimal intervention for stress-related sick leave in general practice: Results of an economic evaluation alongside a pragmatic randomised control trial. <i>Journal of Affective Disorders</i> , 2010, 120, 177-187.	4.1	21
77	A workplace intervention for sick-listed employees with distress: results of a randomised controlled trial. <i>Occupational and Environmental Medicine</i> , 2010, 67, 596-602.	2.8	96
78	Three ways to quantify uncertainty in individually applied "minimally important change" values. <i>Journal of Clinical Epidemiology</i> , 2010, 63, 37-45.	5.0	113
79	The validity of the Dutch K10 and extended K10 screening scales for depressive and anxiety disorders. <i>Psychiatry Research</i> , 2010, 176, 45-50.	3.3	143
80	A Participatory Workplace Intervention for Employees With Distress and Lost Time: A Feasibility Evaluation Within a Randomized Controlled Trial. <i>Journal of Occupational Rehabilitation</i> , 2009, 19, 212-222.	2.2	67
81	Predicting Return to Work in Employees Sick-Listed Due to Minor Mental Disorders. <i>Journal of Occupational Rehabilitation</i> , 2009, 19, 323-332.	2.2	70
82	Detecting depressive and anxiety disorders in distressed patients in primary care; comparative diagnostic accuracy of the Four-Dimensional Symptom Questionnaire (4DSQ) and the Hospital Anxiety and Depression Scale (HADS). <i>BMC Family Practice</i> , 2009, 10, 58.	2.9	91
83	Understanding Ferguson's $\hat{\tau}$: time to say good-bye?. <i>Health and Quality of Life Outcomes</i> , 2009, 7, 38.	2.4	5
84	Cost-effectiveness of a workplace intervention for sick-listed employees with common mental disorders: design of a randomized controlled trial. <i>BMC Public Health</i> , 2008, 8, 12.	2.9	44
85	Cost-effectiveness of usual general practitioner care with or without antidepressant medication for patients with minor or mild-major depression. <i>Journal of Affective Disorders</i> , 2008, 111, 106-112.	4.1	19
86	Commentary: Good primary mental health care is simple. <i>British Journal of General Practice</i> , 2008, 58, 409-409.	1.4	1
87	A Cluster-Randomised Trial Evaluating an Intervention for Patients with Stress-Related Mental Disorders and Sick Leave in Primary Care. <i>PLOS Clinical Trials</i> , 2007, 2, e26.	3.5	65
88	Development of a workplace intervention for sick-listed employees with stress-related mental disorders: Intervention Mapping as a useful tool. <i>BMC Health Services Research</i> , 2007, 7, 127.	2.2	97
89	Clinical effectiveness of usual care with or without antidepressant medication for primary care patients with minor or mild-major depression: a randomized equivalence trial. <i>BMC Medicine</i> , 2007, 5, 36.	5.5	37
90	Effectiveness of an intervention to reduce sickness absence in patients with emotional distress or minor mental disorders: a randomized controlled effectiveness trial. <i>General Hospital Psychiatry</i> , 2006, 28, 223-229.	2.4	86

#	ARTICLE	IF	CITATIONS
91	The Four-Dimensional Symptom Questionnaire (4DSQ): a validation study of a multidimensional self-report questionnaire to assess distress, depression, anxiety and somatization. BMC Psychiatry, 2006, 6, 34.	2.6	369
92	Effectiveness of a Minimal Intervention for Stress-related mental disorders with Sick leave (MISS); study protocol of a cluster randomised controlled trial in general practice [ISRCTN43779641]. BMC Public Health, 2006, 6, 124.	2.9	20
93	The four-dimensional symptom questionnaire (4DSQ): measuring distress and other mental health problems in a working population. Work and Stress, 2004, 18, 187-207.	4.5	125
94	The prognosis of minor depression in the general population: a systematic review. General Hospital Psychiatry, 2004, 26, 453-462.	2.4	62
95	Weg met de poortwachter!. Huisarts En Wetenschap, 2003, 46, 804-805.	0.0	1
96	Reliability and validity of the assessment of depression in general practice: the short depression interview (SDI). General Hospital Psychiatry, 2002, 24, 396-405.	2.4	14
97	Interpretation threshold values for the Oxford Knee Score in patients undergoing unicompartmental knee arthroplasty. Monthly Notices of the Royal Astronomical Society: Letters, 0, 93, 634-642.	3.3	6