

Kyoung-Im Cho

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6703643/publications.pdf>

Version: 2024-02-01

19
papers

217
citations

1040056

9
h-index

1058476

14
g-index

20
all docs

20
docs citations

20
times ranked

419
citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of duration and dosage of statin treatment and epicardial fat thickness on the recurrence of atrial fibrillation after electrical cardioversion. <i>Heart and Vessels</i> , 2015, 30, 490-497.	1.2	25
2	Impact of Gender on the Association of Epicardial Fat Thickness, Obesity, and Circadian Blood Pressure Pattern in Hypertensive Patients. <i>Journal of Diabetes Research</i> , 2015, 2015, 1-10.	2.3	24
3	Effects of High-Intensity Interval Training Versus Moderate-Intensity Continuous Training on Epicardial Fat Thickness and Endothelial Function in Hypertensive Metabolic Syndrome. <i>Metabolic Syndrome and Related Disorders</i> , 2020, 18, 96-102.	1.3	22
4	Effects of exergaming in postmenopausal women with high cardiovascular risk: A randomized controlled trial. <i>Clinical Cardiology</i> , 2020, 43, 363-370.	1.8	20
5	Impact of periodontitis as representative of chronic inflammation on long-term clinical outcomes in patients with atrial fibrillation. <i>Open Heart</i> , 2018, 5, e000708.	2.3	18
6	Simvastatin Attenuates the Oxidative Stress, Endothelial Thrombogenicity and the Inducibility of Atrial Fibrillation in a Rat Model of Ischemic Heart Failure. <i>International Journal of Molecular Sciences</i> , 2014, 15, 14803-14818.	4.1	14
7	Epicardial Fat Thickness and Free Fatty Acid Level are Predictors of Acute Ischemic Stroke with Atrial Fibrillation. <i>Journal of Cardiovascular Imaging</i> , 2018, 26, 65.	0.7	14
8	Exergaming Improves Executive Functions in Patients With Metabolic Syndrome: Randomized Controlled Trial. <i>JMIR Serious Games</i> , 2019, 7, e13575.	3.1	14
9	The Impact of Subclinical Hypothyroidism or Thyroid Autoimmunity on Coronary Vasospasm in Patients without Associated Cardiovascular Risk Factors. <i>Korean Circulation Journal</i> , 2015, 45, 125.	1.9	12
10	Attenuation of Acetylcholine Activated Potassium Current (IKACH) by Simvastatin, Not Pravastatin in Mouse Atrial Cardiomyocyte: Possible Atrial Fibrillation Preventing Effects of Statin. <i>PLoS ONE</i> , 2014, 9, e106570.	2.5	10
11	Association between the N-terminal plasma brain natriuretic peptide levels or elevated left ventricular filling pressure and thromboembolic risk in patients with non-valvular atrial fibrillation. <i>Journal of Cardiology</i> , 2016, 68, 110-116.	1.9	9
12	Association among epicardial fat, heart rate recovery and circadian blood pressure variability in patients with hypertension. <i>Clinical Hypertension</i> , 2015, 21, 24.	2.0	8
13	Heart failure awareness in the Korean general population: Results from the nationwide survey. <i>PLoS ONE</i> , 2019, 14, e0222264.	2.5	8
14	Heart rate recovery and blood pressure response during exercise testing in patients with microvascular angina. <i>Clinical Hypertension</i> , 2019, 25, 4.	2.0	8
15	Cardiac Involvement in Hypereosinophilia Associated with Toxocariasis. <i>Journal of Cardiovascular Imaging</i> , 2014, 22, 224.	0.8	4
16	Factors Associated with Low Awareness of Heart Failure in the General Population of Korea. <i>Korean Circulation Journal</i> , 2020, 50, 586.	1.9	4
17	Blunted heart rate recovery is associated with coronary artery spasm in patients with suspected vasospastic angina. <i>Clinical Hypertension</i> , 2017, 23, 24.	2.0	2
18	Fibroblasts, electrophysiological changes, and atrial fibrillation: Focus on the arrhythmogenic properties of fibroblasts. <i>Journal of Arrhythmia</i> , 2013, 29, 201-203.	1.2	1

#	ARTICLE	IF	CITATIONS
19	Effect of a fixed-dose combination of Telmisartan/S-amlodipine on circadian blood pressure compared with Telmisartan monotherapy: TENUVA-BP study. <i>Clinical Hypertension</i> , 2022, 28, 7.	2.0	0