
Kyoung-Im Cho

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6703643/publications.pdf Version: 2024-02-01



Куонис-Ім Сно

#	Article	IF	CITATIONS
1	Effect of a fixed-dose combination of Telmisartan/S-amlodipine on circadian blood pressure compared with Telmisartan monotherapy: TENUVA-BP study. Clinical Hypertension, 2022, 28, 7.	2.0	0
2	Effects of exergaming in postmenopausal women with high cardiovascular risk: A randomized controlled trial. Clinical Cardiology, 2020, 43, 363-370.	1.8	20
3	Effects of High-Intensity Interval Training Versus Moderate-Intensity Continuous Training on Epicardial Fat Thickness and Endothelial Function in Hypertensive Metabolic Syndrome. Metabolic Syndrome and Related Disorders, 2020, 18, 96-102.	1.3	22
4	Factors Associated with Low Awareness of Heart Failure in the General Population of Korea. Korean Circulation Journal, 2020, 50, 586.	1.9	4
5	Heart failure awareness in the Korean general population: Results from the nationwide survey. PLoS ONE, 2019, 14, e0222264.	2.5	8
6	Heart rate recovery and blood pressure response during exercise testing in patients with microvascular angina. Clinical Hypertension, 2019, 25, 4.	2.0	8
7	Exergaming Improves Executive Functions in Patients With Metabolic Syndrome: Randomized Controlled Trial. JMIR Serious Games, 2019, 7, e13575.	3.1	14
8	Epicardial Fat Thickness and Free Fatty Acid Level are Predictors of Acute Ischemic Stroke with Atrial Fibrillation. Journal of Cardiovascular Imaging, 2018, 26, 65.	0.7	14
9	Impact of periodontitis as representative of chronic inflammation on long-term clinical outcomes in patients with atrial fibrillation. Open Heart, 2018, 5, e000708.	2.3	18
10	Blunted heart rate recovery is associated with coronary artery spasm in patients with suspected vasospastic angina. Clinical Hypertension, 2017, 23, 24.	2.0	2
11	Association between the N-terminal plasma brain natriuretic peptide levels or elevated left ventricular filling pressure and thromboembolic risk in patients with non-valvular atrial fibrillation. Journal of Cardiology, 2016, 68, 110-116.	1.9	9
12	Association among epicardial fat, heart rate recovery and circadian blood pressure variability in patients with hypertension. Clinical Hypertension, 2015, 21, 24.	2.0	8
13	The Impact of Subclinical Hypothyroidism or Thyroid Autoimmunity on Coronary Vasospasm in Patients without Associated Cardiovascular Risk Factors. Korean Circulation Journal, 2015, 45, 125.	1.9	12
14	Impact of Gender on the Association of Epicardial Fat Thickness, Obesity, and Circadian Blood Pressure Pattern in Hypertensive Patients. Journal of Diabetes Research, 2015, 2015, 1-10.	2.3	24
15	Impact of duration and dosage of statin treatment and epicardial fat thickness on the recurrence of atrial fibrillation after electrical cardioversion. Heart and Vessels, 2015, 30, 490-497.	1.2	25
16	Cardiac Involvement in Hypereosinophilia Associated with Toxocariasis. Journal of Cardiovascular Imaging, 2014, 22, 224.	0.8	4
17	Attenuation of Acetylcholine Activated Potassium Current (IKACh) by Simvastatin, Not Pravastatin in Mouse Atrial Cardiomyocyte: Possible Atrial Fibrillation Preventing Effects of Statin. PLoS ONE, 2014, 9, e106570.	2.5	10
18	Simvastatin Attenuates the Oxidative Stress, Endothelial Thrombogenicity and the Inducibility of Atrial Fibrillation in a Rat Model of Ischemic Heart Failure. International Journal of Molecular Sciences, 2014, 15, 14803-14818.	4.1	14

#	Article	IF	CITATIONS
19	Fibroblasts, electrophysiological changes, and atrial fibrillation: Focus on the arrhythmogenic properties of fibroblasts. Journal of Arrhythmia, 2013, 29, 201-203.	1.2	1