Hiroshi Hashizume

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6732828/publications.pdf

Version: 2024-02-01

172457 161849 3,402 107 29 54 citations h-index g-index papers 121 121 121 3184 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Detailed Subphenotyping of Lumbar Modic Changes and Their Association with Low Back Pain in a Large Population-Based Study: The Wakayama Spine Study. Pain and Therapy, 2022, 11, 57-71.	3.2	12
2	Relationship Between Age-Related Spinopelvic Sagittal Alignment and Low Back Pain in Adults of Population-Based Cohorts: The ROAD Study. Journal of Pain Research, 2022, Volume 15, 33-38.	2.0	4
3	The discrepancy between radiographically-assessed and self-recognized hallux valgus in a large population-based cohort. BMC Musculoskeletal Disorders, 2022, 23, 31.	1.9	5
4	Sacroiliac Joint Pain Should Be Suspected In Early Buttock And Groin Pain After Adult Spinal Deformity Surgery: An Observational Study. Spine Surgery and Related Research, 2022, , .	0.7	0
5	Long-Term Outcomes after Selective Microendoscopic Laminotomy for Multilevel Lumbar Spinal Stenosis with and without Remaining Radiographic Stenosis: A 10-Year Follow-Up Study. Spine Surgery and Related Research, 2022, 6, 488-496.	0.7	2
6	Is radiographic lumbar spinal stenosis associated with the quality of life?: The Wakayama Spine Study. PLoS ONE, 2022, 17, e0263930.	2.5	1
7	Lateral interbody release for fused vertebrae via transpsoas approach in adult spinal deformity surgery: a preliminary report of radiographic and clinical outcomes. BMC Musculoskeletal Disorders, 2022, 23, 245.	1.9	O
8	A Prospective, 3-year Longitudinal Study of Modic Changes of the Lumbar Spine in a Population-based Cohort. Spine, 2022, 47, 490-497.	2.0	8
9	Lumbar Fusion including Sacroiliac Joint Fixation Increases the Stress and Angular Motion at the Hip Joint: A Finite Element Study. Spine Surgery and Related Research, 2022, 6, 681-688.	0.7	5
10	A novel technique using ultrasonography in upper airway management after anterior cervical decompression and fusion. BMC Medical Imaging, 2022, 22, 67.	2.7	0
11	Psychometric Evaluation and External Validity of the Japanese Version of Lumbar Stiffness Disability Index. Spine Surgery and Related Research, 2022, , .	0.7	O
12	Association between types of Modic changes in the lumbar region and low back pain in a large cohort: the Wakayama spine study. European Spine Journal, 2021, 30, 1011-1017.	2.2	30
13	Prevalence and associated factors of pistol grip deformity in Japanese local residents. Scientific Reports, 2021, 11, 6025.	3.3	3
14	Does prophylactic use of topical gelatin-thrombin matrix sealant affect postoperative drainage volume and hematoma formation following microendoscopic spine surgery? A randomized controlled trial. Spine Journal, 2021, 21, 446-454.	1.3	4
15	Long-term Clinical Outcomes of Microendoscopic Laminotomy for Cervical Spondylotic Myelopathy. Clinical Spine Surgery, 2021, 34, 383-390.	1.3	5
16	Kitchen elbow sign predicts surgical outcomes in adults with spinal deformity: a retrospective cohort study. Scientific Reports, 2021, 11, 12859.	3.3	6
17	Association between subjective oral dysfunction and locomotive syndrome in community-dwelling older adults. Scientific Reports, 2021, 11, 12591.	3.3	7
18	Adjacent segment disease on hip joint as a complication of spinal fusion surgery including sacroiliac joint fixation. European Spine Journal, 2021, 30, 1314-1319.	2.2	16

#	Article	IF	Citations
19	Lateral lumbar interbody fusion after reduction using the percutaneous pedicle screw system in the lateral position for Meyerding grade II spondylolisthesis: a preliminary report of a new lumbar reconstruction strategy. BMC Musculoskeletal Disorders, 2021, 22, 17.	1.9	3
20	The beneficial effect of physical activity on cognitive function in community-dwelling older persons with locomotive syndrome. PeerJ, 2021, 9, e12292.	2.0	0
21	Relationship of sagittal spinal alignment with low back pain and physical performance in the general population. Scientific Reports, 2021, 11, 20604.	3.3	13
22	Improving effect of microendoscopic decompression surgery on low back pain in patients with lumbar spinal stenosis and predictive factors of postoperative residual low back pain: a single-center retrospective study. BMC Musculoskeletal Disorders, 2021, 22, 954.	1.9	6
23	The incidence and risk factors for adjacent vertebral fractures in community-dwelling people with prevalent vertebral fracture: the 3rd and 4th survey of the ROAD study. Archives of Osteoporosis, 2020, 15, 74.	2.4	2
24	Lumbar high-intensity zones on MRI: imaging biomarkers for severe, prolonged low back pain and sciatica in a population-based cohort. Spine Journal, 2020, 20, 1025-1034.	1.3	26
25	Could automated machine-learned MRI grading aid epidemiological studies of lumbar spinal stenosis? Validation within the Wakayama spine study. BMC Musculoskeletal Disorders, 2020, 21, 158.	1.9	16
26	Prevalence of cervical anterior and posterior spondylolisthesis and its association with degenerative cervical myelopathy in a general population. Scientific Reports, 2020, 10, 10455.	3.3	9
27	Microendoscopic decompression for lumbosacral foraminal stenosis: a novel surgical strategy based on anatomical considerations using 3D image fusion with MRI/CT. Journal of Neurosurgery: Spine, 2020, , 1-7.	1.7	8
28	A short-segment fusion strategy using a wide-foot-plate expandable cage for vertebral pseudarthrosis after an osteoporotic vertebral fracture. Journal of Neurosurgery: Spine, 2020, 33, 862-869.	1.7	12
29	Physical performance decreases in the early stage of cervical myelopathy before the myelopathic signs appear: the Wakayama Spine Study. European Spine Journal, 2019, 28, 1217-1224.	2.2	7
30	Factory and construction work is associated with an increased risk of severe lumbar spinal stenosis on MRI: A case control analysis within the wakayama spine study. American Journal of Industrial Medicine, 2019, 62, 430-438.	2.1	4
31	Microendoscopic Decompression for Lumbar Spinal Stenosis With Degenerative Spondylolisthesis. Clinical Spine Surgery, 2019, 32, E20-E26.	1.3	22
32	<p>Prevalence of Facet Effusion and Its Relationship with Lumbar Spondylolisthesis and Low Back Pain: The Wakayama Spine Study</p> . Journal of Pain Research, 2019, Volume 12, 3521-3528.	2.0	14
33	Is radiographic lumbar spondylolisthesis associated with occupational exposures? Findings from a nested case control study within the Wakayama spine study. BMC Musculoskeletal Disorders, 2019, 20, 618.	1.9	8
34	Differences in prevalence and associated factors between mild and severe vertebral fractures in Japanese men and women: the third survey of the ROAD study. Journal of Bone and Mineral Metabolism, 2019, 37, 844-853.	2.7	25
35	Japanese orthopaedic association cervical myelopathy evaluation questionnaire (JOACMEQ): Part 5. Determination of responsiveness. Journal of Orthopaedic Science, 2019, 24, 57-61.	1.1	9
36	Diffuse idiopathic skeletal hyperostosis is associated with lumbar spinal stenosis requiring surgery. Journal of Bone and Mineral Metabolism, 2019, 37, 118-124.	2.7	29

#	Article	IF	CITATIONS
37	Local Sagittal Alignment of the Lumbar Spine and Range of Motion in 627 Asymptomatic Subjects: Age-Related Changes and Sex-Based Differences. Asian Spine Journal, 2019, 13, 663-671.	2.0	11
38	Serum levels of matrix metalloproteinase-3 and autoantibodies related to rheumatoid arthritis in the general Japanese population and their association with osteoporosis and osteoarthritis: the ROAD study. Journal of Bone and Mineral Metabolism, 2018, 36, 246-253.	2.7	6
39	Spinal Endoscopy for Delayed-Onset Lumbar Radiculopathy Resulting from Foraminal Stenosis after Osteoporotic Vertebral Fracture: A Case Report of a New Surgical Strategy. Case Reports in Orthopedics, 2018, 2018, 1-4.	0.3	5
40	Early versus delayed kyphoplasty for thoracolumbar osteoporotic vertebral fractures: The effect of timing on clinical and radiographic outcomes and subsequent compression fractures. Clinical Neurology and Neurosurgery, 2018, 173, 176-181.	1.4	37
41	Factors associated with lumbar spinal stenosis in a large-scale, population-based cohort: The Wakayama Spine Study. PLoS ONE, 2018, 13, e0200208.	2.5	19
42	Progression, incidence, and risk factors for intervertebral disc degeneration in a longitudinal population-based cohort: the Wakayama Spine Study. Osteoarthritis and Cartilage, 2017, 25, 1122-1131.	1.3	93
43	Minimally invasive decompression surgery for lumbar spinal stenosis with degenerative scoliosis: Predictive factors of radiographic and clinical outcomes. Journal of Orthopaedic Science, 2017, 22, 377-383.	1.1	31
44	Association of Lumbar Spondylolisthesis With Low Back Pain and Symptomatic Lumbar Spinal Stenosis in a Population-based Cohort. Spine, 2017, 42, E666-E671.	2.0	18
45	The prevalence of tandem spinal stenosis and its characteristics in a population-based MRI study: The Wakayama Spine Study. European Spine Journal, 2017, 26, 2529-2535.	2.2	26
46	Verification of the sensitivity of functional scores for treatment results – Substantial clinical benefit thresholds for the Japanese Orthopaedic Association Back Pain Evaluation Questionnaire (JOABPEQ). Journal of Orthopaedic Science, 2017, 22, 665-669.	1.1	20
47	Microendoscopic laminotomy versus conventional laminoplasty for cervical spondylotic myelopathy: 5-year follow-up study. Journal of Neurosurgery: Spine, 2017, 27, 403-409.	1.7	28
48	Long-term survival case of malignant glomus tumor mimicking "dumbbell-shaped―neurogenic tumor. European Spine Journal, 2017, 26, 42-46.	2.2	7
49	Cognitive impairment associated with locomotive syndrome in community-dwelling elderly women in Japan. Clinical Interventions in Aging, 2017, Volume 12, 1451-1457.	2.9	25
50	The Relationship between Locomotive Syndrome and Depression in Community-Dwelling Elderly People. Current Gerontology and Geriatrics Research, 2017, 2017, 1-6.	1.6	19
51	MRI-defined paraspinal muscle morphology in Japanese population: The Wakayama Spine Study. PLoS ONE, 2017, 12, e0187765.	2.5	65
52	Sagittal spino-pelvic alignment in adults: The Wakayama Spine Study. PLoS ONE, 2017, 12, e0178697.	2.5	89
53	Locomotive syndrome is associated with body composition and cardiometabolic disorders in elderly Japanese women. BMC Geriatrics, 2016, 16, 166.	2.7	19
54	Metabolic Syndrome Components Are Associated with Intervertebral Disc Degeneration: The Wakayama Spine Study. PLoS ONE, 2016, 11, e0147565.	2.5	40

#	Article	IF	CITATIONS
55	The Association between the Cross-Sectional Area of the Dural Sac and Low Back Pain in a Large Population: The Wakayama Spine Study. PLoS ONE, 2016, 11, e0160002.	2.5	16
56	Classification of High Intensity Zones of the Lumbar Spine and Their Association with Other Spinal MRI Phenotypes: The Wakayama Spine Study. PLoS ONE, 2016, 11, e0160111.	2.5	30
57	Response to "Revisiting the nomenclature and grading schemes for disc degeneration: issues to be solvedâ€. Spine Journal, 2015, 15, 2595-2596.	1.3	O
58	Physical Performance Measures Associated With Locomotive Syndrome in Middle-Aged and Older Japanese Women. Journal of Geriatric Physical Therapy, 2015, 38, 202-207.	1.1	25
59	Efficacy of Posterior Segmental Decompression Surgery for Pincer Mechanism in Cervical Spondylotic Myelopathy. Spine, 2015, 40, 1807-1815.	2.0	16
60	Clinical outcomes after microendoscopic laminotomy for lumbar spinal stenosis: a 5-year follow-up study. European Spine Journal, 2015, 24, 396-403.	2.2	40
61	Improved accuracy of diagnosis of lumbar intra and/or extra-foraminal stenosis by use of three-dimensional MR imaging: comparison with conventional MR imaging. Journal of Orthopaedic Science, 2015, 20, 287-294.	1.1	25
62	The association of combination of disc degeneration, end plate signal change, and Schmorl node with low back pain in a large population study: the Wakayama Spine Study. Spine Journal, 2015, 15, 622-628.	1.3	91
63	Japanese orthopaedic association back pain evaluation questionnaire (JOABPEQ) as an outcome measure for patients with low back pain: reference values in healthy volunteers. Journal of Orthopaedic Science, 2015, 20, 264-280.	1.1	43
64	<i>In Vivo</i> Patch-Clamp Analysis of the Antinociceptive Actions of TRPA1 Activation in the Spinal Dorsal Horn. Molecular Pain, 2015, 11, s12990-015-0021.	2.1	20
65	Development of a support tool for the clinical diagnosis of symptomatic lumbar intra-and/or extra†foraminal stenosis. Journal of Orthopaedic Science, 2015, 20, 811-817.	1.1	8
66	Radiographic natural course of lumbar degenerative spondylolisthesis and its risk factors related to the progression and onset in a 15-year community-based cohort study: the Miyama study. Journal of Orthopaedic Science, 2015, 20, 978-984.	1.1	26
67	Augmentation of motor evoked potentials using multi-train transcranial electrical stimulation in intraoperative neurophysiologic monitoring during spinal surgery. Journal of Clinical Monitoring and Computing, 2015, 29, 35-39.	1.6	23
68	Prevalence of diffuse idiopathic skeletal hyperostosis (DISH) of the whole spine and its association with lumbar spondylosis and knee osteoarthritis: the ROAD study. Journal of Bone and Mineral Metabolism, 2015, 33, 221-229.	2.7	82
69	The prevalence of cervical myelopathy among subjects with narrow cervical spinal canal in a population-based magnetic resonance imaging study: the Wakayama Spine Study. Spine Journal, 2014, 14, 2811-2817.	1.3	42
70	An outcome measure for patients with cervical myelopathy: the Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire (JOACMEQ): an average score of healthy volunteers. Journal of Orthopaedic Science, 2014, 19, 33-48.	1.1	16
71	Development and evaluation of a video exercise program for locomotive syndrome in the elderly. Modern Rheumatology, 2014, 24, 250-257.	1.8	12
72	Skipping breakfast and less exercise are risk factors for bone loss in young Japanese adults: a 3-year follow-up study. Journal of Bone and Mineral Metabolism, 2014, 32, 420-427.	2.7	3

#	Article	IF	CITATIONS
73	Prevalence and distribution of intervertebral disc degeneration over the entire spine in a population-based cohort: the Wakayama Spine Study. Osteoarthritis and Cartilage, 2014, 22, 104-110.	1.3	339
74	A New Electrophysiological Method for the Diagnosis of Extraforaminal Stenosis at L5-S1. Asian Spine Journal, 2014, 8, 145.	2.0	15
75	Quantification of the proportion of motor neurons recruited by transcranial electrical stimulation during intraoperative motor evoked potential monitoring. Journal of Clinical Monitoring and Computing, 2013, 27, 633-637.	1.6	11
76	Can decompression surgery relieve low back pain in patients with lumbar spinal stenosis combined with degenerative lumbar scoliosis?. European Spine Journal, 2013, 22, 2010-2014.	2.2	29
77	Consuming breakfast and exercising longer during high school increases bone mineral density in young adult men. Journal of Bone and Mineral Metabolism, 2013, 31, 329-336.	2.7	8
78	Endoscope-assisted spinal decompression surgery for lumbar spinal stenosis. Journal of Neurosurgery: Spine, 2013, 19, 664-671.	1.7	74
79	Development and evaluation of a video exercise program for locomotive syndrome in the elderly. Modern Rheumatology, 2013, , 1.	1.8	0
80	Efficacy of Novel Minimally Invasive Surgery Using Spinal Microendoscope For Treating Extraforaminal Stenosis at the Lumbosacral Junction. Journal of Spinal Disorders and Techniques, 2012, 25, 268-276.	1.9	28
81	Prevalence of Cervical Cord Compression and Its Association With Physical Performance in a Population-Based Cohort in Japan. Spine, 2012, 37, 1892-1898.	2.0	48
82	Distinct Degree of Radiculopathy at Different Levels of Peripheral Nerve Injury. Molecular Pain, 2012, 8, 1744-8069-8-31.	2.1	6
83	APrunus mumeExtract Stimulated the Proliferation and Differentiation of Osteoblastic MC3T3-E1 Cells. Bioscience, Biotechnology and Biochemistry, 2011, 75, 1907-1911.	1.3	21
84	A New 3-Dimensional Computed Tomography Imaging Method to Diagnose Extraforaminal Stenosis at the Lumbosacral Junction. Journal of Spinal Disorders and Techniques, 2010, 23, e47-e52.	1.9	17
85	Sarpogrelate Hydrochloride, a 5-HT2A Receptor Antagonist, Attenuates Neurogenic Pain Induced by Nucleus Pulposus in Rats. Spine, 2007, 32, 315-320.	2.0	17
86	Safety of Epidural Administration of Osteogenic Protein-1 (OP-1/BMP-7). Spine, 2007, 32, 1388-1393.	2.0	9
87	The Effects of Bone Morphogenetic Protein and Basic Fibroblast Growth Factor on Cultured Mesenchymal Stem Cells for Spine Fusion. Spine, 2007, 32, 1067-1071.	2.0	43
88	Osteogenic Protein-1 (Osteogenic Protein-1/Bone Morphogenetic Protein-7) Inhibits Degeneration and Pain-Related Behavior Induced by Chronically Compressed Nucleus Pulposus in the Rat. Spine, 2005, 30, 1933-1939.	2.0	56
89	The Use of Cultured Bone Marrow Cells in Type I Collagen Gel and Porous Hydroxyapatite for Posterolateral Lumbar Spine Fusion. Spine, 2005, 30, 1134-1138.	2.0	106
90	Experimental study of carriers of bone morphogenetic protein used for spinal fusion. Journal of Orthopaedic Science, 2004, 9, 142-151.	1.1	22

#	Article	IF	CITATIONS
91	Comparison of neuropathic pain induced by the application of normal and mechanically compressed nucleus pulposus to lumbar nerve roots in the rat. Journal of Orthopaedic Research, 2003, 21, 535-539.	2.3	39
92	A Clinical Case of Endoscopically Assisted Anterior Screw Fixation for the Type II Odontoid Fracture. Spine, 2003, 28, E102-E105.	2.0	29
93	Lumbar Sagittal Balance Influences the Clinical Outcome After Decompression and Posterolateral Spinal Fusion for Degenerative Lumbar Spondylolisthesis. Spine, 2002, 27, 59-64.	2.0	154
94	Epidural injection of cyclooxygenase-2 inhibitor attenuates pain-related behavior following application of nucleus pulposus to the nerve root in the rat. Journal of Orthopaedic Research, 2002, 20, 376-381.	2.3	37
95	Evaluation of Carriers of Bone Morphogenetic Protein for Spinal Fusion. Spine, 2001, 26, 933-939.	2.0	81
96	The use of sintered bone in spinal surgery. European Spine Journal, 2001, 10, S185-S188.	2.2	9
97	An Experimental Approach to Spinal Fusion Using Sintered Bovine Bone in a Pig Model. Journal of Spinal Disorders, 2000, 13, 156-164.	1.1	3
98	Spinal Glial Activation and Cytokine Expression After Lumbar Root Injury in the Rat. Spine, 2000, 25, 1206-1217.	2.0	241
99	Mechanical compression of the lumbar nerve root alters pain-related behaviors induced by the nucleus pulposus in the rat. Journal of Orthopaedic Research, 2000, 18, 257-264.	2.3	37
100	Cyclooxygenase-2 inhibitor SC-236 attenuates mechanical allodynia following nerve root injury in rats. Journal of Orthopaedic Research, 2000, 18, 977-982.	2.3	17
101	Central administration of methotrexate reduces mechanical allodynia in an animal model of radiculopathy/sciatica. Pain, 2000, 87, 159-169.	4.2	75
102	Effects of Basic Fibroblast Growth Factor on Spontaneous Resorption of Herniated Intervertebral Discs. Spine, 1999, 24, 940-945.	2.0	52
103	Experimental Spinal Fusion Using Sintered Bovine Bone Coated With Type I Collagen and Recombinant Human Bone Morphogenetic Protein-2. Spine, 1999, 24, 1863.	2.0	46
104	Effects of Steroid and Lipopolysaccharide on Spontaneous Resorption of Herniated Intervertebral Discs. Spine, 1998, 23, 870-876.	2.0	74
105	The Role of Phospholipase A2 and Nitric Oxide in Pain-Related Behavior Produced by an Allograft of Intervertebral Disc Material to the Sciatic Nerve of the Rat. Spine, 1997, 22, 1074-1079.	2.0	78
106	Histochemical Demonstration of Nitric Oxide in Herniated Lumbar Discs. Spine, 1997, 22, 1080-1084.	2.0	28
107	Pathomechanism of Pain-Related Behavior Produced by Allografts of Intervertebral Disc in the Rat. Spine, 1996, 21, 2101-2107.	2.0	156