

Roger Schneider

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6751333/publications.pdf>

Version: 2024-02-01

93
papers

7,355
citations

94433

37
h-index

58581

82
g-index

99
all docs

99
docs citations

99
times ranked

8909
citing authors

#	ARTICLE	IF	CITATIONS
1	The surface of lipid droplets constitutes a barrier for endoplasmic reticulum-resident integral membrane proteins. <i>Journal of Cell Science</i> , 2022, 135, .	2.0	13
2	Lipid droplets form a network interconnected by the endoplasmic reticulum through which their proteins equilibrate. <i>Journal of Cell Science</i> , 2022, 135, .	2.0	13
3	Prostate secretory protein 94 inhibits sterol binding and export by the mammalian CAP protein CRISP2 in a calcium-sensitive manner. <i>Journal of Biological Chemistry</i> , 2022, 298, 101600.	3.4	4
4	Seipin collaborates with the ER membrane to control the sites of lipid droplet formation. <i>Current Opinion in Cell Biology</i> , 2022, 75, 102070.	5.4	11
5	Pre-existing bilayer stresses modulate triglyceride accumulation in the ER versus lipid droplets. <i>ELife</i> , 2021, 10, .	6.0	55
6	Seipin accumulates and traps diacylglycerols and triglycerides in its ring-like structure. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	7.1	52
7	A Unique Junctional Interface at Contact Sites Between the Endoplasmic Reticulum and Lipid Droplets. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 650186.	3.7	23
8	Retinyl esters form lipid droplets independently of triacylglycerol and seipin. <i>Journal of Cell Biology</i> , 2021, 220, .	5.2	22
9	Membrane-Interacting DNA Nanotubes Induce Cancer Cell Death. <i>Nanomaterials</i> , 2021, 11, 2003.	4.1	8
10	The yeast cell wall protein Pry3 inhibits mating through highly conserved residues within the CAP domain. <i>Biology Open</i> , 2020, 9, .	1.2	5
11	Mitochondrial sphingosine-1-phosphate lyase is essential for phosphatidylethanolamine synthesis and survival of <i>Trypanosoma brucei</i> . <i>Scientific Reports</i> , 2020, 10, 8268.	3.3	8
12	Seipin and Nem1 establish discrete ER subdomains to initiate yeast lipid droplet biogenesis. <i>Journal of Cell Biology</i> , 2020, 219, .	5.2	68
13	Lipid droplet biogenesis from specialized ER subdomains. <i>Microbial Cell</i> , 2020, 7, 218-221.	3.2	11
14	<i>Necator americanus</i> Ancylostoma Secreted Protein-2 (Na-ASP-2) Binds an Ascaroside (ascr#3) in Its Fatty Acid Binding Site. <i>Frontiers in Chemistry</i> , 2020, 8, 608296.	3.6	2
15	Crystal structure of <i>Brugia malayi</i> venom allergen-like protein-1 (BmVAL-1), a vaccine candidate for lymphatic filariasis. <i>International Journal for Parasitology</i> , 2018, 48, 371-378.	3.1	17
16	Architecture of Lipid Droplets in Endoplasmic Reticulum Is Determined by Phospholipid Intrinsic Curvature. <i>Current Biology</i> , 2018, 28, 915-926.e9.	3.9	148
17	<i>Heligmosomoides polygyrus</i> Venom Allergen-like Protein-4 (HpVAL-4) is a sterol binding protein. <i>International Journal for Parasitology</i> , 2018, 48, 359-369.	3.1	18
18	The function of yeast <sc>CAP</sc> family proteins in lipid export, mating, and pathogen defense. <i>FEBS Letters</i> , 2018, 592, 1304-1311.	2.8	18

#	ARTICLE	IF	CITATIONS
19	Secreted venom allergen-like proteins of helminths: Conserved modulators of host responses in animals and plants. <i>PLoS Pathogens</i> , 2018, 14, e1007300.	4.7	41
20	Localization and functional characterization of the pathogenesis-related proteins Rbe1p and Rbt4p in <i>Candida albicans</i> . <i>PLoS ONE</i> , 2018, 13, e0201932.	2.5	3
21	The pathogen-related yeast protein Pry1, a member of the CAP protein superfamily, is a fatty acid-binding protein. <i>Journal of Biological Chemistry</i> , 2017, 292, 8304-8314.	3.4	40
22	Crystal Structure of MpPR-1i, a SCP/TAPS protein from <i>Moniliophthora perniciosa</i> , the fungus that causes Witches'™ Broom Disease of Cacao. <i>Scientific Reports</i> , 2017, 7, 7818.	3.3	11
23	Sphingolipid accumulation causes mitochondrial dysregulation and cell death. <i>Cell Death and Differentiation</i> , 2017, 24, 2044-2053.	11.2	38
24	Crystal Structure of <i>Borrelia turicatae</i> protein, BTA121, a differentially regulated gene in the tick-mammalian transmission cycle of relapsing fever spirochetes. <i>Scientific Reports</i> , 2017, 7, 15310.	3.3	2
25	Plant pathogenesis-related proteins of the cacao fungal pathogen <i>Moniliophthora perniciosa</i> differ in their lipid-binding specificities. <i>Journal of Biological Chemistry</i> , 2017, 292, 20558-20569.	3.4	18
26	The sterol-binding activity of PATHOGENESIS-RELATED PROTEIN 1 reveals the mode of action of an antimicrobial protein. <i>Plant Journal</i> , 2017, 89, 502-509.	5.7	156
27	A Ligand-Binding Assay to Measure the Affinity and Specificity of Sterol-Binding Proteins In Vitro. <i>Methods in Molecular Biology</i> , 2017, 1645, 361-368.	0.9	6
28	Chemical crosslinking and mass spectrometry to elucidate the topology of integral membrane proteins. <i>PLoS ONE</i> , 2017, 12, e0186840.	2.5	11
29	Chemogenetic E-MAP in <i>Saccharomyces cerevisiae</i> for Identification of Membrane Transporters Operating Lipid Flip Flop. <i>PLoS Genetics</i> , 2016, 12, e1006160.	3.5	6
30	Structural and functional characterization of the CAP domain of pathogen-related yeast 1 (Pry1) protein. <i>Scientific Reports</i> , 2016, 6, 28838.	3.3	30
31	Tools for the analysis of metabolic flux through the sphingolipid pathway. <i>Biochimie</i> , 2016, 130, 76-80.	2.6	6
32	Following the flux of long-chain bases through the sphingolipid pathway in vivo using mass spectrometry. <i>Journal of Lipid Research</i> , 2016, 57, 906-915.	4.2	10
33	Mature lipid droplets are accessible to ER luminal proteins. <i>Journal of Cell Science</i> , 2016, 129, 3803-3815.	2.0	42
34	Accumulation of long-chain bases in yeast promotes their conversion to a long-chain base vinyl ether. <i>Journal of Lipid Research</i> , 2016, 57, 2040-2050.	4.2	3
35	Valproate Induces the Unfolded Protein Response by Increasing Ceramide Levels. <i>Journal of Biological Chemistry</i> , 2016, 291, 22253-22261.	3.4	20
36	Cholesterol-Binding by the Yeast CAP Family Member Pry1 Requires the Presence of an Aliphatic Side Chain on Cholesterol. <i>Journal of Steroids & Hormonal Science</i> , 2016, 7, .	0.1	9

#	ARTICLE	IF	CITATIONS
37	Expression of perilipin 5 promotes lipid droplet formation in yeast. <i>Communicative and Integrative Biology</i> , 2015, 8, e1071728.	1.4	6
38	Yeast Integral Membrane Proteins Apq12, Brl1, and Brr6 Form a Complex Important for Regulation of Membrane Homeostasis and Nuclear Pore Complex Biogenesis. <i>Eukaryotic Cell</i> , 2015, 14, 1217-1227.	3.4	36
39	Match-making for posaconazole through systems thinking. <i>Trends in Parasitology</i> , 2015, 31, 46-51.	3.3	9
40	TORC1 Regulates Pah1 Phosphatidate Phosphatase Activity via the Nem1/Spo7 Protein Phosphatase Complex. <i>PLoS ONE</i> , 2014, 9, e104194.	2.5	53
41	The Natural Diyne-Furan Fatty Acid EV-086 Is an Inhibitor of Fungal Delta-9 Fatty Acid Desaturation with Efficacy in a Model of Skin Dermatophytosis. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 455-466.	3.2	12
42	<i>Schistosoma mansoni</i> venom allergen-like protein 4 (SmVAL4) is a novel lipid-binding SCP/TAPS protein that lacks the prototypical CAP motifs. <i>Acta Crystallographica Section D: Biological Crystallography</i> , 2014, 70, 2186-2196.	2.5	36
43	The caveolin-binding motif of the pathogen-related yeast protein Pry1, a member of the CAP protein superfamily, is required for in vivo export of cholesteryl acetate. <i>Journal of Lipid Research</i> , 2014, 55, 883-894.	4.2	35
44	Expression of oleosin and perilipins in yeast promote formation of lipid droplets from the endoplasmic reticulum. <i>Journal of Cell Science</i> , 2013, 126, 5198-209.	2.0	90
45	The CAP protein superfamily: function in sterol export and fungal virulence. <i>Biomolecular Concepts</i> , 2013, 4, 519-525.	2.2	61
46	A Novel Sit4 Phosphatase Complex Is Involved in the Response to Ceramide Stress in Yeast. <i>Oxidative Medicine and Cellular Longevity</i> , 2013, 2013, 1-9.	4.0	6
47	Pathogen-Related Yeast (PRY) proteins and members of the CAP superfamily are secreted sterol-binding proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 16882-16887.	7.1	112
48	Mechanisms of sterol uptake and transport in yeast. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2012, 129, 70-78.	2.5	89
49	Regulation of sphingolipid synthesis via Orm1 and Orm2 in yeast. <i>Journal of Cell Science</i> , 2012, 125, 2428-35.	2.0	77
50	Lipid droplets are functionally connected to the endoplasmic reticulum in <i>Saccharomyces cerevisiae</i>. <i>Journal of Cell Science</i> , 2011, 124, 2424-2437.	2.0	356
51	The topology of the triacylglycerol synthesizing enzyme Lro1 indicates that neutral lipids can be produced within the luminal compartment of the endoplasmic reticulum: Implications for the biogenesis of lipid droplets. <i>Communicative and Integrative Biology</i> , 2011, 4, 781-784.	1.4	40
52	Integral membrane proteins Brr6 and Apq12 link assembly of the nuclear pore complex to lipid homeostasis in the endoplasmic reticulum. <i>Journal of Cell Science</i> , 2010, 123, 141-151.	2.0	72
53	Ypk1, the yeast orthologue of the human serum- and glucocorticoid-induced kinase, is required for efficient uptake of fatty acids. <i>Journal of Cell Science</i> , 2010, 123, 2218-2227.	2.0	33
54	Integrating complex functions. <i>Nucleus</i> , 2010, 1, 387-392.	2.2	12

#	ARTICLE	IF	CITATIONS
55	Orm1 and Orm2 are conserved endoplasmic reticulum membrane proteins regulating lipid homeostasis and protein quality control. Proceedings of the National Academy of Sciences of the United States of America, 2010, 107, 5851-5856.	7.1	245
56	The Cdc42 Effectors Ste20, Cla4, and Skm1 Down-Regulate the Expression of Genes Involved in Sterol Uptake by a Mitogen-activated Protein Kinase-independent Pathway. Molecular Biology of the Cell, 2009, 20, 4826-4837.	2.1	22
57	Membrane Rafts Are Involved in Intracellular Miconazole Accumulation in Yeast Cells. Journal of Biological Chemistry, 2009, 284, 32680-32685.	3.4	31
58	Mitochondrial Outer Membrane Proteins Assist Bid in Bax-mediated Lipidic Pore Formation. Molecular Biology of the Cell, 2009, 20, 2276-2285.	2.1	107
59	Monitoring Sterol Uptake, Acetylation, and Export in Yeast. , 2009, 580, 221-232.		2
60	Lipid signalling in disease. Nature Reviews Molecular Cell Biology, 2008, 9, 162-176.	37.0	1,091
61	Lipid-dependent surface transport of the proton pumping ATPase: A model to study plasma membrane biogenesis in yeast. Biochimie, 2007, 89, 249-254.	2.6	43
62	Intracellular sterol transport in eukaryotes, a connection to mitochondrial function?. Biochimie, 2007, 89, 255-259.	2.6	23
63	An acetylation/deacetylation cycle controls the export of sterols and steroids from <i>S. cerevisiae</i> . EMBO Journal, 2007, 26, 5109-5119.	7.8	87
64	The role of lipids in the biogenesis of integral membrane proteins. Applied Microbiology and Biotechnology, 2007, 73, 1224-1232.	3.6	22
65	Extraction of Yeast Lipids. , 2006, 313, 041-046.		43
66	Analysis of Yeast Lipids. , 2006, 313, 075-084.		37
67	A two-step method for the introduction of single or multiple defined point mutations into the genome of <i>Saccharomyces cerevisiae</i> . Yeast, 2006, 23, 825-831.	1.7	46
68	A Genomewide Screen Reveals a Role of Mitochondria in Anaerobic Uptake of Sterols in Yeast. Molecular Biology of the Cell, 2006, 17, 90-103.	2.1	75
69	Yeh1 Constitutes the Major Steryl Ester Hydrolase under Heme-Deficient Conditions in <i>Saccharomyces cerevisiae</i> . Eukaryotic Cell, 2006, 5, 1018-1025.	3.4	31
70	Very Long-chain Fatty Acid-containing Lipids rather than Sphingolipids per se Are Required for Raft Association and Stable Surface Transport of Newly Synthesized Plasma Membrane ATPase in Yeast. Journal of Biological Chemistry, 2006, 281, 34135-34145.	3.4	79
71	The <i>Saccharomyces cerevisiae</i> YLL012/YEH1, YLR020/YEH2, and TGL1 Genes Encode a Novel Family of Membrane-Anchored Lipases That Are Required for Steryl Ester Hydrolysis. Molecular and Cellular Biology, 2005, 25, 1655-1668.	2.3	124
72	Synthesis of Sphingolipids with Very Long Chain Fatty Acids but Not Ergosterol Is Required for Routing of Newly Synthesized Plasma Membrane ATPase to the Cell Surface of Yeast. Journal of Biological Chemistry, 2005, 280, 22515-22522.	3.4	86

#	ARTICLE	IF	CITATIONS
73	Differential Regulation of Ceramide Synthase Components LAC1 and LAG1 in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2004, 3, 880-892.	3.4	60
74	Identification and biophysical characterization of a very-long-chain-fatty-acid-substituted phosphatidylinositol in yeast subcellular membranes. <i>Biochemical Journal</i> , 2004, 381, 941-949.	3.7	85
75	Acyl-CoA-binding protein, Acb1p, is required for normal vacuole function and ceramide synthesis in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2004, 380, 907-918.	3.7	73
76	The Sur7p Family Defines Novel Cortical Domains in <i>Saccharomyces cerevisiae</i> , Affects Sphingolipid Metabolism, and Is Involved in Sporulation. <i>Molecular and Cellular Biology</i> , 2002, 22, 927-934.	2.3	112
77	Lipid-dependent Subcellular Relocalization of the Acyl Chain Desaturase in Yeast. <i>Molecular Biology of the Cell</i> , 2002, 13, 4429-4442.	2.1	31
78	A Specific Structural Requirement for Ergosterol in Long-chain Fatty Acid Synthesis Mutants Important for Maintaining Raft Domains in Yeast. <i>Molecular Biology of the Cell</i> , 2002, 13, 4414-4428.	2.1	112
79	Bid, Bax, and Lipids Cooperate to Form Supramolecular Openings in the Outer Mitochondrial Membrane. <i>Cell</i> , 2002, 111, 331-342.	28.9	1,337
80	Roles of Phosphatidylethanolamine and of Its Several Biosynthetic Pathways in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2001, 12, 997-1007.	2.1	245
81	Depletion of Acyl-Coenzyme A-Binding Protein Affects Sphingolipid Synthesis and Causes Vesicle Accumulation and Membrane Defects in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2001, 12, 1147-1160.	2.1	128
82	Molecular Identification of <i>virilizer</i> , a Gene Required for the Expression of the Sex-Determining Gene <i>Sex-lethal</i> in <i>Drosophila melanogaster</i> . <i>Genetics</i> , 2001, 157, 679-688.	2.9	34
83	A Novel Cold-Sensitive Allele of the Rate-Limiting Enzyme of Fatty Acid Synthesis, Acetyl Coenzyme A Carboxylase, Affects the Morphology of the Yeast Vacuole through Acylation of Vac8p. <i>Molecular and Cellular Biology</i> , 2000, 20, 2984-2995.	2.3	37
84	Elo1p-Dependent Carboxy-Terminal Elongation of C14:1 ⁿ 9 to C16:1 ⁿ 11 Fatty Acids in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 2000, 182, 3655-3660.	2.2	83
85	Brave little yeast, please guide us to Thebes: sphingolipid function in <i>S. cerevisiae</i> . <i>BioEssays</i> , 1999, 21, 1004-1010.	2.5	57
86	Electrospray Ionization Tandem Mass Spectrometry (Esi-MS/MS) Analysis of the Lipid Molecular Species Composition of Yeast Subcellular Membranes Reveals Acyl Chain-Based Sorting/Remodeling of Distinct Molecular Species En Route to the Plasma Membrane. <i>Journal of Cell Biology</i> , 1999, 146, 741-754.	5.2	449
87	The <i>Saccharomyces cerevisiae</i> Hyperrecombination Mutant <i>hpr1</i> Is Synthetically Lethal with Two Conditional Alleles of the Acetyl Coenzyme A Carboxylase Gene and Causes a Defect in Nuclear Export of Polyadenylated RNA. <i>Molecular and Cellular Biology</i> , 1999, 19, 3415-3422.	2.3	55
88	Brave little yeast, please guide us to Thebes: sphingolipid function in <i>S. cerevisiae</i> . <i>BioEssays</i> , 1999, 21, 1004-1010.	2.5	1
89	Organelle Structure, Function, and Inheritance in Yeast: A Role for Fatty Acid Synthesis?. <i>Cell</i> , 1997, 88, 431-434.	28.9	83
90	The yeast <i>mic2</i> mutant is defective in the formation of mannosyl-diinositolphosphorylceramide1. <i>FEBS Letters</i> , 1997, 411, 211-214.	2.8	30

#	ARTICLE	IF	CITATIONS
91	Phospholipids: synthesis, sorting, subcellular traffic - the yeast approach. Trends in Cell Biology, 1996, 6, 260-266.	7.9	29
92	The nuclear GTPase cycle: promoting peripheralization?. Trends in Cell Biology, 1995, 5, 5-8.	7.9	14
93	Lipid Droplet Biogenesis is Driven by Liquid-Liquid Phase Separation. SSRN Electronic Journal, 0, , .	0.4	10