

Naoko Miyagawa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6773180/publications.pdf>

Version: 2024-02-01

60
papers

1,067
citations

394421

19
h-index

454955

30
g-index

62
all docs

62
docs citations

62
times ranked

1821
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Association between C-Reactive Protein Levels and Functional Disability in the General Older-Population: The Takashima Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2023, 30, 56-65. | 2.0 | 1 |
| 2 | Association between Stress-Coping Strategy and Functional Disability in the General Older Adult Population: The Takashima Study. <i>Gerontology</i> , 2022, 68, 699-706. | 2.8 | 3 |
| 3 | A genome-wide association study on adherence to low-carbohydrate diets in Japanese. <i>European Journal of Clinical Nutrition</i> , 2022, , . | 2.9 | 1 |
| 4 | Dietary Patterns and Their Associations with Intermediate Age-Related Macular Degeneration in a Japanese Population. <i>Journal of Clinical Medicine</i> , 2022, 11, 1617. | 2.4 | 2 |
| 5 | Differential Association of Serum n-3 Polyunsaturated Fatty Acids with Various Cerebrovascular Lesions in Japanese Men. <i>Cerebrovascular Diseases</i> , 2022, 51, 774-780. | 1.7 | 0 |
| 6 | Population-Based Impact of Smoking, Drinking, and Genetic Factors on HDL-Cholesterol Levels in J-MICC Study Participants. <i>Journal of Epidemiology</i> , 2021, , . | 2.4 | 0 |
| 7 | Alcohol drinking and brain morphometry in apparently healthy community-dwelling Japanese men. <i>Alcohol</i> , 2021, 90, 57-65. | 1.7 | 6 |
| 8 | Reproducibility and validity of food group intake in a short food frequency questionnaire for the middle-aged Japanese population. <i>Environmental Health and Preventive Medicine</i> , 2021, 26, 28. | 3.4 | 29 |
| 9 | Association between the prevalence of hypertension and dairy consumption by housing type among survivors of the Great East Japan Earthquake. <i>Journal of Human Hypertension</i> , 2021, , . | 2.2 | 2 |
| 10 | Factors Associated with Lower Cognitive Performance Scores Among Older Japanese Men in Hawaii and Japan. <i>Journal of Alzheimer's Disease</i> , 2021, 81, 403-412. | 2.6 | 1 |
| 11 | Association between socioeconomic status and prolonged television viewing time in a general Japanese population: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2021, 26, 57. | 3.4 | 3 |
| 12 | Association between socioeconomic status and physical inactivity in a general Japanese population: NIPPON DATA2010. <i>PLoS ONE</i> , 2021, 16, e0254706. | 2.5 | 5 |
| 13 | Association between Lifestyle Changes and at-Home Hours during and after the State of Emergency Due to the COVID-19 Pandemic in Japan. <i>Nutrients</i> , 2021, 13, 2698. | 4.1 | 19 |
| 14 | Relationships of Alcohol Consumption with Coronary Risk Factors and Macro- and Micro-Nutrient Intake in Japanese People: The INTERLIPID Study. <i>Journal of Nutritional Science and Vitaminology</i> , 2021, 67, 28-38. | 0.6 | 2 |
| 15 | Proteinuria and Reduced Estimated Glomerular Filtration Rate are Independently Associated With Lower Cognitive Abilities in Apparently Healthy Community-Dwelling Elderly Men in Japan: A Cross-sectional Study. <i>Journal of Epidemiology</i> , 2020, 30, 244-252. | 2.4 | 10 |
| 16 | Elevated Fasting Blood Glucose Levels Are Associated With Lower Cognitive Function, With a Threshold in Non-Diabetic Individuals: A Population-Based Study. <i>Journal of Epidemiology</i> , 2020, 30, 121-127. | 2.4 | 9 |
| 17 | Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80. <i>European Journal of Clinical Nutrition</i> , 2020, 74, 67-76. | 2.9 | 17 |
| 18 | Alcohol consumption and cognitive function in elderly Japanese men. <i>Alcohol</i> , 2020, 85, 145-152. | 1.7 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Association of Red Meat Intake with the Risk of Cardiovascular Mortality in General Japanese Stratified by Kidney Function: NIPPON DATA80. <i>Nutrients</i> , 2020, 12, 3707. | 4.1 | 4 |
| 20 | Relationship Between Step Counts and Cerebral Small Vessel Disease in Japanese Men. <i>Stroke</i> , 2020, 51, 3584-3591. | 2.0 | 19 |
| 21 | Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , 2019, 29, 133-138. | 2.4 | 9 |
| 22 | Association of blood levels of marine omega-3 fatty acids with coronary calcification and calcium density in Japanese men. <i>European Journal of Clinical Nutrition</i> , 2019, 73, 783-792. | 2.9 | 22 |
| 23 | Cardiovascular Risk Assessment Chart by Dietary Factors in Japan. NIPPON DATA80. <i>Circulation Journal</i> , 2019, 83, 1254-1260. | 1.6 | 11 |
| 24 | Associations of Nutrient Patterns with the Prevalence of Metabolic Syndrome: Results from the Baseline Data of the Japan Multi-Institutional Collaborative Cohort Study. <i>Nutrients</i> , 2019, 11, 990. | 4.1 | 24 |
| 25 | The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 417-423. | 2.3 | 8 |
| 26 | The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. <i>Journal of Hypertension</i> , 2019, 37, 1676-1681. | 0.5 | 7 |
| 27 | Factors associated with intra-individual visit-to-visit variability of blood pressure in four countries: the INTERMAP study. <i>Journal of Human Hypertension</i> , 2019, 33, 229-236. | 2.2 | 7 |
| 28 | Having few remaining teeth is associated with a low nutrient intake and low serum albumin levels in middle-aged and older Japanese individuals: findings from the NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019, 24, 1. | 3.4 | 84 |
| 29 | Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality in a 15-Year Follow-Up Study of the General Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 198-206. | 2.0 | 17 |
| 30 | Association of Alcohol Consumption With Fat Deposition in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Epidemiology</i> , 2019, 29, 205-212. | 2.4 | 9 |
| 31 | Overweight or underweight and the risk of decline in activities of daily living in a 22-year cohort study of a Japanese sample. <i>Geriatrics and Gerontology International</i> , 2018, 18, 799-805. | 1.5 | 8 |
| 32 | Serum magnesium, phosphorus, and calcium levels and subclinical calcific aortic valve disease: A population-based study. <i>Atherosclerosis</i> , 2018, 273, 145-152. | 0.8 | 27 |
| 33 | Dietary tofu intake and long-term risk of death from stroke in a general population. <i>Clinical Nutrition</i> , 2018, 37, 182-188. | 5.0 | 10 |
| 34 | Relationships among Food Group Intakes, Household Expenditure, and Education Attainment in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S23-S28. | 2.4 | 10 |
| 35 | Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S29-S34. | 2.4 | 15 |
| 36 | Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S35-S39. | 2.4 | 2 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Associations of High-Density Lipoprotein Particle and High-Density Lipoprotein Cholesterol With Alcohol Intake, Smoking, and Body Mass Indexâ€”The INTERLIPID Study â€”. <i>Circulation Journal</i> , 2018, 82, 2557-2565. | 1.6 | 18 |
| 38 | Effectiveness of a Self-monitoring Device for Urinary Sodium-to-Potassium Ratio on Dietary Improvement in Free-Living Adults: a Randomized Controlled Trial. <i>Journal of Epidemiology</i> , 2018, 28, 41-47. | 2.4 | 30 |
| 39 | Relationship of serum irisin levels to prevalence and progression of coronary artery calcification: A prospective, population-based study. <i>International Journal of Cardiology</i> , 2018, 267, 177-182. | 1.7 | 30 |
| 40 | Food sources of dietary sodium in the Japanese adult population: the international study of macro-/micronutrients and blood pressure (INTERMAP). <i>European Journal of Nutrition</i> , 2017, 56, 1269-1280. | 3.9 | 20 |
| 41 | Significant inverse association of equol-producer status with coronary artery calcification but not dietary isoflavones in healthy Japanese men. <i>British Journal of Nutrition</i> , 2017, 117, 260-266. | 2.3 | 31 |
| 42 | Comparison of weighed food record procedures for the reference methods in two validation studies of food frequency questionnaires. <i>Journal of Epidemiology</i> , 2017, 27, 331-337. | 2.4 | 7 |
| 43 | Overall nutrient and total fat intake among Japanese people: The INTERLIPID Study Japan. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2017, 26, 837-848. | 0.4 | 3 |
| 44 | Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. <i>Journal of the American Heart Association</i> , 2016, 5, . | 3.7 | 39 |
| 45 | Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Preventive Medicine Reports</i> , 2016, 4, 225-232. | 1.8 | 30 |
| 46 | Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. <i>International Journal of Cardiology</i> , 2016, 220, 262-267. | 1.7 | 29 |
| 47 | Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , 2016, 246, 141-147. | 0.8 | 48 |
| 48 | Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia. LOX-1 ligand and IMT in Japanese. <i>Journal of Clinical Lipidology</i> , 2016, 10, 172-180.e1. | 1.5 | 11 |
| 49 | Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , 2016, 10, 1195-1202.e1. | 1.5 | 12 |
| 50 | Relationship of three different types of low-carbohydrate diet to cardiometabolic risk factors in a Japanese population: the INTERMAP/INTERLIPID Study. <i>European Journal of Nutrition</i> , 2016, 55, 1515-1524. | 3.9 | 12 |
| 51 | High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , 2015, 239, 444-450. | 0.8 | 18 |
| 52 | Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980â€”2010. <i>Hypertension Research</i> , 2015, 38, 790-795. | 2.7 | 39 |
| 53 | Six random specimens of daytime casual urine on different days are sufficient to estimate daily sodium/potassium ratio in comparison to 7-day 24-h urine collections. <i>Hypertension Research</i> , 2014, 37, 765-771. | 2.7 | 56 |
| 54 | Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. <i>British Journal of Nutrition</i> , 2014, 112, 916-924. | 2.3 | 59 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: A 24-year follow-up of Japanese general population. <i>Journal of Cardiology</i> , 2014, 64, 218-224. | 1.9 | 11 |
| 56 | Impact of Metabolic Syndrome on the Risk of Cardiovascular Disease Mortality in the United States and in Japan. <i>American Journal of Cardiology</i> , 2014, 113, 84-89. | 1.6 | 69 |
| 57 | Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. <i>Atherosclerosis</i> , 2014, 236, 237-243. | 0.8 | 22 |
| 58 | Long-chain n-3 polyunsaturated fatty acids intake and cardiovascular disease mortality risk in Japanese: A 24-year follow-up of NIPPON DATA80. <i>Atherosclerosis</i> , 2014, 232, 384-389. | 0.8 | 51 |
| 59 | Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. <i>Heart</i> , 2013, 99, 1024-1029. | 2.9 | 7 |
| 60 | Seven-year incidence of new-onset hypertension by frequency of dairy intake among survivors of the Great East Japan Earthquake. <i>Hypertension Research</i> , 0, , . | 2.7 | 1 |