

Rosa Marina Melillo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6813681/publications.pdf>

Version: 2024-02-01

87
papers

6,622
citations

61984

43
h-index

62596

80
g-index

124
all docs

124
docs citations

124
times ranked

7001
citing authors

#	ARTICLE	IF	CITATIONS
1	In PD-1+ human colon cancer cells NIVOLUMAB promotes survival and could protect tumor cells from conventional therapies. , 2022, 10, e004032.		25
2	The Impact of Resolution of Inflammation on Tumor Microenvironment: Exploring New Ways to Control Cancer Progression. Cancers, 2022, 14, 3333.	3.7	6
3	PD-1 blockade delays tumor growth by inhibiting an intrinsic SHP2/Ras/MAPK signalling in thyroid cancer cells. Journal of Experimental and Clinical Cancer Research, 2021, 40, 22.	8.6	37
4	Toll-Like Receptor 7 Mediates Inflammation Resolution and Inhibition of Angiogenesis in Non-Small Cell Lung Cancer. Cancers, 2021, 13, 740.	3.7	8
5	AXL Is a Novel Predictive Factor and Therapeutic Target for Radioactive Iodine Refractory Thyroid Cancer. Cancers, 2019, 11, 785.	3.7	27
6	Recent advances in understanding immune phenotypes of thyroid carcinomas: prognostication and emerging therapies. F1000Research, 2019, 8, 227.	1.6	20
7	RET-mediated modulation of tumor microenvironment and immune response in multiple endocrine neoplasia type 2 (MEN2). Endocrine-Related Cancer, 2018, 25, T105-T119.	3.1	19
8	New perspectives in cancer: Modulation of lipid metabolism and inflammation resolution. Pharmacological Research, 2018, 128, 80-87.	7.1	31
9	Formyl peptide receptor 1 suppresses gastric cancer angiogenesis and growth by exploiting inflammation resolution pathways. Oncoimmunology, 2017, 6, e1293213.	4.6	43
10	Signal Transducer and Activator of Transcription 1 Plays a Pivotal Role in RET/PTC3 Oncogene-induced Expression of Indoleamine 2,3-Dioxygenase 1. Journal of Biological Chemistry, 2017, 292, 1785-1797.	3.4	17
11	Interleukin-8, but Not the Related Chemokine CXCL1, Sustains an Autocrine Circuit Necessary for the Properties and Functions of Thyroid Cancer Stem Cells. Stem Cells, 2017, 35, 135-146.	3.2	40
12	Editorial: Novel Mechanism of Radioactive Iodine Refractivity in Thyroid Cancer. Journal of the National Cancer Institute, 2017, 109, .	6.3	11
13	Multiple anti-tumor effects of Reparixin on thyroid cancer. Oncotarget, 2017, 8, 35946-35961.	1.8	22
14	FRT â€œ FONDATION RENE TOURAINE. Experimental Dermatology, 2015, 24, 803-820.	2.9	0
15	The RET Receptor Family. , 2015, , 559-591.		1
16	Mast cells induce epithelial-to-mesenchymal transition and stem cell features in human thyroid cancer cells through an IL-8â€œAktâ€œSlug pathway. Oncogene, 2015, 34, 5175-5186.	5.9	176
17	Formyl peptide receptors at the interface of inflammation, angiogenesis and tumor growth. Pharmacological Research, 2015, 102, 184-191.	7.1	97
18	The genomic landscape of papillary thyroid carcinoma. Nature Reviews Endocrinology, 2015, 11, 133-134.	9.6	12

#	ARTICLE	IF	CITATIONS
19	The formyl peptide receptor 1 exerts a tumor suppressor function in human gastric cancer by inhibiting angiogenesis. <i>Oncogene</i> , 2015, 34, 3826-3838.	5.9	69
20	AXL is an oncotarget in human colorectal cancer. <i>Oncotarget</i> , 2015, 6, 23281-23296.	1.8	55
21	Indoleamine 2,3-Dioxygenase 1 (IDO1) Is Up-Regulated in Thyroid Carcinoma and Drives the Development of an Immunosuppressant Tumor Microenvironment. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, E832-E840.	3.6	73
22	Molecular Mechanism of 17-Allylamino-17-demethoxygeldanamycin (17-AAG)-induced AXL Receptor Tyrosine Kinase Degradation. <i>Journal of Biological Chemistry</i> , 2013, 288, 17481-17494.	3.4	44
23	Serum soluble ST2 and interleukin-33 levels in patients with pulmonary arterial hypertension. <i>International Journal of Cardiology</i> , 2013, 168, 1545-1547.	1.7	50
24	CXCR4 expression correlates with the degree of tumor infiltration and BRAF status in papillary thyroid carcinomas. <i>Modern Pathology</i> , 2012, 25, 46-55.	5.5	35
25	Molecular Biomarkers in Thyroid FNA Samples. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, 4370-4373.	3.6	18
26	RET: A Multi-Faceted Gene in Human Cancer. <i>Endocrinology and Metabolism</i> , 2012, 27, 173.	3.0	1
27	Inflammation in thyroid oncogenesis. <i>American Journal of Cancer Research</i> , 2012, 2, 286-97.	1.4	25
28	Activation of TYRO3/AXL Tyrosine Kinase Receptors in Thyroid Cancer. <i>Cancer Research</i> , 2011, 71, 1792-1804.	0.9	87
29	Higher Intratumoral Expression of CD1a, Tryptase, and CD68 in a Follicular Variant of Papillary Thyroid Carcinoma Compared to Adenomas: Correlation with Clinical and Pathological Parameters. <i>Thyroid</i> , 2011, 21, 1209-1215.	4.5	39
30	Mast cells have a protumorigenic role in human thyroid cancer. <i>Oncogene</i> , 2010, 29, 6203-6215.	5.9	190
31	Thyroid cancer and inflammation. <i>Molecular and Cellular Endocrinology</i> , 2010, 321, 94-102.	3.2	186
32	CXC Chemokine Receptor 4 Immunodetection in the Follicular Variant of Papillary Thyroid Carcinoma: Comparison to Galectin-3 and Hector Battifora Mesothelial Cell-1. <i>Thyroid</i> , 2010, 20, 495-504.	4.5	24
33	Differential diagnosis of thyroid nodules using fine-needle aspiration cytology and oncogene mutation screening: are we ready?. <i>F1000 Medicine Reports</i> , 2010, 2, 62.	2.9	7
34	Interaction between HMGA1 and Retinoblastoma Protein Is Required for Adipocyte Differentiation. <i>Journal of Biological Chemistry</i> , 2009, 284, 25993-26004.	3.4	16
35	<i>Helicobacter pylori</i> Hp(2â€“20) Promotes Migration and Proliferation of Gastric Epithelial Cells by Interacting with Formyl Peptide Receptors In Vitro and Accelerates Gastric Mucosal Healing In Vivo. <i>Journal of Immunology</i> , 2009, 183, 3761-3769.	0.8	60
36	XB130, a tissue-specific adaptor protein that couples the RET/PTC oncogenic kinase to PI 3-kinase pathway. <i>Oncogene</i> , 2009, 28, 937-949.	5.9	59

#	ARTICLE	IF	CITATIONS
37	Genetica molecolare del carcinoma tiroideo differenziato: implicazioni diagnostiche e terapeutiche. L Endocrinologo, 2009, 10, 114-118.	0.0	0
38	A New Germline Point Mutation in Ret Exon 8 (Cys ⁵¹⁵ Ser) in a Family with Medullary Thyroid Carcinoma. Thyroid, 2008, 18, 775-782.	4.5	27
39	A Cell Proliferation and Chromosomal Instability Signature in Anaplastic Thyroid Carcinoma. Cancer Research, 2007, 67, 10148-10158.	0.9	167
40	RET/Papillary Thyroid Carcinoma Oncogenic Signaling through the Rap1 Small GTPase. Cancer Research, 2007, 67, 381-390.	0.9	50
41	Receptor- and Non-Receptor Tyrosine Kinases Induce Processing of the Amyloid Precursor Protein: Role of the Low-Density Lipoprotein Receptor-Related Protein. Neurodegenerative Diseases, 2007, 4, 94-100.	1.4	7
42	Biological Role and Potential Therapeutic Targeting of the Chemokine Receptor CXCR4 in Undifferentiated Thyroid Cancer. Cancer Research, 2007, 67, 11821-11829.	0.9	100
43	OPN/CD44v6 overexpression in laryngeal dysplasia and correlation with clinical outcome. British Journal of Cancer, 2007, 97, 1545-1551.	6.4	32
44	RET/PTC activation in papillary thyroid carcinoma: European Journal of Endocrinology Prize Lecture. European Journal of Endocrinology, 2006, 155, 645-653.	3.7	176
45	Biochemical and molecular characterization of the novel BRAFV599Ins mutation detected in a classic papillary thyroid carcinoma. Oncogene, 2006, 25, 4235-4240.	5.9	56
46	Thyroid targeting of the N-ras(Gln61Lys) oncogene in transgenic mice results in follicular tumors that progress to poorly differentiated carcinomas. Oncogene, 2006, 25, 5467-5474.	5.9	66
47	HMGA2 induces pituitary tumorigenesis by enhancing E2F1 activity. Cancer Cell, 2006, 9, 459-471.	16.8	226
48	The Receptor-Type Protein Tyrosine Phosphatase J Antagonizes the Biochemical and Biological Effects of RET-Derived Oncoproteins. Cancer Research, 2006, 66, 6280-6287.	0.9	44
49	Activation of the Erk8 Mitogen-activated Protein (MAP) Kinase by RET/PTC3, a Constitutively Active Form of the RET Proto-oncogene. Journal of Biological Chemistry, 2006, 281, 10567-10576.	3.4	42
50	BRAF Is a Therapeutic Target in Aggressive Thyroid Carcinoma. Clinical Cancer Research, 2006, 12, 1623-1629.	7.0	160
51	The RET/PTC-RAS-BRAF linear signaling cascade mediates the motile and mitogenic phenotype of thyroid cancer cells. Journal of Clinical Investigation, 2005, 115, 1068-1081.	8.2	231
52	RAI(ShcC/N-Shc)-dependent recruitment of GAB1 to RET oncoproteins potentiates PI3-K signalling in thyroid tumors. Oncogene, 2005, 24, 6303-6313.	5.9	30
53	Overexpression of the Cytokine Osteopontin Identifies Aggressive Laryngeal Squamous Cell Carcinomas and Enhances Carcinoma Cell Proliferation and Invasiveness. Clinical Cancer Research, 2005, 11, 8019-8027.	7.0	53
54	Osteopontin Is Overexpressed in Human Papillary Thyroid Carcinomas and Enhances Thyroid Carcinoma Cell Invasiveness. Journal of Clinical Endocrinology and Metabolism, 2005, 90, 5270-5278.	3.6	71

#	ARTICLE	IF	CITATIONS
55	The RET/PTC-RAS-BRAF linear signaling cascade mediates the motile and mitogenic phenotype of thyroid cancer cells. <i>Journal of Clinical Investigation</i> , 2005, 115, 1068-1081.	8.2	126
56	Minireview: RET: Normal and Abnormal Functions. <i>Endocrinology</i> , 2004, 145, 5448-5451.	2.8	160
57	A New Germline RET Mutation Apparently Devoid of Transforming Activity Serendipitously Discovered in a Patient with Atrophic Autoimmune Thyroiditis and Primary Ovarian Failure. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004, 89, 4810-4816.	3.6	18
58	Regulation of p27Kip1 Protein Levels Contributes to Mitogenic Effects of the RET/PTC Kinase in Thyroid Carcinoma Cells. <i>Cancer Research</i> , 2004, 64, 3823-3829.	0.9	45
59	Autocrine stimulation by osteopontin plays a pivotal role in the expression of the mitogenic and invasive phenotype of RET/PTC-transformed thyroid cells. <i>Oncogene</i> , 2004, 23, 2188-2196.	5.9	43
60	Functional expression of the CXCR4 chemokine receptor is induced by RET/PTC oncogenes and is a common event in human papillary thyroid carcinomas. <i>Oncogene</i> , 2004, 23, 5958-5967.	5.9	119
61	The Oncogenic Activity of RET Point Mutants for Follicular Thyroid Cells May Account for the Occurrence of Papillary Thyroid Carcinoma in Patients Affected by Familial Medullary Thyroid Carcinoma. <i>American Journal of Pathology</i> , 2004, 165, 511-521.	3.8	35
62	Ras-mediated apoptosis of PC CL 3 rat thyroid cells induced by RET/PTC oncogenes. <i>Oncogene</i> , 2003, 22, 246-255.	5.9	46
63	Protein kinase C α activation by RET: evidence for a negative feedback mechanism controlling RET tyrosine kinase. <i>Oncogene</i> , 2003, 22, 2942-2949.	5.9	27
64	Tyrosine kinase oncoprotein, RET/PTC3, induces the secretion of myeloid growth and chemotactic factors. <i>Oncogene</i> , 2003, 22, 4569-4577.	5.9	67
65	Efficient Inhibition of RET/Papillary Thyroid Carcinoma Oncogenic Kinases by 4-Amino-5-(4-Chloro-Phenyl)-7-(<i>trans</i> -Butyl)Pyrazolo[3,4- <i>d</i>]Pyrimidine (PP2). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003, 88, 1897-1902.	3.6	115
66	The Neuron-Specific Rai (Shc) Adaptor Protein Inhibits Apoptosis by Coupling Ret to the Phosphatidylinositol 3-Kinase/Akt Signaling Pathway. <i>Molecular and Cellular Biology</i> , 2002, 22, 7351-7363.	2.3	84
67	Potent Mitogenicity of the RET/PTC3 Oncogene Correlates with Its Prevalence in Tall-Cell Variant of Papillary Thyroid Carcinoma. <i>American Journal of Pathology</i> , 2002, 160, 247-254.	3.8	103
68	Glial cell line-derived neurotrophic factor induces proliferative inhibition of NT2/D1 cells through RET-mediated up-regulation of the cyclin-dependent kinase inhibitor p27kip 1. <i>Oncogene</i> , 2002, 21, 1739-1749.	5.9	13
69	Cytoplasmic relocation and inhibition of the cyclin-dependent kinase inhibitor p27Kip1 by PKB/Akt-mediated phosphorylation in breast cancer. <i>Nature Medicine</i> , 2002, 8, 1136-1144.	30.7	644
70	Molecular Mechanisms of RET Activation in Human Cancer. <i>Annals of the New York Academy of Sciences</i> , 2002, 963, 116-121.	3.8	137
71	Molecular Mechanisms of RET Activation in Human Neoplasia. , 2002, , 176-183.		0
72	The insulin receptor substrate (IRS)-1 recruits phosphatidylinositol 3-kinase to Ret: evidence for a competition between Shc and IRS-1 for the binding to Ret. <i>Oncogene</i> , 2001, 20, 209-218.	5.9	57

#	ARTICLE	IF	CITATIONS
73	RET/PTC1 oncogene signaling in PC Cl 3 thyroid cells requires the small GTP-binding protein Rho. <i>Oncogene</i> , 2001, 20, 6973-6982.	5.9	45
74	Docking Protein FRS2 Links the Protein Tyrosine Kinase RET and Its Oncogenic Forms with the Mitogen-Activated Protein Kinase Signaling Cascade. <i>Molecular and Cellular Biology</i> , 2001, 21, 4177-4187.	2.3	123
75	Critical Role of the HMGI(Y) Proteins in Adipocytic Cell Growth and Differentiation. <i>Molecular and Cellular Biology</i> , 2001, 21, 2485-2495.	2.3	86
76	Tyrosines 1015 and 1062 Are <i>in Vivo</i> Autophosphorylation Sites in Ret and Ret-Derived Oncoproteins. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2000, 85, 3898-3907.	3.6	54
77	Pivotal Role of the RB Family Proteins in <i>in Vitro</i> Thyroid Cell Transformation. <i>Experimental Cell Research</i> , 2000, 260, 257-267.	2.6	10
78	Different mutations of the RET gene cause different human tumoral diseases. <i>Biochimie</i> , 1999, 81, 397-402.	2.6	12
79	Signalling of the Ret receptor tyrosine kinase through the c-Jun NH2-terminal protein kinases (JNKs): evidence for a divergence of the ERKs and JNKs pathways induced by Ret. <i>Oncogene</i> , 1998, 16, 2435-2445.	5.9	112
80	Molecular biology of the MEN2 gene. <i>Journal of Internal Medicine</i> , 1998, 243, 505-508.	6.0	42
81	Glial Cell Line-Derived Neurotrophic Factor Differentially Stimulates Ret Mutants Associated with the Multiple Endocrine Neoplasia Type 2 Syndromes and Hirschsprung's Disease. <i>Endocrinology</i> , 1998, 139, 3613-3619.	2.8	32
82	The ret/ptc1 Oncogene Is Activated in Familial Adenomatous Polyposis-Associated Thyroid Papillary Carcinomas. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1998, 83, 1003-1006.	3.6	82
83	The ret/ptc1 Oncogene Is Activated in Familial Adenomatous Polyposis-Associated Thyroid Papillary Carcinomas. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1998, 83, 1003-1006.	3.6	49
84	Glial cell line-derived neurotrophic factor (GDNF) stimulates ret activity. <i>Rendiconti Lincei</i> , 1997, 8, 139-149.	2.2	0
85	Only the Substitution of Methionine 918 with a Threonine and Not with Other Residues Activates RET Transforming Potential. <i>Endocrinology</i> , 1997, 138, 1450-1455.	2.8	7
86	Molecular defects in thyroid carcinomas: Role of the RET oncogene in thyroid neoplastic transformation. <i>European Journal of Endocrinology</i> , 1995, 133, 513-522.	3.7	56
87	PTC is a novel rearranged form of the ret proto-oncogene and is frequently detected <i>in vivo</i> in human thyroid papillary carcinomas. <i>Cell</i> , 1990, 60, 557-563.	28.9	905