Jordi Bort

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6828329/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Crop phenotyping in a context of global change: What to measure and how to do it. Journal of Integrative Plant Biology, 2022, 64, 592-618. | 4.1 | 29 |
| 2 | Effect of irrigation salinity and ecotype on the growth, physiological indicators and seed yield and quality of Salicornia europaea. Plant Science, 2021, 304, 110819. | 1.7 | 20 |
| 3 | Use of RGB Vegetation Indexes in Assessing Early Effects of Verticillium Wilt of Olive in Asymptomatic Plants in High and Low Fertility Scenarios. Remote Sensing, 2019, 11, 607. | 1.8 | 17 |
| 4 | The Hydrogen Isotope Composition δ ² H Reflects Plant Performance. Plant Physiology, 2019, 180, 793-812. | 2.3 | 41 |
| 5 | Identification of traits associated with barley yield performance using contrasting nitrogen fertilizations and genotypes. Plant Science, 2019, 282, 83-94. | 1.7 | 7 |
| 6 | Assessment of heavy metal tolerance in two plant species growing in experimental disturbed polluted urban soil. Journal of Soils and Sediments, 2018, 18, 2305-2317. | 1.5 | 31 |
| 7 | Challenges and Bottlenecks in VAV Phenotyping. , 2018, , . | | 1 |
| 8 | Post-green revolution genetic advance in durum wheat: The case of Spain. Field Crops Research, 2018, 228, 158-169. | 2.3 | 49 |
| 9 | Durum wheat ears perform better than the flag leaves under water stress: Gene expression and physiological evidence. Environmental and Experimental Botany, 2018, 153, 271-285. | 2.0 | 52 |
| 10 | Agronomic and physiological responses of Chinese facultative wheat genotypes to high-yielding Mediterranean conditions. Journal of Agricultural Science, 2016, 154, 870-889. | 0.6 | 10 |
| 11 | The combined use of vegetation indices and stable isotopes to predict durum wheat grain yield under contrasting water conditions. Agricultural Water Management, 2015, 158, 196-208. | 2.4 | 39 |
| 12 | Low-cost assessment of wheat resistance to yellow rust through conventional RGB images. Computers and Electronics in Agriculture, 2015, 116, 20-29. | 3.7 | 44 |
| 13 | Comparative performance of the stable isotope signatures of carbon, nitrogen and oxygen in assessing early vigour and grain yield in durum wheat. Journal of Agricultural Science, 2014, 152, 408-426. | 0.6 | 19 |
| 14 | Physiological traits contributed to the recent increase in yield potential of winter wheat from Henan Province, China. Journal of Integrative Plant Biology, 2014, 56, 492-504. | 4.1 | 46 |
| 15 | Contribution of the ear and the flag leaf to grain filling in durum wheat inferred from the carbon isotope signature: Genotypic and growing conditions effects. Journal of Integrative Plant Biology, 2014, 56, 444-454. | 4.1 | 90 |
| 16 | Molecular and physiological mechanisms associated with root exposure to mercury in barley. Metallomics, 2013, 5, 1305. | 1.0 | 22 |
| 17 | Comparative performance of δ13C, δ18O and δ15N for phenotyping durum wheat adaptation to a dryland environment. Functional Plant Biology, 2013, 40, 595. | 1.1 | 88 |
| 18 | Accumulation and toxic effects of chromium and zinc in Iris pseudacorus L. Acta Physiologiae Plantarum, 2012, 34, 1217-1228. | 1.0 | 42 |

Jordi Bort

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Ultrastructure and subcellular distribution of Cr in Iris pseudacorus L. using TEM and X-ray microanalysis. Cell Biology and Toxicology, 2012, 28, 57-68. | 2.4 | 58 |
| 20 | Mixed model association scans of multi-environmental trial data reveal major loci controlling yield and yield related traits in Hordeum vulgare in Mediterranean environments. Theoretical and Applied Genetics, 2011, 122, 1363-1373. | 1.8 | 75 |
| 21 | NDVI as a potential tool for predicting biomass, plant nitrogen content and growth in wheat genotypes subjected to different water and nitrogen conditions. Cereal Research Communications, 2011, 39, 147-159. | 0.8 | 147 |
| 22 | Physiological responses of Eichhornia crassipes [Mart.] Solms to the combined exposure to excess nutrients and Hg. Brazilian Journal of Plant Physiology, 2009, 21, 1-12. | 0.5 | 12 |
| 23 | Patterns of genetic diversity and linkage disequilibrium in a highly structured Hordeum vulgare association-mapping population for the Mediterranean basin. Theoretical and Applied Genetics, 2009, 119, 175-187. | 1.8 | 99 |
| 24 | Mapping adaptation of barley to droughted environments. Euphytica, 2008, 161, 35-45. | 0.6 | 44 |
| 25 | Barley adaptation and improvement in the Mediterranean basin. Plant Breeding, 2008, 127, 554-560. | 1.0 | 40 |
| 26 | Quantitative Trait Loci for Grain Yield and Adaptation of Durum Wheat (<i>Triticum durum</i> Desf.) Across a Wide Range of Water Availability. Genetics, 2008, 178, 489-511. | 1.2 | 397 |
| 27 | The Photosynthetic Role of Ears in C3 Cereals: Metabolism, Water Use Efficiency and Contribution to Grain Yield. Critical Reviews in Plant Sciences, 2007, 26, 1-16. | 2.7 | 196 |
| 28 | Relationships of grain ?13C and ?18O with wheat phenology and yield under water-limited conditions. Annals of Applied Biology, 2007, 150, 207-215. | 1.3 | 61 |
| 29 | Using vegetation indices derived from conventional digital cameras as selection criteria for wheat breeding in water-limited environments. Annals of Applied Biology, 2007, 150, 227-236. | 1.3 | 150 |
| 30 | Can wheat yield be assessed by early measurements of Normalized Difference Vegetation Index?. Annals of Applied Biology, 2007, 150, 253-257. | 1.3 | 164 |
| 31 | Water use efficiency in C3cereals under Mediterranean conditions: a review of physiological aspects. Annals of Applied Biology, 2007, 150, 307-321. | 1.3 | 192 |
| 32 | The combined effect of constant water deficit and nitrogen supply on WUE, NUE and Δ ¹³ C in durum wheat potted plants. Annals of Applied Biology, 2007, 151, 277-289. | 1.3 | 116 |
| 33 | A panel of elite accessions of durum wheat (Triticum durum Desf.) suitable for association mapping studies. Plant Genetic Resources: Characterisation and Utilisation, 2006, 4, 79-85. | 0.4 | 54 |
| 34 | Detection and Quantification of Unbound Phytochelatin 2 in Plant Extracts of Brassica napus Grown with Different Levels of Mercury. Plant Physiology, 2006, 142, 742-749. | 2.3 | 59 |
| 35 | Water management practices and climate in ancient agriculture: inferences from the stable isotope composition of archaeobotanical remains. Vegetation History and Archaeobotany, 2005, 14, 510-517. | 1.0 | 185 |
| 36 | Factors affecting the grain yield predicting attributes of spectral reflectance indices in durum wheat: growing conditions, genotype variability and date of measurement. International Journal of Remote Sensing, 2005, 26, 2337-2358. | 1.3 | 39 |

Jordi Bort

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Comparison of flag leaf and ear photosynthesis with biomass and grain yield of durum wheat under various water conditions and genotypes. Agronomy for Sustainable Development, 2004, 24, 19-28. | 0.8 | 87 |
| 38 | Relationships between early vigour, grain yield, leaf structure and stable isotope composition in field grown barley. Plant Physiology and Biochemistry, 1998, 36, 889-897. | 2.8 | 36 |
| 39 | Refixation of respiratory CO2in the ears of C3cereals. Journal of Experimental Botany, 1996, 47, 1567-1575. | 2.4 | 73 |
| 40 | Lack of C4 photosynthetic metabolism in ears of C3 cereals. Plant, Cell and Environment, 1995, 18, 697-702. | 2.8 | 24 |
| 41 | Role of awns in ear water-use efficiency and grain weight in barley. Agronomy for Sustainable Development, 1994, 14, 133-139. | 0.8 | 59 |
| 42 | Immunocytochemical localization of phosphoenolpyruvate carboxylase and photosynthetic gas-exchange characteristics in ears of Triticum durum Desf Planta, 1993, 191, 507. | 1.6 | 36 |