

# Pascal DollÃ©

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6842980/publications.pdf>

Version: 2024-02-01

138  
papers

18,980  
citations

17405

63  
h-index

15218

126  
g-index

150  
all docs

150  
docs citations

150  
times ranked

16280  
citing authors

#	ARTICLE	IF	CITATIONS
1	Loss of morphine-induced analgesia, reward effect and withdrawal symptoms in mice lacking the $\mu$ -opioid-receptor gene. <i>Nature</i> , 1996, 383, 819-823.	13.7	1,652
2	Embryonic retinoic acid synthesis is essential for early mouse post-implantation development. <i>Nature Genetics</i> , 1999, 21, 444-448.	9.4	963
3	Retinoic acid signalling during development. <i>Development (Cambridge)</i> , 2012, 139, 843-858.	1.2	693
4	Genetic analysis of RXR $\beta$ developmental function: Convergence of RXR and RAR signaling pathways in heart and eye morphogenesis. <i>Cell</i> , 1994, 78, 987-1003.	13.5	671
5	A homeotic transformation is generated in the rostral branchial region of the head by disruption of <i>Hoxa-2</i> , which acts as a selector gene. <i>Cell</i> , 1993, 75, 1333-1349.	13.5	612
6	Retinoic acid in development: towards an integrated view. <i>Nature Reviews Genetics</i> , 2008, 9, 541-553.	7.7	603
7	Poly(ADP-ribose) Polymerase-2 (PARP-2) Is Required for Efficient Base Excision DNA Repair in Association with PARP-1 and XRCC1. <i>Journal of Biological Chemistry</i> , 2002, 277, 23028-23036.	1.6	602
8	Coordinate expression of the murine <i>Hox-5</i> complex homeobox-containing genes during limb pattern formation. <i>Nature</i> , 1989, 342, 767-772.	13.7	593
9	A High-Resolution Anatomical Atlas of the Transcriptome in the Mouse Embryo. <i>PLoS Biology</i> , 2011, 9, e1000582.	2.6	552
10	Differential expression of genes encoding $\hat{1}$ , $\hat{2}$ and $\hat{3}$ retinoic acid receptors and CRABP in the developing limbs of the mouse. <i>Nature</i> , 1989, 342, 702-705.	13.7	496
11	The retinoic acid-metabolizing enzyme, CYP26A1, is essential for normal hindbrain patterning, vertebral identity, and development of posterior structures. <i>Genes and Development</i> , 2001, 15, 226-240.	2.7	492
12	Studies of human, mouse and yeast homologues indicate a mitochondrial function for frataxin. <i>Nature Genetics</i> , 1997, 16, 345-351.	9.4	489
13	Restricted expression and retinoic acid-induced downregulation of the retinaldehyde dehydrogenase type 2 (RALDH-2) gene during mouse development. <i>Mechanisms of Development</i> , 1997, 62, 67-78.	1.7	486
14	Identification and characterization of rod-derived cone viability factor. <i>Nature Genetics</i> , 2004, 36, 755-759.	9.4	463
15	Disruption of the <i>Hoxd-13</i> gene induces localized heterochrony leading to mice with neotenic limbs. <i>Cell</i> , 1993, 75, 431-441.	13.5	443
16	Oral-facial-digital type I protein is required for primary cilia formation and left-right axis specification. <i>Nature Genetics</i> , 2006, 38, 112-117.	9.4	299
17	Homeotic transformation of the occipital bones of the skull by ectopic expression of a homeobox gene. <i>Nature</i> , 1992, 359, 835-841.	13.7	285
18	Developmental expression of murine retinoid X receptor (RXR) genes. <i>Mechanisms of Development</i> , 1994, 45, 91-104.	1.7	285

#	ARTICLE	IF	CITATIONS
19	Cardiac T-box factor Tbx20 directly interacts with Nkx2-5, GATA4, and GATA5 in regulation of gene expression in the developing heart. <i>Developmental Biology</i> , 2003, 262, 206-224.	0.9	260
20	Dorsal pancreas agenesis in retinoic acid-deficient Raldh2 mutant mice. <i>Developmental Biology</i> , 2005, 284, 399-411.	0.9	226
21	Differential expression of retinoic acid-synthesizing (RALDH) enzymes during fetal development and organ differentiation in the mouse. <i>Mechanisms of Development</i> , 2002, 110, 165-171.	1.7	220
22	Expression of GHF-1 protein in mouse pituitaries correlates both temporally and spatially with the onset of growth hormone gene activity. <i>Cell</i> , 1990, 60, 809-820.	13.5	216
23	Retinoic Acid Controls the Bilateral Symmetry of Somite Formation in the Mouse Embryo. <i>Science</i> , 2005, 308, 563-566.	6.0	214
24	Genetic evidence that oxidative derivatives of retinoic acid are not involved in retinoid signaling during mouse development. <i>Nature Genetics</i> , 2002, 31, 84-88.	9.4	213
25	Cloning of a novel retinoic-acid metabolizing cytochrome P450, Cyp26B1, and comparative expression analysis with Cyp26A1 during early murine development. <i>Mechanisms of Development</i> , 2001, 107, 195-201.	1.7	208
26	The regional pattern of retinoic acid synthesis by RALDH2 is essential for the development of posterior pharyngeal arches and the enteric nervous system. <i>Development (Cambridge)</i> , 2003, 130, 2525-2534.	1.2	200
27	Developmental expression pattern of Stra6, a retinoic acid-responsive gene encoding a new type of membrane protein. <i>Mechanisms of Development</i> , 1997, 63, 173-186.	1.7	184
28	AP-2.2, a novel gene related to AP-2, is expressed in the forebrain, limbs and face during mouse embryogenesis. <i>Mechanisms of Development</i> , 1996, 54, 83-94.	1.7	175
29	Differential expression of the retinoic acid-metabolizing enzymes CYP26A1 and CYP26B1 during murine organogenesis. <i>Mechanisms of Development</i> , 2002, 110, 173-177.	1.7	172
30	Efficient Cloning of cDNAs of Retinoic Acid-Responsive Genes in P19 Embryonal Carcinoma Cells and Characterization of a Novel Mouse Gene, Stra1 (Mouse LERK-2/Eplg2). <i>Developmental Biology</i> , 1995, 170, 420-433.	0.9	168
31	Expression of the murine Dlx-1 homeobox gene during facial, ocular and limb development. <i>Differentiation</i> , 1992, 49, 93-99.	1.0	159
32	Non-cell-autonomous retinoid signaling is crucial for renal development. <i>Development (Cambridge)</i> , 2010, 137, 283-292.	1.2	149
33	Mouse Lbx1 and human LBX1 define a novel mammalian homeobox gene family related to the Drosophila ladybird genes. <i>Mechanisms of Development</i> , 1995, 53, 345-356.	1.7	147
34	Hox genes define distinct progenitor sub-domains within the second heart field. <i>Developmental Biology</i> , 2011, 353, 266-274.	0.9	144
35	Decreased embryonic retinoic acid synthesis results in a DiGeorge syndrome phenotype in newborn mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 1763-1768.	3.3	143
36	miRNeasy: a microRNA expression atlas of the mouse eye. <i>BMC Genomics</i> , 2010, 11, 715.	1.2	140

#	ARTICLE	IF	CITATIONS
37	Developmental roles of the retinoic acid receptors. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1995, 53, 475-486.	1.2	137
38	Retinoic acid regulates morphogenesis and patterning of posterior foregut derivatives. <i>Developmental Biology</i> , 2006, 297, 433-445.	0.9	136
39	The Hox-4.8 gene is localized at the 5' extremity of the Hox-4 complex and is expressed in the most posterior parts of the body during development. <i>Mechanisms of Development</i> , 1991, 36, 3-13.	1.7	134
40	Cyp26C1 encodes a novel retinoic acid-metabolizing enzyme expressed in the hindbrain, inner ear, first branchial arch and tooth buds during murine development. <i>Gene Expression Patterns</i> , 2003, 3, 449-454.	0.3	133
41	CTIP1 and CTIP2 are differentially expressed during mouse embryogenesis. <i>Gene Expression Patterns</i> , 2004, 4, 733-739.	0.3	133
42	CYP26A1 and CYP26C1 cooperatively regulate anterior-posterior patterning of the developing brain and the production of migratory cranial neural crest cells in the mouse. <i>Developmental Biology</i> , 2007, 302, 399-411.	0.9	128
43	Tissue-specific expression of retinoic acid receptor isoform transcripts in the mouse embryo. <i>Mechanisms of Development</i> , 2000, 94, 223-232.	1.7	117
44	Developmental expression of retinoic acid receptors (RARs). <i>Nuclear Receptor Signaling</i> , 2009, 7, nrs.07006.	1.0	116
45	Defects of the Chorioallantoic Placenta in Mouse RXR $\alpha$ Null Fetuses. <i>Developmental Biology</i> , 1997, 191, 29-41.	0.9	115
46	Retinaldehyde dehydrogenase 2 (RALDH2)-mediated retinoic acid synthesis regulates early mouse embryonic forebrain development by controlling FGF and sonic hedgehog signaling. <i>Development (Cambridge)</i> , 2006, 133, 351-361.	1.2	114
47	Retinaldehyde dehydrogenase 2 (RALDH2)- independent patterns of retinoic acid synthesis in the mouse embryo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 16111-16116.	3.3	109
48	Signaling hierarchy downstream of retinoic acid that independently regulates vascular remodeling and endothelial cell proliferation. <i>Genes and Development</i> , 2004, 18, 1345-1358.	2.7	108
49	AP-2.2: A Novel AP-2-Related Transcription Factor Induced by Retinoic Acid during Differentiation of P19 Embryonal Carcinoma Cells. <i>Experimental Cell Research</i> , 1996, 225, 338-347.	1.2	106
50	Retinoids regulate the anterior expression boundaries of 5' Hoxb genes in posterior hindbrain. <i>EMBO Journal</i> , 2003, 22, 262-269.	3.5	103
51	Sequence and expression pattern of the Stra7 (Gbx-2) homeobox-containing gene induced by retinoic acid in P19 embryonal carcinoma cells. <i>Developmental Dynamics</i> , 1995, 204, 372-382.	0.8	100
52	Early mouse caudal development relies on crosstalk between retinoic acid, Shh and Fgf signalling pathways. <i>Development (Cambridge)</i> , 2009, 136, 665-676.	1.2	98
53	Retinoids control anterior and dorsal properties in the developing forebrain. <i>Developmental Biology</i> , 2007, 303, 362-375.	0.9	97
54	Involvement of retinol dehydrogenase 10 in embryonic patterning and rescue of its loss of function by maternal retinaldehyde treatment. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 16687-16692.	3.3	97

#	ARTICLE	IF	CITATIONS
55	Endogenous retinoic acid regulates cardiac progenitor differentiation. Proceedings of the National Academy of Sciences of the United States of America, 2010, 107, 9234-9239.	3.3	96
56	Distinct roles for retinoic acid receptors alpha and beta in early lung morphogenesis. Developmental Biology, 2006, 291, 12-24.	0.9	93
57	Meis2, a novel mousePbx-related homeobox gene induced by retinoic acid during differentiation of P19 embryonal carcinoma cells. , 1997, 210, 173-183.		88
58	Retinoic acid signalling is required for specification of pronephric cell fate. Developmental Biology, 2006, 299, 35-51.	0.9	80
59	The Expression Pattern of the Mouse Receptor Tyrosine Kinase Gene MDK1 Is Conserved through Evolution and Requires Hoxa-2 for Rhombomere-Specific Expression in Mouse Embryos. Developmental Biology, 1996, 177, 397-412.	0.9	79
60	Developing with lethal RA levels: genetic ablation of Rarg can restore the viability of mice lacking Cyp26a1. Development (Cambridge), 2003, 130, 1449-1459.	1.2	74
61	Genetic disruption of CYP26B1 severely affects development of neural crest derived head structures, but does not compromise hindbrain patterning. Developmental Dynamics, 2009, 238, 732-745.	0.8	73
62	Differential expression of transcripts encoding retinoid binding proteins and retinoic acid receptors during placentation of the mouse. , 1997, 208, 199-210.		72
63	Retinaldehyde dehydrogenase 2 and Hoxc8 are required in the murine brachial spinal cord for the specification of Lim1+ motoneurons and the correct distribution of Islet1+ motoneurons. Development (Cambridge), 2005, 132, 1611-1621.	1.2	70
64	Insertion of a targeting construct in a Hoxd-10 allele can influence the control of Hoxd-9 expression. Developmental Dynamics, 1994, 201, 366-377.	0.8	66
65	Differential expression of the TEF family of transcription factors in the murine placenta and during differentiation of primary human trophoblasts in vitro. , 1998, 212, 423-436.		66
66	Direct crossregulation between retinoic acid receptor $\hat{r}^2$ and Hox genes during hindbrain segmentation. Development (Cambridge), 2005, 132, 503-513.	1.2	65
67	Retinoid signaling in inner ear development. Journal of Neurobiology, 2006, 66, 687-704.	3.7	63
68	Embryonic retinoic acid synthesis is required for forelimb growth and anteroposterior patterning in the mouse. Development (Cambridge), 2002, 129, 3563-74.	1.2	62
69	Rescue of cytochrome P450 oxidoreductase (Por) mouse mutants reveals functions in vasculogenesis, brain and limb patterning linked to retinoic acid homeostasis. Developmental Biology, 2007, 303, 66-81.	0.9	61
70	Expression of the murine retinol dehydrogenase 10 ( <i>Rdh10</i> ) gene correlates with many sites of retinoid signalling during embryogenesis and organ differentiation. Developmental Dynamics, 2007, 236, 2899-2908.	0.8	60
71	Retinoic acid signaling regulates murine bronchial tubule formation. Mechanisms of Development, 2003, 120, 691-700.	1.7	50
72	FGF Signalling Regulates Chromatin Organisation during Neural Differentiation via Mechanisms that Can Be Uncoupled from Transcription. PLoS Genetics, 2013, 9, e1003614.	1.5	50

#	ARTICLE	IF	CITATIONS
73	Expression patterns of the Ets transcription factors from the PEA3 group during early stages of mouse development. <i>Mechanisms of Development</i> , 2001, 108, 191-195.	1.7	49
74	The oxidizing enzyme CYP26a1 tightly regulates the availability of retinoic acid in the gastrulating mouse embryo to ensure proper head development and vasculogenesis. <i>Developmental Dynamics</i> , 2007, 236, 644-653.	0.8	48
75	Truncation of the Catalytic Domain of the Cylindromatosis Tumor Suppressor Impairs Lung Maturation. <i>Neoplasia</i> , 2009, 11, 469-476.	2.3	47
76	Retinoic Acid Receptor $\hat{1}^2$ Controls Development of Striatonigral Projection Neurons through FGF-Dependent and Meis1-Dependent Mechanisms. <i>Journal of Neuroscience</i> , 2015, 35, 14467-14475.	1.7	47
77	Morphological and Molecular Characterization of Retinoic Acid-Induced Limb Duplications in Mice. <i>Developmental Biology</i> , 1996, 176, 185-198.	0.9	46
78	Kidney-specific inactivation of <i>Ofd1</i> leads to renal cystic disease associated with upregulation of the mTOR pathway. <i>Human Molecular Genetics</i> , 2010, 19, 2792-2803.	1.4	46
79	Stage and tissue-specific expression of the alcohol dehydrogenase 1 ( <i>Adh-1</i> ) gene during mouse development. <i>Developmental Dynamics</i> , 1994, 199, 199-213.	0.8	45
80	A Bidirectional Promoter Connects the Poly(ADP-ribose) Polymerase 2 (PARP-2) Gene to the Gene for RNase P RNA. <i>Journal of Biological Chemistry</i> , 2001, 276, 11092-11099.	1.6	43
81	Combinatorial signalling controls <i>Neurogenin2</i> expression at the onset of spinal neurogenesis. <i>Developmental Biology</i> , 2008, 321, 470-481.	0.9	43
82	Molars and incisors: show your microarray IDs. <i>BMC Research Notes</i> , 2013, 6, 113.	0.6	43
83	Differential expression of retinoic acid-inducible ( <i>Stra</i> ) genes during mouse placentation. <i>Mechanisms of Development</i> , 2000, 92, 295-299.	1.7	42
84	External Genitalia Formation. <i>Annals of the New York Academy of Sciences</i> , 2001, 948, 13-31.	1.8	42
85	The retinoic acid receptors $RAR^{\hat{1}\pm}$ and $RAR^{\hat{1}3}$ are required for inner ear development. <i>Mechanisms of Development</i> , 2002, 119, 213-223.	1.7	40
86	Expression Analysis of Murine Genes Using <i>In Situ</i> Hybridization With Radioactive and Nonradioactively Labeled RNA Probes. , 2006, 326, 61-88.		40
87	Mutations in the latent TGF-beta binding protein 3 ( <i>LTBP3</i> ) gene cause brachyolmia with amelogenesis imperfecta. <i>Human Molecular Genetics</i> , 2015, 24, 3038-3049.	1.4	40
88	<i>Vax2</i> regulates retinoic acid distribution and cone opsin expression in the vertebrate eye. <i>Development (Cambridge)</i> , 2011, 138, 261-271.	1.2	39
89	Developmental expression of the mouse <i>Evx-2</i> gene: relationship with the evolution of the HOM/Hox complex. <i>Development (Cambridge)</i> , 1994, 1994, 143-153.	1.2	38
90	Restricted expression of the <i>ron</i> gene encoding the macrophage stimulating protein receptor during mouse development. <i>Developmental Dynamics</i> , 1995, 204, 383-390.	0.8	37

#	ARTICLE	IF	CITATIONS
91	Dynamic expression of the retinoic acid-synthesizing enzyme retinol dehydrogenase 10 (rdh10) in the developing mouse brain and sensory organs. <i>Journal of Comparative Neurology</i> , 2008, 508, 879-892.	0.9	37
92	Retinoic acid regulates olfactory progenitor cell fate and differentiation. <i>Neural Development</i> , 2013, 8, 13.	1.1	35
93	Retinoic acid controls early neurogenesis in the developing mouse cerebral cortex. <i>Developmental Biology</i> , 2017, 430, 129-141.	0.9	35
94	Genome-wide Analysis of RAR $\beta$ Transcriptional Targets in Mouse Striatum Links Retinoic Acid Signaling with Huntington's Disease and Other Neurodegenerative Disorders. <i>Molecular Neurobiology</i> , 2017, 54, 3859-3878.	1.9	34
95	Molecular cloning, genomic structure, and expression analysis of the mouse transcriptional intermediary factor 1 gamma gene. <i>Gene</i> , 2004, 334, 3-13.	1.0	33
96	Dynamic expression of retinoic acid-synthesizing and -metabolizing enzymes in the developing mouse inner ear. <i>Journal of Comparative Neurology</i> , 2006, 496, 643-654.	0.9	33
97	Sox2 acts as a rheostat of epithelial to mesenchymal transition during neural crest development. <i>Frontiers in Physiology</i> , 2014, 5, 345.	1.3	33
98	Expression of T $\alpha$ : $\beta$ mismatch-specific thymidine-DNA glycosylase and DNA methyl transferase genes during development and tumorigenesis. <i>Oncogene</i> , 1998, 17, 1577-1585.	2.6	28
99	Transcriptomic Analysis of Murine Embryos Lacking Endogenous Retinoic Acid Signaling. <i>PLoS ONE</i> , 2013, 8, e62274.	1.1	27
100	Endogenous retinoic acid signaling is required for maintenance and regeneration of cornea. <i>Experimental Eye Research</i> , 2017, 154, 190-195.	1.2	27
101	Fate of retinoic acid-activated embryonic cell lineages. <i>Developmental Dynamics</i> , 2010, 239, 3260-3274.	0.8	26
102	The macroPARP genes <i>parp9</i> and <i>parp14</i> are developmentally and differentially regulated in mouse tissues. <i>Developmental Dynamics</i> , 2008, 237, 209-215.	0.8	25
103	Retinoic Acid Deficiency Impairs the Vestibular Function. <i>Journal of Neuroscience</i> , 2013, 33, 5856-5866.	1.7	25
104	Genetic Inactivation of Prokineticin Receptor-1 Leads to Heart and Kidney Disorders. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 842-850.	1.1	24
105	Retinoic Acid-Dependent Signaling Pathways and Lineage Events in the Developing Mouse Spinal Cord. <i>PLoS ONE</i> , 2012, 7, e32447.	1.1	24
106	Prokineticin receptor-1 signaling promotes Epicardial to Mesenchymal Transition during heart development. <i>Scientific Reports</i> , 2016, 6, 25541.	1.6	24
107	Complementary expression patterns of retinoid acid-synthesizing and -metabolizing enzymes in pre-natal mouse inner ear structures. <i>Gene Expression Patterns</i> , 2004, 4, 123-133.	0.3	23
108	RSK2 Is a Modulator of Craniofacial Development. <i>PLoS ONE</i> , 2014, 9, e84343.	1.1	23

#	ARTICLE	IF	CITATIONS
109	Specific expression of the retinoic acid-synthesizing enzyme RALDH2 during mouse inner ear development. <i>Mechanisms of Development</i> , 2001, 106, 185-189.	1.7	22
110	In Situ Hybridization with 35S-Labeled Probes for Retinoid Receptors. , 1998, 89, 247-267.		21
111	Essential role of the TFIID subunit TAF4 in murine embryogenesis and embryonic stem cell differentiation. <i>Nature Communications</i> , 2016, 7, 11063.	5.8	21
112	Local retinoic acid signaling directs emergence of the extraocular muscle functional unit. <i>PLoS Biology</i> , 2020, 18, e3000902.	2.6	21
113	Meningeal retinoic acid contributes to neocortical lamination and radial migration during mouse brain development. <i>Biology Open</i> , 2017, 6, 148-160.	0.6	20
114	Distinct retinoic acid receptor (RAR) isoforms control differentiation of embryonal carcinoma cells to dopaminergic or striatopallidal medium spiny neurons. <i>Scientific Reports</i> , 2017, 7, 13671.	1.6	19
115	Restricted expression of a novel retinoic acid responsive gene during limb bud dorsoventral patterning and endochondral ossification. , 1996, 19, 66-73.		18
116	Expression of the transcriptional intermediary factor TIF1± during mouse development and in the reproductive organs. <i>Mechanisms of Development</i> , 1999, 88, 111-117.	1.7	16
117	Deficiency of the SMOC2 matricellular protein impairs bone healing and produces age-dependent bone loss. <i>Scientific Reports</i> , 2020, 10, 14817.	1.6	16
118	Regulation of expression of the retinoic acid metabolizing enzyme CYP26A1 in uteri of ovariectomized mice after treatment with ovarian steroid hormones. <i>Molecular Reproduction and Development</i> , 2007, 74, 258-264.	1.0	15
119	Conditional (loxP-flanked) allele for the gene encoding the retinoic acid-synthesizing enzyme retinaldehyde dehydrogenase 2 (RALDH2). <i>Genesis</i> , 2006, 44, 155-158.	0.8	14
120	Retinoic Acid Excess Impairs Amelogenesis Inducing Enamel Defects. <i>Frontiers in Physiology</i> , 2016, 7, 673.	1.3	14
121	Retinoic acid signaling is directly activated in cardiomyocytes and protects mouse hearts from apoptosis after myocardial infarction. <i>ELife</i> , 2021, 10, .	2.8	14
122	Regulation of expression of the retinoic acid-synthesising enzymes retinaldehyde dehydrogenases in the uteri of ovariectomised mice after treatment with oestrogen, gestagen and their combination. <i>Reproduction, Fertility and Development</i> , 2006, 18, 339.	0.1	13
123	Enamel and dental anomalies in latent transforming growth factor beta binding protein 3 mutant mice. <i>European Journal of Oral Sciences</i> , 2017, 125, 8-17.	0.7	13
124	Retinoic Acid Receptor (RAR)-± Is Not Critically Required for Mediating Retinoic Acid Effects in the Developing Mouse Retina. , 2010, 51, 3281.		11
125	Rescue of morphogenetic defects and of retinoic acid signaling in retinaldehyde dehydrogenase 2 (Raldh2) mouse mutants by chimerism with wild-type cells. <i>Differentiation</i> , 2006, 74, 661-668.	1.0	10
126	Prokineticin receptor 1 is required for mesenchymal epithelial transition in kidney development. <i>FASEB Journal</i> , 2016, 30, 2733-2740.	0.2	7



#	ARTICLE	IF	CITATIONS
127	The homeodomain factor <i>Gbx1</i> is required for locomotion and cell specification in the dorsal spinal cord. <i>PeerJ</i> , 2013, 1, e142.	0.9	7
128	Molecular mediators of retinoic acid signaling during development. <i>Advances in Developmental Biology</i> (Amsterdam, Netherlands), 2006, , 105-143.	0.4	6
129	Structural and Functional Aspects of Mammalian Hox Genes. <i>Advances in Developmental Biochemistry</i> , 1993, , 57-109.	0.9	3
130	Retinoids and mouse placentation. <i>Placenta</i> , 1998, 19, 57-76.	0.7	3
131	Retinoic acid receptor beta protects striatopallidal medium spiny neurons from mitochondrial dysfunction and neurodegeneration. <i>Progress in Neurobiology</i> , 2022, 212, 102246.	2.8	3
132	Teratogenic effects of ethanol: Interaction with retinoid metabolism. <i>Toxicology Letters</i> , 2006, 164, S49-S50.	0.4	1
133	Retinoids and Heart Development. , 2010, , 237-253.		1
134	A Comparison of the Expression Domains of the Murine Hox-4, RARs and CRABP Genes Suggests Possible Functional Relationships During Patterning of the Vertebrate Limb. , 1991, , 65-73.		1
135	Malformaciones congénitas de las extremidades: embriología, etiología. <i>EMC Pediatría</i> , 2003, 38, 1-7.	0.0	0
136	Malformations congénitales des membres: embryologie, étiologie. <i>EMC - Pédiatrie - Maladies Infectieuses</i> , 2006, 1, 1-8.	0.0	0
137	ISDN2014_0036: REMOVED: Craniofacial development is fine tuned by Sox2. <i>International Journal of Developmental Neuroscience</i> , 2015, 47, 7-7.	0.7	0
138	Integrated Annotation and Analysis of In Situ Hybridization Images Using the ImAnno System: Application to the Ear and Sensory Organs of the Fetal Mouse. <i>PLoS ONE</i> , 2015, 10, e0118024.	1.1	0