
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6875433/publications.pdf Version: 2024-02-01



HENK | SCHEDS

#	Article	IF	CITATIONS
1	Comprehensive Health Assessment 3 Months After Recovery From Acute Coronavirus Disease 2019 (COVID-19). Clinical Infectious Diseases, 2021, 73, e1089-e1098.	5.8	332
2	Implementation Barriers for General Practice Guidelines on Low Back Pain. Spine, 2001, 26, E348-E353.	2.0	142
3	How unique is continuity of care? A review of continuity and related concepts. Family Practice, 2012, 29, 264-271.	1.9	117
4	Prevalence of physical symptoms of itch, pain and fatigue in patients with skin diseases in general practice. British Journal of Dermatology, 2007, 156, 1346-1349.	1.5	106
5	"Beating osteoARThritis†Development of a stepped care strategy to optimize utilization and timing of non-surgical treatment modalities for patients with hip or knee osteoarthritis. Clinical Rheumatology, 2011, 30, 1623-1629.	2.2	102
6	Continuity of Care: Is the Personal Doctor Still Important? A Survey of General Practitioners and Family Physicians in England and Wales, the United States, and the Netherlands. Annals of Family Medicine, 2005, 3, 353-359.	1.9	101
7	The effect of weekly specialist palliative care teleconsultations in patients with advanced cancer –a randomized clinical trial. BMC Medicine, 2017, 15, 119.	5.5	80
8	Impact of Interprofessional Education on Collaboration Attitudes, Skills, and Behavior Among Primary Care Professionals. Journal of Continuing Education in the Health Professions, 2012, 32, 196-204.	1.3	79
9	Familiarity with a GP and patients' evaluations of care. A cross-sectional study. Family Practice, 2004, 22, 15-19.	1.9	71
10	Nijmegen Continuity Questionnaire: Development and testing of a questionnaire that measures continuity of care. Journal of Clinical Epidemiology, 2011, 64, 1391-1399.	5.0	71
11	Health Care in The Netherlands. Journal of the American Board of Family Medicine, 2012, 25, S12-S17.	1.5	71
12	Measurement Properties of Questionnaires Measuring Continuity of Care: A Systematic Review. PLoS ONE, 2012, 7, e42256.	2.5	70
13	Skin Diseases in Family Medicine: Prevalence and Health Care Use. Annals of Family Medicine, 2008, 6, 349-354.	1.9	67
14	Continuity of care in general practice: a survey of patients' views. British Journal of General Practice, 2002, 52, 459-62.	1.4	66
15	Development of an instrument for the identification of frail older people as a target population for integrated care. British Journal of General Practice, 2013, 63, e225-e231.	1.4	54
16	Multicomponent Program to Reduce Functional Decline in Frail Elderly People: A Cluster Controlled Trial. Journal of the American Board of Family Medicine, 2016, 29, 209-217.	1.5	52
17	Home visits for frail older people: a qualitative study on the needs and preferences of frail older people and their informal caregivers. British Journal of General Practice, 2012, 62, e554-e560.	1.4	46
18	Barriers Impeding the Use of Non-pharmacological, Non-surgical Care in Hip and Knee Osteoarthritis. Journal of Clinical Rheumatology, 2017, 23, 405-410.	0.9	46

#	Article	IF	CITATIONS
19	Evaluation of an eHealth Intervention in Chronic Care for Frail Older People: Why Adherence is the First Target. Journal of Medical Internet Research, 2014, 16, e156.	4.3	46
20	Reasons for Treatment Choices in Knee and Hip Osteoarthritis: A Qualitative Study. Arthritis Care and Research, 2016, 68, 1260-1267.	3.4	44
21	Implementation of an innovative web-based conference table for community-dwelling frail older people, their informal caregivers and professionals: a process evaluation. BMC Health Services Research, 2012, 12, 251.	2.2	43
22	Health Care Use of Patients With Osteoarthritis of the Hip or Knee After Implementation of a Steppedâ€Care Strategy: An Observational Study. Arthritis Care and Research, 2014, 66, 817-827.	3.4	42
23	Measuring continuity of care: psychometric properties of the Nijmegen Continuity Questionnaire. British Journal of General Practice, 2012, 62, e949-e957.	1.4	39
24	The role of the general practitioner in multidisciplinary teams: a qualitative study in elderly care. BMC Family Practice, 2018, 19, 40.	2.9	39
25	Construct validity and reliability of a two-step tool for the identification of frail older people in primary care. Journal of Clinical Epidemiology, 2014, 67, 176-183.	5.0	36
26	Educating Dutch General Practitioners in Dementia Advance Care Planning: A Cluster Randomized Controlled Trial. Journal of the American Medical Directors Association, 2020, 21, 837-842.e4.	2.5	36
27	What went and what came? Morbidity trends in general practice from the Netherlands. European Journal of General Practice, 2008, 14, 13-24.	2.0	35
28	Preferences for receiving information among frail older adults and their informal caregivers: a qualitative study. Family Practice, 2012, 29, 742-747.	1.9	35
29	Increased incidence of squamous cell carcinoma of the skin after longâ€ŧerm treatment with azathioprine in patients with autoâ€immune inflammatory rheumatic diseases. Journal of the European Academy of Dermatology and Venereology, 2014, 28, 27-33.	2.4	34
30	The COVID-19 Pandemic in Nijmegen, the Netherlands: Changes in Presented Health Problems and Demand for Primary Care. Annals of Family Medicine, 2021, 19, 44-47.	1.9	33
31	Incidence, Causative Mechanisms, and Anatomic Localization of Stroke in Pituitary Adenoma Patients Treated With Postoperative Radiation Therapy Versus Surgery Alone. International Journal of Radiation Oncology Biology Physics, 2013, 87, 53-59.	0.8	31
32	Is the quality of data in an electronic medical record sufficient for assessing the quality of primary care?. Journal of the American Medical Informatics Association: JAMIA, 2014, 21, 692-698.	4.4	30
33	The predictive value of gait speed and maximum step length for falling in community-dwelling older persons. Age and Ageing, 2015, 44, 294-299.	1.6	29
34	Psychosocial well-being of patients with skin diseases in general practice. Journal of the European Academy of Dermatology and Venereology, 2007, 21, 070206173308001-???.	2.4	27
35	Predictive validity of a two-step tool to map frailty in primary care. BMC Medicine, 2015, 13, 287.	5.5	27
36	Filling the Gaps in a Fragmented Health Care System: Development of the Health and Welfare Information Portal (ZWIP). JMIR Research Protocols, 2012, 1, e10.	1.0	25

#	Article	IF	CITATIONS
37	Patient reported barriers and facilitators to using a self-management booklet for hip and knee osteoarthritis in primary care: results of a qualitative interview study. BMC Family Practice, 2013, 14, 181.	2.9	24
38	Heart failure patients' experiences with continuity of care and its relation to medication adherence: a cross-sectional study. BMC Family Practice, 2012, 13, 86.	2.9	23
39	Continuity in different care modes and its relationship to quality of life: a randomised controlled trial in patients with COPD. British Journal of General Practice, 2012, 62, e422-e428.	1.4	22
40	The CareWell-primary care program: design of a cluster controlled trial and process evaluation of a complex intervention targeting community-dwelling frail elderly. BMC Family Practice, 2012, 13, 115.	2.9	22
41	Continuity of care preferably measured from the patients' perspective. Journal of Clinical Epidemiology, 2010, 63, 998-999.	5.0	21
42	Agreement of general practitioners with the guideline-based stepped-care strategy for patients with osteoarthritis of the hip or knee: a cross-sectional study. BMC Family Practice, 2013, 14, 33.	2.9	21
43	Non-surgical care in patients with hip or knee osteoarthritis is modestly consistent with a stepped care strategy after its implementation. International Journal for Quality in Health Care, 2014, 26, 490-498.	1.8	21
44	The development of an ICF-based questionnaire for patients with chronic conditions in primary care. Journal of Clinical Epidemiology, 2018, 103, 92-100.	5.0	19
45	Advance care planning for patients with cancer in the palliative phase in Dutch general practices. Family Practice, 2019, 36, 587-593.	1.9	19
46	Patients' Needs for Contact With Their GP at the Time of Hospital Admission and Other Life Events: A Quantitative and Qualitative Exploration. Annals of Family Medicine, 2004, 2, 462-468.	1.9	18
47	Diagnosis of Frailty after a Comprehensive Geriatric Assessment: Differences between Family Physicians and Geriatricians. Journal of the American Board of Family Medicine, 2015, 28, 240-248.	1.5	18
48	Psychosocial, musculoskeletal and somatoform comorbidity in patients with chronic low back pain: original results from the Dutch Transition Project. Family Practice, 2015, 32, 297-304.	1.9	18
49	Which Frail Older Patients Use Online Health Communities and Why? A Mixed Methods Process Evaluation of Use of the Health and Welfare Portal. Journal of Medical Internet Research, 2014, 16, e278.	4.3	18
50	International variation in GP treatment strategies for subclinical hypothyroidism in older adults: a case-based survey. British Journal of General Practice, 2015, 65, e121-e132.	1.4	17
51	Factors Associated With Referral to Secondary Care in Patients With Osteoarthritis of the Hip or Knee After Implementation of a Steppedâ€Care Strategy. Arthritis Care and Research, 2017, 69, 216-225.	3.4	17
52	Family medicine trainees still value continuity of care. Family Medicine, 2004, 36, 51-4.	0.5	17
53	Study protocol: optimization of complex palliative care at home via telemedicine. A cluster randomized controlled trial. BMC Palliative Care, 2011, 10, 13.	1.8	16
54	The Diagnostic Value of the Patient's Reason for Encounter for Diagnosing Cancer in Primary Care. Journal of the American Board of Family Medicine, 2017, 30, 806-812.	1.5	16

#	Article	IF	CITATIONS
55	Incidence, course and risk factors of head injury: a retrospective cohort study. BMJ Open, 2018, 8, e020364.	1.9	16
56	Structured chronic primary care and health-related quality of life in chronic heart failure. BMC Health Services Research, 2009, 9, 104.	2.2	15
57	Cost-effectiveness of a multicomponent primary care program targeting frail elderly people. BMC Family Practice, 2018, 19, 62.	2.9	15
58	Preliminary effects of a regional approached multidisciplinary educational program on healthcare utilization in patients with hip or knee osteoarthritis: an observational study. BMC Family Practice, 2018, 19, 82.	2.9	14
59	Feasibility evaluation of a stepped procedure to identify community-dwelling frail older people in general practice. A mixed methods study. European Journal of General Practice, 2014, 20, 107-113.	2.0	13
60	Purposeful Incorporation of Patient Narratives in the Medical Record in the Netherlands. Journal of the American Board of Family Medicine, 2021, 34, 709-723.	1.5	13
61	Hierarchical structure and importance of patients' reasons for treatment choices in knee and hip osteoarthritis: a concept mapping study. Rheumatology, 2017, 56, 271-278.	1.9	12
62	Treatment Beliefs Underlying Intended Treatment Choices in Knee and Hip Osteoarthritis. International Journal of Behavioral Medicine, 2018, 25, 198-206.	1.7	12
63	Professional Care Networks of Frail Older People: An Explorative Survey Study from the Patient Perspective. International Journal of Integrated Care, 2020, 20, 1.	0.2	12
64	Monitoring Opioids in Europe: The Need for Shared Definitions and Measuring Drivers of Opioid Use and Related Harms. European Addiction Research, 2022, 28, 231-240.	2.4	12
65	Continuity of information in general practice. Scandinavian Journal of Primary Health Care, 2003, 21, 21-26.	1.5	11
66	Assessment of Appropriateness of Screening Communityâ€Dwelling Older People to Prevent Functional Decline. Journal of the American Geriatrics Society, 2012, 60, 42-50.	2.6	11
67	Timely identification of patients in need of palliative care using the Double Surprise Question: A prospective study on outpatients with cancer. Palliative Medicine, 2021, 35, 592-602.	3.1	11
68	Physician's prescribing preference as an instrumental variable. Epidemiology, 2015, 27, 1.	2.7	11
69	Experienced continuity of care in patients at risk for depression in primary care. European Journal of General Practice, 2014, 20, 161-166.	2.0	10
70	Effect of stepped care on health outcomes in patients with osteoarthritis: an observational study in Dutch general practice. British Journal of General Practice, 2014, 64, e538-e544.	1.4	10
71	Most Important Frequently Asked Questions From Patients With Hip or Knee Osteoarthritis: A Bestâ€Worst Scaling Exercise. Arthritis Care and Research, 2019, 71, 885-892.	3.4	10
72	Preparing for an orthopedic consultation using an eHealth tool: a randomized controlled trial in patients with hip and knee osteoarthritis. BMC Medical Informatics and Decision Making, 2020, 20, 92.	3.0	9

#	Article	IF	CITATIONS
73	First experiences with a twoâ€ <del>s</del> tep method for discussing goals with communityâ€dwelling frail older people. Health Expectations, 2015, 18, 1559-1566.	2.6	8
74	Development of the "Treatment beliefs in knee and hip OsteoArthritis (TOA)―questionnaire. BMC Musculoskeletal Disorders, 2017, 18, 402.	1.9	8
75	Continuity of care through medical recordsan explorative study on GPs' management considerations. Family Practice, 2006, 23, 349-352.	1.9	7
76	Primary care research–an international responsibility. Family Practice, 2012, 29, 499-500.	1.9	6
77	Mirror meetings with frail older people and multidisciplinary primary care teams: Process and impact analysis. Health Expectations, 2019, 22, 993-1002.	2.6	6
78	Advance care planning with people with dementia: a process evaluation of an educational intervention for general practitioners. BMC Family Practice, 2020, 21, 199.	2.9	6
79	General practitioners' views on the influence of long-term care reforms on integrated elderly care in the Netherlands: a qualitative interview study. Health Policy, 2021, 125, 930-940.	3.0	6
80	An Electronic Health Tool to Prepare for the First Orthopedic Consultation: Use and Usability Study. JMIR Formative Research, 2019, 3, e13577.	1.4	6
81	Healthcare utilisation of patients with cholecystolithiasis in primary care: a multipractice comparative analysis. BMJ Open, 2021, 11, e053188.	1.9	6
82	Which questionnaire to use when measuring continuity of care. Journal of Clinical Epidemiology, 2012, 65, 577-578.	5.0	5
83	Depressive disorder in the last phase of life in patients with cardiovascular disease, cancer, and COPD: data from a 20-year follow-up period in general practice. British Journal of General Practice, 2013, 63, e303-e308.	1.4	5
84	Respiratory Tract Infection Management and Antibiotic Prescription in Children: A Unique Study Comparing Three Levels of Healthcare in The Netherlands. Pediatric Infectious Disease Journal, 2021, 40, e100-e105.	2.0	5
85	Assessing the accuracy of a computerized decision support system for digoxin dosing in primary care: an observational study. Journal of Clinical Pharmacy and Therapeutics, 2005, 30, 279-283.	1.5	4
86	A different way of looking at depression. International Journal of Clinical Practice, 2010, 64, 1493-1495.	1.7	4
87	The association between implementation and outcome of a complex care program for frail elderly people. Family Practice, 2018, 35, 47-52.	1.9	4
88	Primary care functioning scale showed validity and reliability in patients with chronic conditions: a psychometric study. Journal of Clinical Epidemiology, 2020, 125, 130-137.	5.0	4
89	Addressing Palliative Sedation during Expert Consultation: A Descriptive Analysis of the Practice of Dutch Palliative Care Consultation Teams. PLoS ONE, 2015, 10, e0136309.	2.5	4
90	GPs' attitudes to personal continuity: findings from everyday practice differ from postal surveys. British Journal of General Practice, 2006, 56, 536-8.	1.4	4

#	Article	IF	CITATIONS
91	Recently enlisted patients in general practice use more health care resources. BMC Family Practice, 2007, 8, 64.	2.9	3
92	Palliative care consultations in primary care: a cross-sectional survey among Dutch general practitioners. BMJ Supportive and Palliative Care, 2016, 9, bmjspcare-2015-000967.	1.6	3
93	General practitioners' evaluations of optimal timing to initiate advance care planning for patients with cancer, organ failure, or multimorbidity: A health records survey study. Palliative Medicine, 2022, 36, 510-518.	3.1	3
94	Roles of general practitioners in shared decisionâ€making for patients with cancer: A qualitative study. European Journal of Cancer Care, 2022, 31, e13594.	1.5	3
95	The Watchman. Annals of Internal Medicine, 2012, 156, 602.	3.9	2
96	General practitioners' adherence to work-up and referral recommendations in fertility care. Human Reproduction, 2017, 32, 1249-1257.	0.9	2
97	Reasons for treatment choices in knee and hip osteoarthritis: a qualitative study. Osteoarthritis and Cartilage, 2016, 24, S489.	1.3	1
98	Incidence, presentation and management of Lyme disease in Dutch general practice. Family Practice, 2019, 36, 110-116.	1.9	1
99	Optimising personal continuity for older patients in general practice: a study protocol for a cluster randomised stepped wedge pragmatic trial. BMC Family Practice, 2021, 22, 207.	2.9	1
100	Assessment of functioning in Dutch primary care: Development study of a consultation tool for patients with chronic conditions and multimorbidity. Health Expectations, 0, , .	2.6	1
101	THU0204â€Factors Associated with the Setting and Content of Care in Patients with Osteoarthritis of the Hip or Knee. Annals of the Rheumatic Diseases, 2014, 73, 252.1-252.	0.9	0
102	FRI0638-HPRâ€Reasons Underlying Treatment Choices in Knee and Hip Osteoarthritis: A Qualitative Study. Annals of the Rheumatic Diseases, 2015, 74, 1332.2-1332.	0.9	0
103	OP0098-HPRâ€Structure and Importance of Patients' Reasons for Treatment Choices in HIP and Knee Osteoarthritis: a Concept Mapping Study. Annals of the Rheumatic Diseases, 2015, 74, 105.3-106.	0.9	0
104	Structure and importance of patients' reasons for treatment choices in hip and knee osteoarthritis: a concept mapping study. Osteoarthritis and Cartilage, 2016, 24, S485-S486.	1.3	0
105	AB1087-HPRâ€Non-Pharmacological, Non-Surgical Care in Hip and Knee Osteoarthritis: The View of Healthcare Providers. Annals of the Rheumatic Diseases, 2016, 75, 1303.2-1303.	0.9	0
106	Non-pharmacological, non-surgical care in hip and knee osteoarthritis: the view of healthcare providers. Osteoarthritis and Cartilage, 2016, 24, S494.	1.3	0
107	THU0752-HPRâ€Development of the "treatment beliefs in knee and hip osteoarthritis (TOA)― questionnaire. , 2017, , .		0
108	OP0138-HPRâ€Do patients' treatment beliefs affect treatment choices in knee and hip osteoarthritis?. , 2017, , .		0

#	Article	IF	CITATIONS
109	THU0430â€PREPARING AN ORTHOPAEDIC CONSULTATION USING AN EHEALTH TOOL: A RANDOMIZED CONTROLLED TRIAL IN PATIENTS WITH HIP AND KNEE OSTEOARTHRITIS. , 2019, , .		0
110	AB0789â€AN EHEALTH TOOL TO PREPARE A FIRST ORTHOPAEDIC CONSULTATIONS: A USE AND USABILITY STUDY. , 2019, , .		0
111	The perspective of frail elderly on their care networks in primary care. International Journal of Integrated Care, 2016, 16, 45.	0.2	0
112	The role of the general practitioner in multidisciplinary teams: a qualitative study in elderly care. International Journal of Integrated Care, 2018, 18, 259.	0.2	0