

Henk J Schers

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6875433/publications.pdf>

Version: 2024-02-01

112
papers

2,991
citations

159585

30
h-index

197818

49
g-index

124
all docs

124
docs citations

124
times ranked

4294
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Monitoring Opioids in Europe: The Need for Shared Definitions and Measuring Drivers of Opioid Use and Related Harms. <i>European Addiction Research</i> , 2022, 28, 231-240. | 2.4 | 12 |
| 2 | General practitioners'™ evaluations of optimal timing to initiate advance care planning for patients with cancer, organ failure, or multimorbidity: A health records survey study. <i>Palliative Medicine</i> , 2022, 36, 510-518. | 3.1 | 3 |
| 3 | Roles of general practitioners in shared decision-making for patients with cancer: A qualitative study. <i>European Journal of Cancer Care</i> , 2022, 31, e13594. | 1.5 | 3 |
| 4 | Comprehensive Health Assessment 3 Months After Recovery From Acute Coronavirus Disease 2019 (COVID-19). <i>Clinical Infectious Diseases</i> , 2021, 73, e1089-e1098. | 5.8 | 332 |
| 5 | Timely identification of patients in need of palliative care using the Double Surprise Question: A prospective study on outpatients with cancer. <i>Palliative Medicine</i> , 2021, 35, 592-602. | 3.1 | 11 |
| 6 | General practitioners'™ views on the influence of long-term care reforms on integrated elderly care in the Netherlands: a qualitative interview study. <i>Health Policy</i> , 2021, 125, 930-940. | 3.0 | 6 |
| 7 | Purposeful Incorporation of Patient Narratives in the Medical Record in the Netherlands. <i>Journal of the American Board of Family Medicine</i> , 2021, 34, 709-723. | 1.5 | 13 |
| 8 | The COVID-19 Pandemic in Nijmegen, the Netherlands: Changes in Presented Health Problems and Demand for Primary Care. <i>Annals of Family Medicine</i> , 2021, 19, 44-47. | 1.9 | 33 |
| 9 | Respiratory Tract Infection Management and Antibiotic Prescription in Children: A Unique Study Comparing Three Levels of Healthcare in The Netherlands. <i>Pediatric Infectious Disease Journal</i> , 2021, 40, e100-e105. | 2.0 | 5 |
| 10 | Optimising personal continuity for older patients in general practice: a study protocol for a cluster randomised stepped wedge pragmatic trial. <i>BMC Family Practice</i> , 2021, 22, 207. | 2.9 | 1 |
| 11 | Healthcare utilisation of patients with cholecystolithiasis in primary care: a multipractice comparative analysis. <i>BMJ Open</i> , 2021, 11, e053188. | 1.9 | 6 |
| 12 | Educating Dutch General Practitioners in Dementia Advance Care Planning: A Cluster Randomized Controlled Trial. <i>Journal of the American Medical Directors Association</i> , 2020, 21, 837-842.e4. | 2.5 | 36 |
| 13 | Advance care planning with people with dementia: a process evaluation of an educational intervention for general practitioners. <i>BMC Family Practice</i> , 2020, 21, 199. | 2.9 | 6 |
| 14 | Preparing for an orthopedic consultation using an eHealth tool: a randomized controlled trial in patients with hip and knee osteoarthritis. <i>BMC Medical Informatics and Decision Making</i> , 2020, 20, 92. | 3.0 | 9 |
| 15 | Primary care functioning scale showed validity and reliability in patients with chronic conditions: a psychometric study. <i>Journal of Clinical Epidemiology</i> , 2020, 125, 130-137. | 5.0 | 4 |
| 16 | Professional Care Networks of Frail Older People: An Explorative Survey Study from the Patient Perspective. <i>International Journal of Integrated Care</i> , 2020, 20, 1. | 0.2 | 12 |
| 17 | Most Important Frequently Asked Questions From Patients With Hip or Knee Osteoarthritis: A Best-Worst Scaling Exercise. <i>Arthritis Care and Research</i> , 2019, 71, 885-892. | 3.4 | 10 |
| 18 | Incidence, presentation and management of Lyme disease in Dutch general practice. <i>Family Practice</i> , 2019, 36, 110-116. | 1.9 | 1 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Mirror meetings with frail older people and multidisciplinary primary care teams: Process and impact analysis. <i>Health Expectations</i> , 2019, 22, 993-1002. | 2.6 | 6 |
| 20 | THU0430â€¦PREPARING AN ORTHOPAEDIC CONSULTATION USING AN EHEALTH TOOL: A RANDOMIZED CONTROLLED TRIAL IN PATIENTS WITH HIP AND KNEE OSTEOARTHRITIS. , 2019, , . | | 0 |
| 21 | AB0789â€¦AN EHEALTH TOOL TO PREPARE A FIRST ORTHOPAEDIC CONSULTATIONS: A USE AND USABILITY STUDY. , 2019, , . | | 0 |
| 22 | Advance care planning for patients with cancer in the palliative phase in Dutch general practices. <i>Family Practice</i> , 2019, 36, 587-593. | 1.9 | 19 |
| 23 | An Electronic Health Tool to Prepare for the First Orthopedic Consultation: Use and Usability Study. <i>JMIR Formative Research</i> , 2019, 3, e13577. | 1.4 | 6 |
| 24 | Treatment Beliefs Underlying Intended Treatment Choices in Knee and Hip Osteoarthritis. <i>International Journal of Behavioral Medicine</i> , 2018, 25, 198-206. | 1.7 | 12 |
| 25 | The association between implementation and outcome of a complex care program for frail elderly people. <i>Family Practice</i> , 2018, 35, 47-52. | 1.9 | 4 |
| 26 | Cost-effectiveness of a multicomponent primary care program targeting frail elderly people. <i>BMC Family Practice</i> , 2018, 19, 62. | 2.9 | 15 |
| 27 | Preliminary effects of a regional approached multidisciplinary educational program on healthcare utilization in patients with hip or knee osteoarthritis: an observational study. <i>BMC Family Practice</i> , 2018, 19, 82. | 2.9 | 14 |
| 28 | The role of the general practitioner in multidisciplinary teams: a qualitative study in elderly care. <i>BMC Family Practice</i> , 2018, 19, 40. | 2.9 | 39 |
| 29 | The development of an ICF-based questionnaire for patients with chronic conditions in primary care. <i>Journal of Clinical Epidemiology</i> , 2018, 103, 92-100. | 5.0 | 19 |
| 30 | Incidence, course and risk factors of head injury: a retrospective cohort study. <i>BMJ Open</i> , 2018, 8, e020364. | 1.9 | 16 |
| 31 | The role of the general practitioner in multidisciplinary teams: a qualitative study in elderly care. <i>International Journal of Integrated Care</i> , 2018, 18, 259. | 0.2 | 0 |
| 32 | Factors Associated With Referral to Secondary Care in Patients With Osteoarthritis of the Hip or Knee After Implementation of a Steppedâ€œCare Strategy. <i>Arthritis Care and Research</i> , 2017, 69, 216-225. | 3.4 | 17 |
| 33 | General practitionersâ€™ adherence to work-up and referral recommendations in fertility care. <i>Human Reproduction</i> , 2017, 32, 1249-1257. | 0.9 | 2 |
| 34 | Barriers Impeding the Use of Non-pharmacological, Non-surgical Care in Hip and Knee Osteoarthritis. <i>Journal of Clinical Rheumatology</i> , 2017, 23, 405-410. | 0.9 | 46 |
| 35 | THU0752-HPRâ€¦Development of the â€œtreatment beliefs in knee and hip osteoarthritis (TOA)â€œ questionnaire. , 2017, , . | | 0 |
| 36 | The effect of weekly specialist palliative care teleconsultations in patients with advanced cancer â€œa randomized clinical trial. <i>BMC Medicine</i> , 2017, 15, 119. | 5.5 | 80 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Hierarchical structure and importance of patients' reasons for treatment choices in knee and hip osteoarthritis: a concept mapping study. <i>Rheumatology</i> , 2017, 56, 271-278. | 1.9 | 12 |
| 38 | Development of the 'Treatment beliefs in knee and hip OsteoArthritis (TOA)' questionnaire. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 402. | 1.9 | 8 |
| 39 | OP0138-HPR...Do patients' treatment beliefs affect treatment choices in knee and hip osteoarthritis?. , 2017, , . | | 0 |
| 40 | The Diagnostic Value of the Patient's Reason for Encounter for Diagnosing Cancer in Primary Care. <i>Journal of the American Board of Family Medicine</i> , 2017, 30, 806-812. | 1.5 | 16 |
| 41 | Reasons for Treatment Choices in Knee and Hip Osteoarthritis: A Qualitative Study. <i>Arthritis Care and Research</i> , 2016, 68, 1260-1267. | 3.4 | 44 |
| 42 | Multicomponent Program to Reduce Functional Decline in Frail Elderly People: A Cluster Controlled Trial. <i>Journal of the American Board of Family Medicine</i> , 2016, 29, 209-217. | 1.5 | 52 |
| 43 | Palliative care consultations in primary care: a cross-sectional survey among Dutch general practitioners. <i>BMJ Supportive and Palliative Care</i> , 2016, 9, bmjspcare-2015-000967. | 1.6 | 3 |
| 44 | Structure and importance of patients' reasons for treatment choices in hip and knee osteoarthritis: a concept mapping study. <i>Osteoarthritis and Cartilage</i> , 2016, 24, S485-S486. | 1.3 | 0 |
| 45 | Reasons for treatment choices in knee and hip osteoarthritis: a qualitative study. <i>Osteoarthritis and Cartilage</i> , 2016, 24, S489. | 1.3 | 1 |
| 46 | AB1087-HPR...Non-Pharmacological, Non-Surgical Care in Hip and Knee Osteoarthritis: The View of Healthcare Providers. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 1303.2-1303. | 0.9 | 0 |
| 47 | Non-pharmacological, non-surgical care in hip and knee osteoarthritis: the view of healthcare providers. <i>Osteoarthritis and Cartilage</i> , 2016, 24, S494. | 1.3 | 0 |
| 48 | The perspective of frail elderly on their care networks in primary care. <i>International Journal of Integrated Care</i> , 2016, 16, 45. | 0.2 | 0 |
| 49 | Predictive validity of a two-step tool to map frailty in primary care. <i>BMC Medicine</i> , 2015, 13, 287. | 5.5 | 27 |
| 50 | FRIO638-HPR...Reasons Underlying Treatment Choices in Knee and Hip Osteoarthritis: A Qualitative Study. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 1332.2-1332. | 0.9 | 0 |
| 51 | OP0098-HPR...Structure and Importance of Patients' Reasons for Treatment Choices in HIP and Knee Osteoarthritis: a Concept Mapping Study. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 105.3-106. | 0.9 | 0 |
| 52 | Diagnosis of Frailty after a Comprehensive Geriatric Assessment: Differences between Family Physicians and Geriatricians. <i>Journal of the American Board of Family Medicine</i> , 2015, 28, 240-248. | 1.5 | 18 |
| 53 | First experiences with a two-step method for discussing goals with community-dwelling frail older people. <i>Health Expectations</i> , 2015, 18, 1559-1566. | 2.6 | 8 |
| 54 | Psychosocial, musculoskeletal and somatoform comorbidity in patients with chronic low back pain: original results from the Dutch Transition Project. <i>Family Practice</i> , 2015, 32, 297-304. | 1.9 | 18 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | International variation in GP treatment strategies for subclinical hypothyroidism in older adults: a case-based survey. <i>British Journal of General Practice</i> , 2015, 65, e121-e132. | 1.4 | 17 |
| 56 | The predictive value of gait speed and maximum step length for falling in community-dwelling older persons. <i>Age and Ageing</i> , 2015, 44, 294-299. | 1.6 | 29 |
| 57 | Physician's prescribing preference as an instrumental variable. <i>Epidemiology</i> , 2015, 27, 1. | 2.7 | 11 |
| 58 | Addressing Palliative Sedation during Expert Consultation: A Descriptive Analysis of the Practice of Dutch Palliative Care Consultation Teams. <i>PLoS ONE</i> , 2015, 10, e0136309. | 2.5 | 4 |
| 59 | Experienced continuity of care in patients at risk for depression in primary care. <i>European Journal of General Practice</i> , 2014, 20, 161-166. | 2.0 | 10 |
| 60 | Health Care Use of Patients With Osteoarthritis of the Hip or Knee After Implementation of a Stepped-Care Strategy: An Observational Study. <i>Arthritis Care and Research</i> , 2014, 66, 817-827. | 3.4 | 42 |
| 61 | Increased incidence of squamous cell carcinoma of the skin after long-term treatment with azathioprine in patients with autoimmune inflammatory rheumatic diseases. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2014, 28, 27-33. | 2.4 | 34 |
| 62 | Feasibility evaluation of a stepped procedure to identify community-dwelling frail older people in general practice. A mixed methods study. <i>European Journal of General Practice</i> , 2014, 20, 107-113. | 2.0 | 13 |
| 63 | Is the quality of data in an electronic medical record sufficient for assessing the quality of primary care?. <i>Journal of the American Medical Informatics Association: JAMIA</i> , 2014, 21, 692-698. | 4.4 | 30 |
| 64 | Non-surgical care in patients with hip or knee osteoarthritis is modestly consistent with a stepped care strategy after its implementation. <i>International Journal for Quality in Health Care</i> , 2014, 26, 490-498. | 1.8 | 21 |
| 65 | Construct validity and reliability of a two-step tool for the identification of frail older people in primary care. <i>Journal of Clinical Epidemiology</i> , 2014, 67, 176-183. | 5.0 | 36 |
| 66 | Effect of stepped care on health outcomes in patients with osteoarthritis: an observational study in Dutch general practice. <i>British Journal of General Practice</i> , 2014, 64, e538-e544. | 1.4 | 10 |
| 67 | THU0204...Factors Associated with the Setting and Content of Care in Patients with Osteoarthritis of the Hip or Knee. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 252.1-252. | 0.9 | 0 |
| 68 | Evaluation of an eHealth Intervention in Chronic Care for Frail Older People: Why Adherence is the First Target. <i>Journal of Medical Internet Research</i> , 2014, 16, e156. | 4.3 | 46 |
| 69 | Which Frail Older Patients Use Online Health Communities and Why? A Mixed Methods Process Evaluation of Use of the Health and Welfare Portal. <i>Journal of Medical Internet Research</i> , 2014, 16, e278. | 4.3 | 18 |
| 70 | Agreement of general practitioners with the guideline-based stepped-care strategy for patients with osteoarthritis of the hip or knee: a cross-sectional study. <i>BMC Family Practice</i> , 2013, 14, 33. | 2.9 | 21 |
| 71 | Incidence, Causative Mechanisms, and Anatomic Localization of Stroke in Pituitary Adenoma Patients Treated With Postoperative Radiation Therapy Versus Surgery Alone. <i>International Journal of Radiation Oncology Biology Physics</i> , 2013, 87, 53-59. | 0.8 | 31 |
| 72 | Patient reported barriers and facilitators to using a self-management booklet for hip and knee osteoarthritis in primary care: results of a qualitative interview study. <i>BMC Family Practice</i> , 2013, 14, 181. | 2.9 | 24 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Development of an instrument for the identification of frail older people as a target population for integrated care. <i>British Journal of General Practice</i> , 2013, 63, e225-e231. | 1.4 | 54 |
| 74 | Depressive disorder in the last phase of life in patients with cardiovascular disease, cancer, and COPD: data from a 20-year follow-up period in general practice. <i>British Journal of General Practice</i> , 2013, 63, e303-e308. | 1.4 | 5 |
| 75 | How unique is continuity of care? A review of continuity and related concepts. <i>Family Practice</i> , 2012, 29, 264-271. | 1.9 | 117 |
| 76 | Continuity in different care modes and its relationship to quality of life: a randomised controlled trial in patients with COPD. <i>British Journal of General Practice</i> , 2012, 62, e422-e428. | 1.4 | 22 |
| 77 | Health Care in The Netherlands. <i>Journal of the American Board of Family Medicine</i> , 2012, 25, S12-S17. | 1.5 | 71 |
| 78 | Preferences for receiving information among frail older adults and their informal caregivers: a qualitative study. <i>Family Practice</i> , 2012, 29, 742-747. | 1.9 | 35 |
| 79 | Primary care research—an international responsibility. <i>Family Practice</i> , 2012, 29, 499-500. | 1.9 | 6 |
| 80 | Which questionnaire to use when measuring continuity of care. <i>Journal of Clinical Epidemiology</i> , 2012, 65, 577-578. | 5.0 | 5 |
| 81 | Impact of Interprofessional Education on Collaboration Attitudes, Skills, and Behavior Among Primary Care Professionals. <i>Journal of Continuing Education in the Health Professions</i> , 2012, 32, 196-204. | 1.3 | 79 |
| 82 | The CareWell-primary care program: design of a cluster controlled trial and process evaluation of a complex intervention targeting community-dwelling frail elderly. <i>BMC Family Practice</i> , 2012, 13, 115. | 2.9 | 22 |
| 83 | Heart failure patients' experiences with continuity of care and its relation to medication adherence: a cross-sectional study. <i>BMC Family Practice</i> , 2012, 13, 86. | 2.9 | 23 |
| 84 | Implementation of an innovative web-based conference table for community-dwelling frail older people, their informal caregivers and professionals: a process evaluation. <i>BMC Health Services Research</i> , 2012, 12, 251. | 2.2 | 43 |
| 85 | Home visits for frail older people: a qualitative study on the needs and preferences of frail older people and their informal caregivers. <i>British Journal of General Practice</i> , 2012, 62, e554-e560. | 1.4 | 46 |
| 86 | The Watchman. <i>Annals of Internal Medicine</i> , 2012, 156, 602. | 3.9 | 2 |
| 87 | Measuring continuity of care: psychometric properties of the Nijmegen Continuity Questionnaire. <i>British Journal of General Practice</i> , 2012, 62, e949-e957. | 1.4 | 39 |
| 88 | Assessment of Appropriateness of Screening Community-Dwelling Older People to Prevent Functional Decline. <i>Journal of the American Geriatrics Society</i> , 2012, 60, 42-50. | 2.6 | 11 |
| 89 | Measurement Properties of Questionnaires Measuring Continuity of Care: A Systematic Review. <i>PLoS ONE</i> , 2012, 7, e42256. | 2.5 | 70 |
| 90 | Filling the Gaps in a Fragmented Health Care System: Development of the Health and Welfare Information Portal (ZWIP). <i>JMIR Research Protocols</i> , 2012, 1, e10. | 1.0 | 25 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Nijmegen Continuity Questionnaire: Development and testing of a questionnaire that measures continuity of care. <i>Journal of Clinical Epidemiology</i> , 2011, 64, 1391-1399. | 5.0 | 71 |
| 92 | “Beating osteoARthritis”: Development of a stepped care strategy to optimize utilization and timing of non-surgical treatment modalities for patients with hip or knee osteoarthritis. <i>Clinical Rheumatology</i> , 2011, 30, 1623-1629. | 2.2 | 102 |
| 93 | Study protocol: optimization of complex palliative care at home via telemedicine. A cluster randomized controlled trial. <i>BMC Palliative Care</i> , 2011, 10, 13. | 1.8 | 16 |
| 94 | A different way of looking at depression. <i>International Journal of Clinical Practice</i> , 2010, 64, 1493-1495. | 1.7 | 4 |
| 95 | Continuity of care preferably measured from the patients' perspective. <i>Journal of Clinical Epidemiology</i> , 2010, 63, 998-999. | 5.0 | 21 |
| 96 | Structured chronic primary care and health-related quality of life in chronic heart failure. <i>BMC Health Services Research</i> , 2009, 9, 104. | 2.2 | 15 |
| 97 | Skin Diseases in Family Medicine: Prevalence and Health Care Use. <i>Annals of Family Medicine</i> , 2008, 6, 349-354. | 1.9 | 67 |
| 98 | What went and what came? Morbidity trends in general practice from the Netherlands. <i>European Journal of General Practice</i> , 2008, 14, 13-24. | 2.0 | 35 |
| 99 | Recently enlisted patients in general practice use more health care resources. <i>BMC Family Practice</i> , 2007, 8, 64. | 2.9 | 3 |
| 100 | Prevalence of physical symptoms of itch, pain and fatigue in patients with skin diseases in general practice. <i>British Journal of Dermatology</i> , 2007, 156, 1346-1349. | 1.5 | 106 |
| 101 | Psychosocial well-being of patients with skin diseases in general practice. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2007, 21, 070206173308001-??? | 2.4 | 27 |
| 102 | Continuity of care through medical records—an explorative study on GPs' management considerations. <i>Family Practice</i> , 2006, 23, 349-352. | 1.9 | 7 |
| 103 | GPs' attitudes to personal continuity: findings from everyday practice differ from postal surveys. <i>British Journal of General Practice</i> , 2006, 56, 536-8. | 1.4 | 4 |
| 104 | Assessing the accuracy of a computerized decision support system for digoxin dosing in primary care: an observational study. <i>Journal of Clinical Pharmacy and Therapeutics</i> , 2005, 30, 279-283. | 1.5 | 4 |
| 105 | Continuity of Care: Is the Personal Doctor Still Important? A Survey of General Practitioners and Family Physicians in England and Wales, the United States, and the Netherlands. <i>Annals of Family Medicine</i> , 2005, 3, 353-359. | 1.9 | 101 |
| 106 | Patients' Needs for Contact With Their GP at the Time of Hospital Admission and Other Life Events: A Quantitative and Qualitative Exploration. <i>Annals of Family Medicine</i> , 2004, 2, 462-468. | 1.9 | 18 |
| 107 | Familiarity with a GP and patients' evaluations of care. A cross-sectional study. <i>Family Practice</i> , 2004, 22, 15-19. | 1.9 | 71 |
| 108 | Family medicine trainees still value continuity of care. <i>Family Medicine</i> , 2004, 36, 51-4. | 0.5 | 17 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Continuity of information in general practice. <i>Scandinavian Journal of Primary Health Care</i> , 2003, 21, 21-26. | 1.5 | 11 |
| 110 | Continuity of care in general practice: a survey of patients' views. <i>British Journal of General Practice</i> , 2002, 52, 459-62. | 1.4 | 66 |
| 111 | Implementation Barriers for General Practice Guidelines on Low Back Pain. <i>Spine</i> , 2001, 26, E348-E353. | 2.0 | 142 |
| 112 | Assessment of functioning in Dutch primary care: Development study of a consultation tool for patients with chronic conditions and multimorbidity. <i>Health Expectations</i> , 0, , . | 2.6 | 1 |