

Paola Dugo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6908580/publications.pdf>

Version: 2024-02-01

331
papers

13,944
citations

19657

61
h-index

45317

90
g-index

340
all docs

340
docs citations

340
times ranked

11068
citing authors

#	ARTICLE	IF	CITATIONS
1	Phytochemical Characterization of <i>Rhus coriaria</i> L. Extracts by Headspace Solid-Phase Micro Extraction Gas Chromatography, Comprehensive Two-Dimensional Liquid Chromatography, and Antioxidant Activity Evaluation. <i>Molecules</i> , 2022, 27, 1727.	3.8	15
2	Comparison of lipid profile of Italian Extra Virgin Olive Oils by using rapid chromatographic approaches. <i>Journal of Food Composition and Analysis</i> , 2022, 110, 104531.	3.9	4
3	Supercritical fluid chromatography-tandem mass spectrometry of oxygen heterocyclic compounds in Citrus essential oils. <i>Analytical and Bioanalytical Chemistry</i> , 2022, 414, 4821-4836.	3.7	4
4	Elucidation of the Lipid Composition of Hemp (<i>Cannabis sativa</i> L.) Products by Means of Gas Chromatography and Ultra-High Performance Liquid Chromatography Coupled to Mass Spectrometry Detection. <i>Molecules</i> , 2022, 27, 3358.	3.8	16
5	Determination of the polyphenolic content of <i>Ammodaucus leucotrichus</i> Cosson and Durieu by liquid chromatography coupled with mass spectrometry and evaluation of the antioxidant and antiglycation properties. <i>Journal of Separation Science</i> , 2022, 45, 3301-3309.	2.5	7
6	Apocarotenoids profiling in different <i>Capsicum</i> species. <i>Food Chemistry</i> , 2021, 334, 127595.	8.2	24
7	Multidimensional liquid chromatography approaches for analysis of food contaminants. <i>Journal of Separation Science</i> , 2021, 44, 17-34.	2.5	15
8	Comprehensive two-dimensional liquid chromatography-based qualitative screening of aqueous phases from pyrolysis bio-oils. <i>Electrophoresis</i> , 2021, 42, 58-67.	2.4	15
9	Reliable identification and quantification of anabolic androgenic steroids in dietary supplements by using gas chromatography coupled to triple quadrupole mass spectrometry. <i>Drug Testing and Analysis</i> , 2021, 13, 128-139.	2.6	9
10	<i>Cannabis Sativa</i> L.: a comprehensive review on the analytical methodologies for cannabinoids and terpenes characterization. <i>Journal of Chromatography A</i> , 2021, 1637, 461864.	3.7	49
11	Influence of Citrus Flavor Addition in Brewing Process: Characterization of the Volatile and Non-Volatile Profile to Prevent Frauds and Adulterations. <i>Separations</i> , 2021, 8, 18.	2.4	13
12	Development of a Novel Microwave Distillation Technique for the Isolation of <i>Cannabis sativa</i> L. Essential Oil and Gas Chromatography Analyses for the Comprehensive Characterization of Terpenes and Terpenoids, Including Their Enantio-Distribution. <i>Molecules</i> , 2021, 26, 1588.	3.8	20
13	Identification of high-value generating molecules from the wastes of tuna fishery industry by liquid chromatography and gas chromatography hyphenated techniques with automated sample preparation. <i>Journal of Separation Science</i> , 2021, 44, 1571-1580.	2.5	15
14	The retention index approach in liquid chromatography: An historical review and recent advances. <i>Journal of Chromatography A</i> , 2021, 1640, 461963.	3.7	18
15	Characterization of <i>Rubus fruticosus</i> L. berries growing wild in Morocco: phytochemical screening, antioxidant activity and chromatography analysis. <i>European Food Research and Technology</i> , 2021, 247, 1689-1699.	3.3	6
16	Reversed phase versus hydrophilic interaction liquid chromatography as first dimension of comprehensive two-dimensional liquid chromatography systems for the elucidation of the polyphenolic content of food and natural products. <i>Journal of Chromatography A</i> , 2021, 1645, 462129.	3.7	28
17	Pattern-Type Separation of Triacylglycerols by Silver Thiolate-Non-Aqueous Reversed Phase Comprehensive Liquid Chromatography. <i>Separations</i> , 2021, 8, 88.	2.4	11
18	Dietary Intake of Coumarins and Furocoumarins through Citrus Beverages: A Detailed Estimation by a HPLC-MS/MS Method Combined with the Linear Retention Index System. <i>Foods</i> , 2021, 10, 1533.	4.3	13

#	ARTICLE	IF	CITATIONS
19	Linear retention index approach applied to liquid chromatography coupled to triple quadrupole mass spectrometry to determine oxygen heterocyclic compounds at trace level in finished cosmetics. <i>Journal of Chromatography A</i> , 2021, 1649, 462183.	3.7	15
20	The Digestibility of Hibiscus sabdariffa L. Polyphenols Using an In Vitro Human Digestion Model and Evaluation of Their Antimicrobial Activity. <i>Nutrients</i> , 2021, 13, 2360.	4.1	10
21	Determination of bioactive compounds in extra virgin olive oils from 19 Moroccan areas using liquid chromatography coupled to mass spectrometry: a study over two successive years. <i>European Food Research and Technology</i> , 2021, 247, 2993-3012.	3.3	16
22	Coumarins, Psoralens and Polymethoxyflavones in Cold-pressed Citrus Essential Oils: a Review. <i>Journal of Essential Oil Research</i> , 2021, 33, 221-239.	2.7	18
23	Comparative study of the phenolic profile, antioxidant and antimicrobial activities of leaf extracts of five <i>Juniperus</i> L. (Cupressaceae) taxa growing in Turkey. <i>Natural Product Research</i> , 2020, 34, 1636-1641.	1.8	25
24	Evaluation of antioxidant and anti-inflammatory activity of green coffee beans methanolic extract in rat skin. <i>Natural Product Research</i> , 2020, 34, 1535-1541.	1.8	24
25	Characterization of the polyphenolic fraction of pomegranate samples by comprehensive two-dimensional liquid chromatography coupled to mass spectrometry detection. <i>Natural Product Research</i> , 2020, 34, 39-45.	1.8	34
26	<i>Inula viscosa</i> (L.) Aiton leaves and flower buds: Effect of extraction solvent/technique on their antioxidant ability, antimicrobial properties and phenolic profile. <i>Natural Product Research</i> , 2020, 34, 46-52.	1.8	22
27	Combining linear retention index and electron ionization mass spectrometry for a reliable identification in nano liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1610, 460581.	3.7	17
28	<i>Silene vulgaris</i> subsp. <i>macrocarpa</i> leaves and roots from Morocco: assessment of the efficiency of different extraction techniques and solvents on their antioxidant capacity, brine shrimp toxicity and phenolic characterization. <i>Plant Biosystems</i> , 2020, 154, 692-699.	1.6	10
29	Characterization of monoacylglycerols and diacylglycerols rich in polyunsaturated fatty acids produced by hydrolysis of <i>Mustelus mustelus</i> liver oil catalyzed by an immobilized bacterial lipase. <i>Journal of Chromatography A</i> , 2020, 1613, 460692.	3.7	9
30	Application of compressed fluid-based extraction and purification procedures to obtain astaxanthin-enriched extracts from <i>Haematococcus pluvialis</i> and characterization by comprehensive two-dimensional liquid chromatography coupled to mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2020, 412, 589-599.	3.7	19
31	Characterization of Phenolic Compounds, Vitamin E and Fatty Acids from Monovarietal Virgin Olive Oils of <i>Picholine marocaine</i> Cultivar. <i>Molecules</i> , 2020, 25, 5428.	3.8	15
32	Physico-Chemical and Phytochemical Characterization of Moroccan Wild Jujube <i>Zizyphus lotus</i> (L.) Fruit Crude Extract and Fractions. <i>Molecules</i> , 2020, 25, 5237.	3.8	14
33	Polyphenolic compounds with biological activity in guabiroba fruits (<i>Campomanesia</i>) <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50</i> 2020, 41, 1784-1792.	2.4	19
34	Concentration of Potentially Bioactive Compounds in Italian Extra Virgin Olive Oils from Various Sources by Using LC-MS and Multivariate Data Analysis. <i>Foods</i> , 2020, 9, 1120.	4.3	20
35	Determination of the Metabolite Content of <i>Brassica juncea</i> Cultivars Using Comprehensive Two-Dimensional Liquid Chromatography Coupled with a Photodiode Array and Mass Spectrometry Detection. <i>Molecules</i> , 2020, 25, 1235.	3.8	29
36	Hyphenations of 2D capillary-based LC with mass spectrometry. , 2020, , 369-412.		1

#	ARTICLE	IF	CITATIONS
37	Flavors and odors analysis. , 2020, , 697-727.		0
38	Wild strawberry (<i>Arbutus unedo</i>): Phytochemical screening and antioxidant properties of fruits collected in northern Morocco. <i>Arabian Journal of Chemistry</i> , 2020, 13, 6299-6311.	4.9	18
39	Rapid and miniaturized qualitative and quantitative gas chromatography profiling of human blood total fatty acids. <i>Analytical and Bioanalytical Chemistry</i> , 2020, 412, 2327-2337.	3.7	23
40	Comprehensive two-dimensional liquid chromatography as a powerful tool for the analysis of food and food products. <i>TrAC - Trends in Analytical Chemistry</i> , 2020, 127, 115894.	11.4	52
41	Evaluation of Italian extra virgin olive oils based on the phenolic compounds composition using multivariate statistical methods. <i>European Food Research and Technology</i> , 2020, 246, 1241-1249.	3.3	11
42	Lipid profile of fish species by liquid chromatography coupled to mass spectrometry and a novel linear retention index database. <i>Journal of Separation Science</i> , 2020, 43, 1773-1780.	2.5	11
43	Evaluation of matrix effect in one-dimensional and comprehensive two-dimensional liquid chromatography for the determination of the phenolic fraction in extra virgin olive oils. <i>Journal of Separation Science</i> , 2020, 43, 1781-1789.	2.5	19
44	Fingerprinting of the Unsaponifiable Fraction of Vegetable Oils by Using Cryogenically-Modulated Comprehensive Two-Dimensional Gas Chromatography-High Resolution Time-of-Flight Mass Spectrometry. <i>Food Analytical Methods</i> , 2020, 13, 1523-1529.	2.6	12
45	Free carotenoids and carotenoids esters composition in Spanish orange and mandarin juices from diverse varieties. <i>Food Chemistry</i> , 2019, 300, 125139.	8.2	16
46	Oxygen heterocyclic compound screening in <i>Citrus</i> essential oils by linear retention index approach applied to liquid chromatography coupled to photodiode array detector. <i>Flavour and Fragrance Journal</i> , 2019, 34, 349-364.	2.6	12
47	Evaluation of the availability of delphinidin and cyanidin-3-O-sambubioside from <i>Hibiscus sabdariffa</i> and 6-gingerol from <i>Zingiber officinale</i> in colon using liquid chromatography and mass spectrometry detection. <i>European Food Research and Technology</i> , 2019, 245, 2425-2433.	3.3	9
48	High-performance liquid chromatography combined with electron ionization mass spectrometry: A review. <i>TrAC - Trends in Analytical Chemistry</i> , 2019, 118, 112-122.	11.4	54
49	The Phenolic Fraction of Italian Extra Virgin Olive Oils: Elucidation Through Combined Liquid Chromatography and NMR Approaches. <i>Food Analytical Methods</i> , 2019, 12, 1759-1770.	2.6	38
50	Determination of the polyphenolic fraction of <i>Pistacia vera</i> L. kernel extracts by comprehensive two-dimensional liquid chromatography coupled to mass spectrometry detection. <i>Analytical and Bioanalytical Chemistry</i> , 2019, 411, 4819-4829.	3.7	30
51	Blood orange (<i>Citrus sinensis</i>) as a rich source of nutraceuticals: investigation of bioactive compounds in different parts of the fruit by HPLC-PDA/MS. <i>Natural Product Research</i> , 2019, 35, 1-5.	1.8	18
52	Green Extraction Approaches for Carotenoids and Esters: Characterization of Native Composition from Orange Peel. <i>Antioxidants</i> , 2019, 8, 613.	5.1	37
53	Comprehensive lipid profiling in the Mediterranean mussel (<i>Mytilus galloprovincialis</i>) using hyphenated and multidimensional chromatography techniques coupled to mass spectrometry detection. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 3297-3313.	3.7	35
54	Use of an Online Extraction Technique Coupled to Liquid Chromatography for Determination of Caffeine in Coffee, Tea, and Cocoa. <i>Food Analytical Methods</i> , 2018, 11, 2637-2644.	2.6	17

#	ARTICLE	IF	CITATIONS
55	Proposal of a Linear Retention Index System for Improving Identification Reliability of Triacylglycerol Profiles in Lipid Samples by Liquid Chromatography Methods. <i>Analytical Chemistry</i> , 2018, 90, 3313-3320.	6.5	31
56	Untargeted profiling of <i>Glycyrrhiza glabra</i> extract with comprehensive two-dimensional liquid chromatography-mass spectrometry using multi-segmented shift gradients in the second dimension: Expanding the metabolic coverage. <i>Electrophoresis</i> , 2018, 39, 1993-2000.	2.4	27
57	Analysis of phenolic compounds in different parts of pomegranate (<i>Punica granatum</i>) fruit by HPLC-PDA-ESI/MS and evaluation of their antioxidant activity: application to different Italian varieties. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 3507-3520.	3.7	111
58	Partial characterization of the pigments produced by the marine-derived fungus <i>Talaromyces albobiverticillius</i> 30548. Towards a new fungal red colorant for the food industry. <i>Journal of Food Composition and Analysis</i> , 2018, 67, 38-47.	3.9	53
59	Recent Analytical Techniques Advances in the Carotenoids and Their Derivatives Determination in Various Matrixes. <i>Journal of Agricultural and Food Chemistry</i> , 2018, 66, 3302-3307.	5.2	33
60	Comparison of different analytical techniques for the analysis of carotenoids in tamarillo (<i>Solanum</i>)	3.0	42
61	Phenolic profile, antioxidant and cytotoxic properties of polar extracts from leaves and flowers of <i>Isatis tinctoria</i> L. (<i>Brassicaceae</i>) growing in Sicily. <i>Plant Biosystems</i> , 2018, 152, 795-803.	1.6	24
62	Authentication of citrus volatiles based on carbon isotope ratios. <i>Journal of Essential Oil Research</i> , 2018, 30, 1-15.	2.7	21
63	Novel comprehensive multidimensional liquid chromatography approach for elucidation of the microsphere of shikimate-producing <i>Escherichia coli</i> SP1.1/pKD15.071 strain. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 3473-3482.	3.7	8
64	7. Applications of supercritical fluid chromatography in the field of edible lipids. , 2018, , 163-188.		0
65	Comprehensive Two-Dimensional Liquid Chromatography Coupled to Mass Spectrometry. <i>Comprehensive Analytical Chemistry</i> , 2018, 79, 81-123.	1.3	3
66	Supercritical Fluid Chromatography—Ultra-High Pressure Liquid Chromatography for Red Chilli Pepper Fingerprinting by Photodiode Array, Quadrupole-Time-of-Flight and Ion Mobility Mass Spectrometry (SFC—RP-UHPLC-PDA-Q-ToF MS-IMS). <i>Food Analytical Methods</i> , 2018, 11, 3331-3341.	2.6	20
67	Phenolic profile and biological properties of the leaves of <i>Ficus vasta</i> Forssk. (<i>Moraceae</i>) growing in Egypt. <i>BMC Complementary and Alternative Medicine</i> , 2018, 18, 161.	3.7	13
68	Bioactives Screening in Overripe Fruits and Vegetables by Liquid Chromatography Coupled to Photodiode Array and Mass Spectrometry Detection. <i>Food Analytical Methods</i> , 2018, 11, 3053-3070.	2.6	2
69	Development and characterisation of carotenoid-rich microencapsulates from tropical fruit by-products and yellow tamarillo (<i>Solanum betaceum</i> Cav.). <i>Powder Technology</i> , 2018, 339, 702-709.	4.2	18
70	Characterization of Limonoids in Citrus Essential Oils by Means of Supercritical Fluid Chromatography Tandem Mass Spectrometry. <i>Food Analytical Methods</i> , 2018, 11, 3257-3266.	2.6	10
71	Multidimensional gas chromatographic techniques applied to the analysis of lipids from wild-caught and farmed marine species. <i>European Journal of Lipid Science and Technology</i> , 2017, 119, 1600043.	1.5	20
72	Apocarotenoids determination in <i>Capsicum chinense</i> Jacq. cv. Habanero, by supercritical fluid chromatography-triple-quadrupole/mass spectrometry. <i>Food Chemistry</i> , 2017, 231, 316-323.	8.2	48

#	ARTICLE	IF	CITATIONS
73	Chemical Characterization and Biological Activities of Phenolic-Rich Fraction from Cauline Leaves of <i>Isatis tinctoria</i> L. (Brassicaceae) Growing in Sicily, Italy. <i>Chemistry and Biodiversity</i> , 2017, 14, e1700073.	2.1	29
74	Highly informative multiclass profiling of lipids by ultra-high performance liquid chromatography coupled with low resolution (quadrupole) mass spectrometry by using electrospray ionization and atmospheric pressure chemical ionization interfaces. <i>Journal of Chromatography A</i> , 2017, 1509, 69-82.	3.7	18
75	Ionic liquids as stationary phases for fatty acid analysis by gas chromatography. <i>Analyst</i> , 2017, 142, 4601-4612.	3.5	36
76	Direct online extraction and determination by supercritical fluid extraction with chromatography and mass spectrometry of targeted carotenoids from red Habanero peppers (<i>Capsicum chinense</i>)	0.0	0
77	Separation of lipids. , 2017, , 201-243.		4
78	Comprehensive two-dimensional liquid chromatography. , 2017, , 403-415.		2
79	Multidimensional liquid chromatography in food analysis. <i>TrAC - Trends in Analytical Chemistry</i> , 2017, 96, 116-123.	11.4	59
80	Comprehensive Liquid Chromatography and Other Liquid-Based Comprehensive Techniques Coupled to Mass Spectrometry in Food Analysis. <i>Analytical Chemistry</i> , 2017, 89, 414-429.	6.5	46
81	Comprehensive two-dimensional liquid chromatography for polyphenol analysis in foodstuffs. <i>Journal of Separation Science</i> , 2017, 40, 7-24.	2.5	48
82	Detailed Profiling of the Volatile Oxygenated Fraction of Mandarin Essential Oils by Using the Off-Line Combination of High-Performance Liquid Chromatography and Comprehensive Two-Dimensional Gas Chromatography-Mass Spectrometry. <i>Food Analytical Methods</i> , 2017, 10, 1106-1116.	2.6	7
83	Recent Advances in Comprehensive Two-Dimensional Liquid Chromatography for the Analysis of Natural Products. , 2017, , 287-307.		1
84	Green Sample-Preparation Techniques in Comprehensive Two-Dimensional Chromatography. <i>Comprehensive Analytical Chemistry</i> , 2017, 76, 601-623.	1.3	0
85	Analysis of lipid profile in lipid storage myopathy. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016, 1029-1030, 157-168.	2.3	6
86	Enhanced resolution of <i>Mentha piperita</i> volatile fraction using a novel medium-polarity ionic liquid gas chromatography stationary phase. <i>Journal of Separation Science</i> , 2016, 39, 537-544.	2.5	10
87	Rapid isolation, reliable characterization, and water solubility improvement of polymethoxyflavones from cold-pressed mandarin essential oil. <i>Journal of Separation Science</i> , 2016, 39, 2018-2027.	2.5	20
88	Reuse of Dairy Product: Evaluation of the Lipid Profile Evolution During and After Their Shelf-Life. <i>Food Analytical Methods</i> , 2016, 9, 3143-3154.	2.6	11
89	Characterization of the pigment fraction in sweet bell peppers (<i>Capsicum annuum</i>) harvested at green and overripe yellow and red stages by offline multidimensional convergence chromatography/liquid chromatography-mass spectrometry. <i>Journal of Separation Science</i> , 2016, 39, 3281-3291.	2.5	30
90	Comprehensive two-dimensional liquid chromatography-tandem mass spectrometry for the simultaneous determination of wine polyphenols and target contaminants. <i>Journal of Chromatography A</i> , 2016, 1458, 54-62.	3.7	69

#	ARTICLE	IF	CITATIONS
91	Comprehensive two-dimensional gas chromatography-mass spectrometry: Recent evolution and current trends. <i>Mass Spectrometry Reviews</i> , 2016, 35, 524-534.	5.4	100
92	Reliability of the $\delta^{13}C$ ECN42 limit and global method for extra virgin olive oil purity assessment using different analytical approaches. <i>Food Chemistry</i> , 2016, 190, 216-225.	8.2	9
93	Characterisation of the C50 carotenoids produced by strains of the cheese-ripening bacterium <i>Arthrobacter arilaitensis</i> . <i>International Dairy Journal</i> , 2016, 55, 10-16.	3.0	30
94	<i>Helichrysum italicum</i> (Roth) G. Don fil. subsp. <i>italicum</i> oil analysis by gas chromatography $\delta^{13}C$ carbon isotope ratio mass spectrometry (GC-C-IRMS): a rapid method of genotype differentiation?. <i>Journal of Essential Oil Research</i> , 2016, 28, 193-201.	2.7	10
95	Chemical characterisation of old cabbage (<i>Brassica oleracea</i> L. var. <i>acephala</i>) seed oil by liquid chromatography and different spectroscopic detection systems. <i>Natural Product Research</i> , 2016, 30, 1646-1654.	1.8	22
96	Application of Comprehensive Two-Dimensional Liquid Chromatography for Carotenoid Analysis in Red Mamey (<i>Pouteria sapote</i>) Fruit. <i>Food Analytical Methods</i> , 2016, 9, 2335-2341.	2.6	33
97	Antimicrobial activities, toxicity and phenolic composition of <i>Asphodeline anatolica</i> E. Tuzlaci leaf extracts from Turkey. <i>Natural Product Research</i> , 2016, 30, 2620-2623.	1.8	12
98	Bergamot (<i>Citrus bergamia</i> Risso) as a source of nutraceuticals: Limonoids and flavonoids. <i>Journal of Functional Foods</i> , 2016, 20, 10-19.	3.4	62
99	Capsaicinoids and Carotenoids in <i>Capsicum annum</i> L.: Optimization of the Extraction Method, Analytical Characterization, and Evaluation of its Biological Properties. <i>Food Analytical Methods</i> , 2016, 9, 1381-1390.	2.6	22
100	Role of the flavonoid-rich fraction in the antioxidant and cytotoxic activities of <i>Bauhinia forficata</i> Link. (Fabaceae) leaves extract. <i>Natural Product Research</i> , 2016, 30, 1229-1239.	1.8	40
101	Multidimensional preparative liquid chromatography to isolate flavonoids from bergamot juice and evaluation of their anti-inflammatory potential. <i>Journal of Separation Science</i> , 2015, 38, 4196-4203.	2.5	9
102	Analysis of the sesquiterpene fraction of citrus essential oils by using the off-line combination of high performance liquid chromatography and gas chromatography-based methods: a comparative study. <i>Flavour and Fragrance Journal</i> , 2015, 30, 411-422.	2.6	15
103	Non-polar lipids characterization of Quinoa (<i>Chenopodium quinoa</i>) seed by comprehensive two-dimensional gas chromatography with flame ionization/mass spectrometry detection and non-aqueous reversed-phase liquid chromatography with atmospheric pressure chemical ionization mass spectrometry detection. <i>Journal of Separation Science</i> , 2015, 38, 3151-3160.	2.5	17
104	Carbon isotope ratios of selected volatiles in <i>Citrus sinensis</i> and in orange-flavoured food. <i>Journal of the Science of Food and Agriculture</i> , 2015, 95, 2944-2950.	3.5	13
105	Lipidomics. <i>Comprehensive Analytical Chemistry</i> , 2015, 68, 395-439.	1.3	4
106	Determination of the triacylglycerol fraction in fish oil by comprehensive liquid chromatography techniques with the support of gas chromatography and mass spectrometry data. <i>Analytical and Bioanalytical Chemistry</i> , 2015, 407, 5211-5225.	3.7	36
107	Evolution and status of preparative gas chromatography as a green sample-preparation technique. <i>TrAC - Trends in Analytical Chemistry</i> , 2015, 71, 65-73.	11.4	21
108	The penetration of green sample-preparation techniques in comprehensive two-dimensional gas chromatography. <i>TrAC - Trends in Analytical Chemistry</i> , 2015, 71, 74-84.	11.4	25

#	ARTICLE	IF	CITATIONS
109	Reduced time HPLC analyses for fast quality control of citrus essential oils. <i>Journal of Essential Oil Research</i> , 2015, 27, 307-315.	2.7	32
110	Sample preparation techniques coupled to advanced chromatographic methods for marine organisms investigation. <i>Analytica Chimica Acta</i> , 2015, 875, 41-53.	5.4	25
111	Underestimated sources of flavonoids, limonoids and dietary fiber: Availability in orange's by-products. <i>Journal of Functional Foods</i> , 2015, 12, 150-157.	3.4	53
112	Analysis of human plasma lipids by using comprehensive two-dimensional gas chromatography with dual detection and with the support of high-resolution time-of-flight mass spectrometry for structural elucidation. <i>Journal of Separation Science</i> , 2015, 38, 267-275.	2.5	18
113	Determination of the polyphenolic content of a <i>Capsicum annum</i> L. extract by liquid chromatography coupled to photodiode array and mass spectrometry detection and evaluation of its biological activity. <i>Journal of Separation Science</i> , 2015, 38, 171-178.	2.5	54
114	Study of the carotenoid composition in membrillo, guanabana toreta, jobo and mamey fruits. <i>Fruits</i> , 2015, 70, 163-172.	0.4	10
115	Complementary Analytical Liquid Chromatography Methods for the Characterization of Aqueous Phase from Pyrolysis of Lignocellulosic Biomasses. <i>Analytical Chemistry</i> , 2014, 86, 11255-11262.	6.5	51
116	Flow-modulation low-pressure comprehensive two-dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2014, 1372, 236-244.	3.7	44
117	Thorough investigation of the oxygen heterocyclic fraction of lime (<i>Citrus aurantifolia</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10	2.5	13
118	Underestimated sources of flavonoids, limonoids and dietary fibre: Availability in lemon's by-products. <i>Journal of Functional Foods</i> , 2014, 9, 18-26.	3.4	71
119	Determination of new bioflavonoids in bergamot (<i>Citrus bergamia</i>) peel oil by liquid chromatography coupled to tandem ion trap time-of-flight mass spectrometry. <i>Flavour and Fragrance Journal</i> , 2014, 29, 131-136.	2.6	13
120	Characterisation of lipid fraction of marine macroalgae by means of chromatography techniques coupled to mass spectrometry. <i>Food Chemistry</i> , 2014, 145, 932-940.	8.2	55
121	High performance characterization of triacylglycerols in milk and milk-related samples by liquid chromatography and mass spectrometry. <i>Journal of Chromatography A</i> , 2014, 1360, 172-187.	3.7	54
122	Use of greatly-reduced gas flows in flow-modulated comprehensive two-dimensional gas chromatography-mass spectrometry. <i>Journal of Chromatography A</i> , 2014, 1359, 271-276.	3.7	48
123	Rapid Isolation of High Solute Amounts Using an Online Four-Dimensional Preparative System: Normal Phase-Liquid Chromatography Coupled to Methyl Siloxane-Ionic Liquid-Wax Phase Gas Chromatography. <i>Analytical Chemistry</i> , 2014, 86, 4295-4301.	6.5	20
124	Evaluation of carotenoid and capsaicinoid contents in powder of red chili peppers during one year of storage. <i>Food Research International</i> , 2014, 65, 163-170.	6.2	49
125	Continuous vs. segmented second-dimension system gradients for comprehensive two-dimensional liquid chromatography of sugarcane (<i>Saccharum</i> spp.). <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 4315-4324.	3.7	33
126	Profiling and quantifying polar lipids in milk by hydrophilic interaction liquid chromatography coupled with evaporative light-scattering and mass spectrometry detection. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 4617-4626.	3.7	49

#	ARTICLE	IF	CITATIONS
127	Qualitative and quantitative analysis of the unsaponifiable fraction of vegetable oils by using comprehensive 2D GC with dual MS/FID detection. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 4655-4663.	3.7	27
128	Comparison of two different multidimensional liquidâ€“gas chromatography interfaces for determination of mineral oil saturated hydrocarbons in foodstuffs. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 1077-1084.	3.7	24
129	Analysis of the unsaponifiable fraction of lipids belonging to various milk-types by using comprehensive two-dimensional gas chromatography with dual mass spectrometry/flame ionization detection and with the support of high resolution time-of-flight mass spectrometry for structural elucidation. <i>Journal of Chromatography A</i> , 2013, 1313, 194-201.	3.7	35
130	<i>Juniperus oxycedrus</i> L. subsp. <i>oxycedrus</i> and <i>Juniperus oxycedrus</i> L. subsp. <i>macrocarpa</i> (Sibth. & amp;) Tj ETQq0 0 0 rgBT /Overlock 10 TF and antimicrobial activities. <i>Food and Chemical Toxicology</i> , 2013, 58, 22-29.	3.6	49
131	Multiple headspace-solid-phase microextraction: An application to quantification of mushroom volatiles. <i>Analytica Chimica Acta</i> , 2013, 770, 1-6.	5.4	65
132	Rapid collection and identification of a novel component from <i>Clausena lansium</i> Skeels leaves by means of three-dimensional preparative gas chromatography and nuclear magnetic resonance/infrared/mass spectrometric analysis. <i>Analytica Chimica Acta</i> , 2013, 785, 119-125.	5.4	36
133	<i>Betula pendula</i> Roth leaves: gastroprotective effects of an HPLC-fingerprinted methanolic extract. <i>Natural Product Research</i> , 2013, 27, 1569-1575.	1.8	9
134	The off-line combination of high performance liquid chromatography and comprehensive two-dimensional gas chromatographyâ€“mass spectrometry: A powerful approach for highly detailed essential oil analysis. <i>Journal of Chromatography A</i> , 2013, 1305, 276-284.	3.7	38
135	Untargeted and targeted comprehensive two-dimensional GC analysis using a novel unified high-speed triple quadrupole mass spectrometer. <i>Journal of Chromatography A</i> , 2013, 1278, 153-159.	3.7	43
136	Potential of comprehensive chromatography in food analysis. <i>TrAC - Trends in Analytical Chemistry</i> , 2013, 52, 186-205.	11.4	91
137	Characterization of 12 <i>Capsicum</i> varieties by evaluation of their carotenoid profile and pungency determination. <i>Food Chemistry</i> , 2013, 140, 794-802.	8.2	158
138	Analysis of bovine milk caseins on organic monolithic columns: An integrated capillary liquid chromatographyâ€“high resolution mass spectrometry approach for the study of time-dependent casein degradation. <i>Journal of Chromatography A</i> , 2013, 1313, 259-269.	3.7	29
139	Ultra high performance liquid chromatography with ionâ€“trap <sc>TOF</sc>â€“<sc>MS</sc> for the fast characterization of flavonoids in <i>C</i>itrus bergamia</i> juice. <i>Journal of Separation Science</i>, 2013, 36, 3351-3355.</i>	2.5	19
140	Capillary-liquid chromatography (CLC) and nano-LC in food analysis. <i>TrAC - Trends in Analytical Chemistry</i> , 2013, 52, 226-238.	11.4	71
141	Measurement of fundamental chromatography parameters in conventional and split-flow comprehensive two-dimensional gas chromatography-mass spectrometry: A focus on the importance of second-dimension injection efficiency. <i>Journal of Separation Science</i> , 2013, 36, 212-218.	2.5	8
142	Fast gas chromatography combined with a highâ€“speed triple quadrupole mass spectrometer for the analysis of unknown and target citrus essential oil volatiles. <i>Journal of Separation Science</i> , 2013, 36, 511-516.	2.5	11
143	Advances in Food Analysis. <i>Journal of Chromatography A</i> , 2013, 1313, 1.	3.7	3
144	Gas velocity at the point of re-injection: An additional parameter in comprehensive two-dimensional gas chromatography optimization. <i>Journal of Chromatography A</i> , 2013, 1314, 216-223.	3.7	17

#	ARTICLE	IF	CITATIONS
145	Stop-flow comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection for phospholipid analysis. <i>Journal of Chromatography A</i> , 2013, 1278, 46-53.	3.7	69
146	Native carotenoids composition of some tropical fruits. <i>Food Chemistry</i> , 2013, 140, 825-836.	8.2	85
147	Solid-phase microextraction with fast GC combined with a high-speed triple quadrupole mass spectrometer for targeted and untargeted food analysis. <i>Journal of Separation Science</i> , 2013, 36, 2145-2150.	2.5	13
148	Separation of Lipids. , 2013, , 203-248.		4
149	Determination of the carotenoid profile in peach fruits, juice and jam. <i>Fruits</i> , 2013, 68, 39-44.	0.4	21
150	Bioaccessibility of pistachio polyphenols, xanthophylls, and tocopherols during simulated human digestion. <i>Nutrition</i> , 2013, 29, 338-344.	2.4	111
151	Electronic nose and GC-MS analysis of volatile compounds in Tuber magnatum Pico: Evaluation of different storage conditions. <i>Food Chemistry</i> , 2013, 136, 668-674.	8.2	57
152	Evaluation of Gas Chromatography-Combustion-Isotope Ratio Mass Spectrometry (GC-C-IRMS) for the Quality Assessment of Citrus Liqueurs. <i>Journal of Agricultural and Food Chemistry</i> , 2013, 61, 1661-1670.	5.2	30
153	Mechanisms Underlying the Anti-Tumoral Effects of Citrus bergamia Juice. <i>PLoS ONE</i> , 2013, 8, e61484.	2.5	60
154	Determination of Bioactive Compounds in the Juice of Pummelo (Citrus grandis Osbeck). <i>Natural Product Communications</i> , 2013, 8, 1934578X1300800.	0.5	5
155	Sampling and Sample Preparation Techniques for the Determination of the Volatile Components of Milk and Dairy Products. , 2012, , 43-59.		3
156	A new HPLC method developed for the analysis of oxygen heterocyclic compounds in Citrus essential oils. <i>Journal of Essential Oil Research</i> , 2012, 24, 119-129.	2.7	31
157	Characterization of cold-pressed and processed bergamot oils by using GC-FID, GC-MS, GC-C-IRMS, enantio-GC, MDGC, HPLC and HPLC-MS-IT-TOF. <i>Journal of Essential Oil Research</i> , 2012, 24, 93-117.	2.7	32
158	Recent Developments in High-Performance Liquid Chromatography. , 2012, , 1-32.		0
159	Mass spectrometric elucidation of triacylglycerol content of Brevoortia tyrannus (menhaden) oil using non-aqueous reversed-phase liquid chromatography under ultra high pressure conditions. <i>Journal of Chromatography A</i> , 2012, 1259, 227-236.	3.7	34
160	A flow-modulated comprehensive gas chromatography-mass spectrometry method for the analysis of fatty acid profiles in marine and biological samples. <i>Journal of Chromatography A</i> , 2012, 1255, 171-176.	3.7	31
161	Ultra high pressure in the second dimension of a comprehensive two-dimensional liquid chromatographic system for carotenoid separation in red chili peppers. <i>Journal of Chromatography A</i> , 2012, 1255, 244-251.	3.7	63
162	Increasing the Isolated Quantities and Purities of Volatile Compounds by Using a Triple Deans-Switch Multidimensional Preparative Gas Chromatographic System with an Apolar-Wax-Ionic Liquid Stationary-Phase Combination. <i>Analytical Chemistry</i> , 2012, 84, 7092-7098.	6.5	36

#	ARTICLE	IF	CITATIONS
163	Heart-cutting multidimensional gas chromatography: A review of recent evolution, applications, and future prospects. <i>Analytica Chimica Acta</i> , 2012, 716, 66-75.	5.4	90
164	Protective effects of an extract from <i>Citrus bergamia</i> against inflammatory injury in interferon-gamma and histamine exposed human keratinocytes. <i>Life Sciences</i> , 2012, 90, 968-974.	4.3	41
165	Flavors and Odors. , 2012, , 599-663.		0
166	Use of ionic liquids as stationary phases in hyphenated gas chromatography techniques. <i>Journal of Chromatography A</i> , 2012, 1255, 130-144.	3.7	94
167	Determination of Carotenoids and their Esters in Fruits of Sea Buckthorn (<i>Hippophae</i>) Tj ETQq1 1 0.784314 rgBJ /Overlock_10 Tf 50	2.4	48
168	Multidimensional liquid chromatography for the determination of chiral coumarins and furocoumarins in <i>Citrus</i> essential oils. <i>Journal of Separation Science</i> , 2012, 35, 1828-1836.	2.5	29
169	Mass spectrometry detection in comprehensive liquid chromatography: Basic concepts, instrumental aspects, applications and trends. <i>Mass Spectrometry Reviews</i> , 2012, 31, 523-559.	5.4	86
170	Multidimensional enantio gas chromatography/mass spectrometry and gas chromatography-combustion-isotopic ratio mass spectrometry for the authenticity assessment of lime essential oils (<i>C. aurantifolia</i> Swingle and <i>C. latifolia</i> Tanaka). <i>Journal of Chromatography A</i> , 2012, 1226, 87-95.	3.7	26
171	<i>Betula pendula</i> leaves: Polyphenolic characterization and potential innovative use in skin whitening products. <i>Fito-terapia</i> , 2012, 83, 877-882.	2.2	60
172	Authenticity control on lemon essential oils employing Gas Chromatography-Combustion-Isotope Ratio Mass Spectrometry (GC-C-IRMS). <i>Food Chemistry</i> , 2012, 131, 1523-1530.	8.2	29
173	In vitro antimycoplasmal activity of citrus bergamia essential oil and its major components. <i>European Journal of Medicinal Chemistry</i> , 2012, 52, 66-69.	5.5	45
174	Analysis of <i>Citrus</i> essential oils: state of the art and future perspectives. A review.. <i>Flavour and Fragrance Journal</i> , 2012, 27, 98-123.	2.6	91
175	Development of an online capillary comprehensive 2D-LC system for the analysis of proteome samples. <i>Journal of Separation Science</i> , 2012, 35, 530-533.	2.5	22
176	Oxycarotenoids (Xanthophylls). , 2012, , 267-286.		1
177	Evaluation of a Medium-Polarity Ionic Liquid Stationary Phase in the Analysis of Flavor and Fragrance Compounds. <i>Analytical Chemistry</i> , 2011, 83, 7947-7954.	6.5	77
178	Online Comprehensive RPLC-RPLC with Mass Spectrometry Detection for the Analysis of Proteome Samples. <i>Analytical Chemistry</i> , 2011, 83, 2485-2491.	6.5	60
179	Chemical Characterization of Sacha Inchi (<i>Plukenetia volubilis</i> L.) Oil. <i>Journal of Agricultural and Food Chemistry</i> , 2011, 59, 13043-13049.	5.2	111
180	Phenolic composition and biological activities of <i>Juniperus drupacea</i> Labill. berries from Turkey. <i>Food and Chemical Toxicology</i> , 2011, 49, 2600-2608.	3.6	53

#	ARTICLE	IF	CITATIONS
181	Comparison of major lipid components in human and donkey milk: new perspectives for a hypoallergenic diet in humans. <i>Immunopharmacology and Immunotoxicology</i> , 2011, 33, 633-644.	2.4	26
182	Reliable Identification and Quantification of Volatile Components of Sage Essential Oil Using Ultra HPLC. <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600.	0.5	1
183	Analysis of Native Carotenoid Composition of Sweet Bell Peppers by Serially Coupled C30 Columns. <i>Natural Product Communications</i> , 2011, 6, 1934578X1100601.	0.5	4
184	Enantiomeric distribution of key volatile components in Citrus essential oils. <i>Revista Brasileira De Farmacognosia</i> , 2011, 21, 841-849.	1.4	33
185	Modulators for comprehensive two-dimensional gas chromatography. <i>TrAC - Trends in Analytical Chemistry</i> , 2011, 30, 1437-1461.	11.4	115
186	Comprehensive chromatographic separations in proteomics. <i>Journal of Chromatography A</i> , 2011, 1218, 8777-8790.	3.7	39
187	A rapid multidimensional liquid-gas chromatography method for the analysis of mineral oil saturated hydrocarbons in vegetable oils. <i>Journal of Chromatography A</i> , 2011, 1218, 7476-7480.	3.7	42
188	Determination of phospholipids in milk samples by means of hydrophilic interaction liquid chromatography coupled to evaporative light scattering and mass spectrometry detection. <i>Journal of Chromatography A</i> , 2011, 1218, 6476-6482.	3.7	110
189	Headspace-solid phase microextraction coupled to gas chromatography-combustion-isotope ratio mass spectrometer and to enantioselective gas chromatography for strawberry flavoured food quality control. <i>Journal of Chromatography A</i> , 2011, 1218, 7481-7486.	3.7	42
190	Advances in food analysis. <i>Journal of Chromatography A</i> , 2011, 1218, 7385.	3.7	4
191	Identification of the Bacterial Cellular Lipid Fraction by Using Fast GC-MS and Innovative MS Libraries. <i>NATO Science for Peace and Security Series A: Chemistry and Biology</i> , 2011, , 231-244.	0.5	1
192	Analysis of anthocyanins in commercial fruit juices by using nano-liquid chromatography-electrospray-mass spectrometry and high-performance liquid chromatography with UV-vis detector. <i>Journal of Separation Science</i> , 2011, 34, 150-159.	2.5	59
193	Comprehensive two-dimensional liquid chromatography with evaporative light-scattering detection for the analysis of triacylglycerols in <i>Borago officinalis</i> . <i>Journal of Separation Science</i> , 2011, 34, 688-692.	2.5	24
194	Determination of flavanones in <i>Citrus</i> juices by means of one- and two-dimensional liquid chromatography. <i>Journal of Separation Science</i> , 2011, 34, 681-687.	2.5	46
195	Performance evaluation of a rapid-scanning quadrupole mass spectrometer in the comprehensive two-dimensional gas chromatography analysis of pesticides in water. <i>Journal of Separation Science</i> , 2011, 34, 2411-2417.	2.5	35
196	Analytical characterization of mandarin (<i>Citrus deliciosa</i> Ten.) essential oil. <i>Flavour and Fragrance Journal</i> , 2011, 26, 34-46.	2.6	28
197	Employing ultra high pressure liquid chromatography as the second dimension in a comprehensive two-dimensional system for analysis of <i>Stevia rebaudiana</i> extracts. <i>Journal of Chromatography A</i> , 2011, 1218, 2012-2018.	3.7	90
198	Application of a multidimensional gas chromatography system with simultaneous mass spectrometric and flame ionization detection to the analysis of sandalwood oil. <i>Journal of Chromatography A</i> , 2011, 1218, 137-142.	3.7	42

#	ARTICLE	IF	CITATIONS
199	A flexible loop-type flow modulator for comprehensive two-dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 3140-3145.	3.7	35
200	Characterization of Oils from the Fruits, Leaves and Flowers of the Bitter Orange Tree. <i>Journal of Essential Oil Research</i> , 2011, 23, 45-59.	2.7	55
201	Authentication of Bergamot Essential Oil by Gas Chromatography-Combustion-Isotope Ratio Mass Spectrometer (GC-C-IRMS). <i>Journal of Essential Oil Research</i> , 2011, 23, 60-71.	2.7	32
202	Analytical Characterization of Industrial Essential Oils from Fruits and Leaves of <i>C. aurantifolia</i> and <i>C. latifolia</i> Swing. <i>Journal of Essential Oil Research</i> , 2011, 23, 68-79.	2.7	13
203	Advances of Modern Gas Chromatography and Hyphenated Techniques for Analysis of Plant Extracts. <i>Current Organic Chemistry</i> , 2010, 14, 1752-1768.	1.6	8
204	Evaluation of tea tree oil quality and ascaridole: A deep study by means of chiral and multi heart-cuts multidimensional gas chromatography system coupled to mass spectrometry detection. <i>Journal of Chromatography A</i> , 2010, 1217, 6422-6427.	3.7	42
205	Study on the chemical composition variability of some processed bergamot (<i>Citrus bergamia</i>) essential oils. <i>Flavour and Fragrance Journal</i> , 2010, 25, 4-12.	2.6	53
206	Genuineness assessment of mandarin essential oils employing gas chromatography-combustion-isotope ratio MS (GC-C-IRMS). <i>Journal of Separation Science</i> , 2010, 33, 617-625.	2.5	48
207	RP-LC-RP-LC analysis of a tryptic digest using a combination of totally porous and partially porous stationary phases. <i>Journal of Separation Science</i> , 2010, 33, 1454-1461.	2.5	38
208	Characterization of bacterial lipid profiles by using rapid sample preparation and fast comprehensive two-dimensional gas chromatography in combination with mass spectrometry. <i>Journal of Separation Science</i> , 2010, 33, 2334-2340.	2.5	38
209	Accurate quadrupole MS peak reconstruction in optimized gas-flow comprehensive two-dimensional gas chromatography. <i>Journal of Separation Science</i> , 2010, 33, 2791-2795.	2.5	4
210	Sicilian lemon oil: Composition of volatile and oxygen heterocyclic fractions and enantiomeric distribution of volatile components. <i>Journal of Separation Science</i> , 2010, 33, 3374-3385.	2.5	33
211	Characterization of polyphenols, lipids and dietary fibre from almond skins (<i>Amygdalus communis</i> L.). <i>Journal of Food Composition and Analysis</i> , 2010, 23, 166-174.	3.9	131
212	Thorough evaluation of the validity of conventional enantio-gas chromatography in the analysis of volatile chiral compounds in mandarin essential oil: A comparative investigation with multidimensional gas chromatography. <i>Journal of Chromatography A</i> , 2010, 1217, 1101-1105.	3.7	42
213	Optimized use of a 50 μ m ID secondary column in comprehensive two-dimensional gas chromatography-mass spectrometry. <i>Journal of Chromatography A</i> , 2010, 1217, 4160-4166.	3.7	28
214	Volatiles from Steam-distilled Leaves of Some Plant Species from Madagascar and New Zealand and Evaluation of Their Biological Activity. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000501.	0.5	4
215	Evaluation of a Rapid-Scanning Quadrupole Mass Spectrometer in an Apolar Å -Ionic-Liquid Comprehensive Two-Dimensional Gas Chromatography System. <i>Analytical Chemistry</i> , 2010, 82, 8583-8590.	6.5	88
216	Analysis of Fresh and Aged Tea Tree Essential Oils By Using GCxGC-qMS. <i>Journal of Chromatographic Science</i> , 2010, 48, 262-266.	1.4	42

#	ARTICLE	IF	CITATIONS
217	Free carotenoid and carotenoid ester composition in native orange juices of different varieties. <i>Fruits</i> , 2010, 65, 277-284.	0.4	43
218	Characterization of Mandarin (<i>Citrus deliciosa</i> Ten.) Essential Oil. Determination of Volatiles, Non-Volatiles, Physico-Chemical Indices and Enantiomeric Ratios. <i>Natural Product Communications</i> , 2009, 4, 1934578X0900401.	0.5	7
219	An investigation on the volatile composition of some <i>Artemisia</i> species from Iran. <i>Flavour and Fragrance Journal</i> , 2009, 24, 75-82.	2.6	26
220	Essential oil composition of <i>Citrus medica</i> L. Cv. Diamante (Diamante citron) determined after using different extraction methods. <i>Journal of Separation Science</i> , 2009, 32, 99-108.	2.5	27
221	High peak capacity separation of peptides through the serial connection of LC shellâ€packed columns. <i>Journal of Separation Science</i> , 2009, 32, 1129-1136.	2.5	34
222	Epoxy-carotenoids esters analysis in intact orange juices using twoâ€dimensional comprehensive liquid chromatography. <i>Journal of Separation Science</i> , 2009, 32, 973-980.	2.5	49
223	Characterization of the yerba mate (<i>Ilex paraguariensis</i>) volatile fraction using solidâ€phase microextractionâ€comprehensive 2â€ GCâ€MS. <i>Journal of Separation Science</i> , 2009, 32, 3755-3763.	2.5	27
224	Characterization of the polyphenolic fraction of <i>Morus alba</i> leaves extracts by HPLC coupled to a hybrid ITâ€OF MS system. <i>Journal of Separation Science</i> , 2009, 32, 3627-3634.	2.5	56
225	Separation of organophosphorus pesticides by using nano-liquid chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 3970-3976.	3.7	61
226	Comprehensive two-dimensional liquid chromatography to quantify polyphenols in red wines. <i>Journal of Chromatography A</i> , 2009, 1216, 7483-7487.	3.7	74
227	Enhanced resolution comprehensive two-dimensional gas chromatography applied to the analysis of roasted coffee volatiles. <i>Journal of Chromatography A</i> , 2009, 1216, 7301-7306.	3.7	35
228	High efficiency liquid chromatography techniques coupled to mass spectrometry for the characterization of mate extracts. <i>Journal of Chromatography A</i> , 2009, 1216, 7213-7221.	3.7	89
229	Optimized Use of a 50 Î¼m Internal Diameter Secondary Column in a Comprehensive Two-Dimensional Gas Chromatography System. <i>Analytical Chemistry</i> , 2009, 81, 8529-8537.	6.5	17
230	Determination of Oxygen Heterocyclic Components in Citrus Products by HPLC with UV Detection. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 6543-6551.	5.2	57
231	Comparative Analysis of Flavonoid Profile, Antioxidant and Antimicrobial Activity of the Berries of <i>Juniperus communis</i> L. var. <i>communis</i> and <i>Juniperus communis</i> L. var. <i>saxatilis</i> Pall. from Turkey. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 6570-6577.	5.2	91
232	Evaluation of Use of a Dicationic Liquid Stationary Phase in the Fast and Conventional Gas Chromatographic Analysis of Health-Hazardous C ₁₈ Cis/Trans Fatty Acids. <i>Analytical Chemistry</i> , 2009, 81, 5561-5568.	6.5	67
233	Chapter 10 Analysis of Food Constituents. <i>Comprehensive Analytical Chemistry</i> , 2009, , 215-241.	1.3	1
234	Comprehensive twoâ€dimensional gas chromatographyâ€mass spectrometry: A review. <i>Mass Spectrometry Reviews</i> , 2008, 27, 101-124.	5.4	350

#	ARTICLE	IF	CITATIONS
235	Elucidation of fatty acid profiles in vegetable oils exploiting group-type patterning and enhanced sensitivity of comprehensive two-dimensional gas chromatography. <i>Journal of Separation Science</i> , 2008, 31, 1797-1802.	2.5	32
236	Analysis of native carotenoid composition in orange juice using C ₃₀ columns in tandem. <i>Journal of Separation Science</i> , 2008, 31, 2151-2160.	2.5	46
237	Evaluation of use of a very short polar microbore column segment in high-speed gas chromatography analysis. <i>Journal of Separation Science</i> , 2008, 31, 2634-2639.	2.5	17
238	Use of partially porous column as second dimension in comprehensive two-dimensional system for analysis of polyphenolic antioxidants. <i>Journal of Separation Science</i> , 2008, 31, 3297-3308.	2.5	72
239	Acquisition of deeper knowledge on the human plasma fatty acid profile exploiting comprehensive 2D GC. <i>Journal of Separation Science</i> , 2008, 31, 3347-3351.	2.5	35
240	Offline LC-GC-MS in combination with rapid-scanning quadrupole mass spectrometry. <i>Journal of Separation Science</i> , 2008, 31, 3329-3336.	2.5	15
241	GC-MS, GC-O and enantio-GC investigation of the essential oil of <i>Tarhomonanthus camphoratus</i> L. <i>Flavour and Fragrance Journal</i> , 2008, 23, 40-48.	2.6	99
242	Linear retention indices in gas chromatographic analysis: a review. <i>Flavour and Fragrance Journal</i> , 2008, 23, 297-314.	2.6	192
243	Comprehensive multidimensional liquid chromatography: Theory and applications. <i>Journal of Chromatography A</i> , 2008, 1184, 353-368.	3.7	299
244	Gas chromatography-olfactometry in food flavour analysis. <i>Journal of Chromatography A</i> , 2008, 1186, 123-143.	3.7	214
245	Reliable identification of pesticides using linear retention indices as an active tool in gas chromatographic-mass spectrometric analysis. <i>Journal of Chromatography A</i> , 2008, 1186, 430-433.	3.7	15
246	Comprehensive normal-phase-reversed-phase liquid chromatography coupled to photodiode array and mass spectrometry detection for the analysis of free carotenoids and carotenoid esters from mandarin. <i>Journal of Chromatography A</i> , 2008, 1189, 196-206.	3.7	82
247	Serial coupled columns reversed-phase separations in high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2008, 1188, 208-215.	3.7	45
248	Application of Comprehensive Two-Dimensional Liquid Chromatography To Elucidate the Native Carotenoid Composition in Red Orange Essential Oil. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 3478-3485.	5.2	64
249	Quantification in Comprehensive Two-Dimensional Liquid Chromatography. <i>Analytical Chemistry</i> , 2008, 80, 5418-5424.	6.5	53
250	Release of Protein, Lipid, and Vitamin E from Almond Seeds during Digestion. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 3409-3416.	5.2	160
251	The Protective Effect of Bergamot Oil Extract on Lecitine-like OxyLDL Receptor-1 Expression in Balloon Injury-related Neointima Formation. <i>Journal of Cardiovascular Pharmacology and Therapeutics</i> , 2008, 13, 120-129.	2.0	48
252	Multidimensional Liquid Chromatographic Separations Applied to the Analysis of Food Samples. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2008, 31, 1758-1807.	1.0	40

#	ARTICLE	IF	CITATIONS
253	Characterization of <i>Artemisia arborescens</i> L. (Asteraceae) leaf-derived essential oil from Southern Italy. <i>Journal of Essential Oil Research</i> , 2007, 19, 218-224.	2.7	18
254	Hypolipidemic Effects of Citrus bergamia Risso et Poiteau Juice in Rats Fed a Hypercholesterolemic Diet. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 10671-10677.	5.2	78
255	Generation of Improved Gas Linear Velocities in a Comprehensive Two-Dimensional Gas Chromatography System. <i>Analytical Chemistry</i> , 2007, 79, 2266-2275.	6.5	54
256	Reliable Identification of Terpenoids and Related Compounds by using Linear Retention Indices Interactively with Mass Spectrometry Search. <i>Natural Product Communications</i> , 2007, 2, 1934578X0700200.	0.5	7
257	Determination of trans-resveratrol in wine by micro-HPLC with fluorescence detection. <i>Journal of Separation Science</i> , 2007, 30, 669-672.	2.5	16
258	Superheated water as chromatographic eluent for parabens separation on octadecyl coated zirconia stationary phase. <i>Journal of Separation Science</i> , 2007, 30, 1125-1130.	2.5	21
259	Rapid analysis of food products by means of high speed gas chromatography. <i>Journal of Separation Science</i> , 2007, 30, 508-526.	2.5	40
260	Fast gas chromatography-full scan quadrupole mass spectrometry for the determination of allergens in fragrances. <i>Journal of Separation Science</i> , 2007, 30, 1905-1911.	2.5	39
261	Comprehensive chromatographic methods for the analysis of lipids. <i>TrAC - Trends in Analytical Chemistry</i> , 2007, 26, 191-205.	11.4	73
262	A comprehensive study on the chemical composition and aromatic characteristics of lemon liquor. <i>Food Chemistry</i> , 2007, 105, 771-783.	8.2	37
263	Odour fingerprint acquisition by means of comprehensive two-dimensional gas chromatography-olfactometry and comprehensive two-dimensional gas chromatography/mass spectrometry. <i>Journal of Chromatography A</i> , 2007, 1141, 279-286.	3.7	59
264	Antifungal activity of essential oils against filamentous fungi determined by broth microdilution and vapour contact methods. <i>Journal of Applied Microbiology</i> , 2007, 102, 1544-1550.	3.1	155
265	Comprehensive gas chromatography coupled to mass spectrometry for the separation of pesticides in a very complex matrix. <i>Analytical and Bioanalytical Chemistry</i> , 2007, 389, 1755-1763.	3.7	39
266	Elucidation of Carotenoid Patterns in Citrus Products by Means of Comprehensive Normal-Phase and Reversed-Phase Liquid Chromatography. <i>Analytical Chemistry</i> , 2006, 78, 7743-7750.	6.5	107
267	Evaluation of Leaf-Derived Extracts as an Environmentally Sustainable Source of Essential Oils by Using Gas Chromatography-Mass Spectrometry and Enantioselective Gas Chromatography-Olfactometry. <i>Analytical Chemistry</i> , 2006, 78, 883-890.	6.5	29
268	Inhibition of nitric oxide biosynthesis by anthocyanin fraction of blackberry extract. <i>Nitric Oxide - Biology and Chemistry</i> , 2006, 15, 30-39.	2.7	140
269	Fast enantiomeric analysis of a complex essential oil with an innovative multidimensional gas chromatographic system. <i>Journal of Chromatography A</i> , 2006, 1105, 11-16.	3.7	31
270	Comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection in the analyses of triacylglycerols in natural lipidic matrixes. <i>Journal of Chromatography A</i> , 2006, 1112, 269-275.	3.7	135

#	ARTICLE	IF	CITATIONS
271	High-throughput analysis of bergamot essential oil by fast solid-phase microextractionâ€“capillary gas chromatographyâ€“flame ionization detection. <i>Journal of Chromatography A</i> , 2006, 1103, 162-165.	3.7	28
272	Rapid, micro-scale preparation and very fast gas chromatographic separation of cod liver oil fatty acid methyl esters. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2006, 41, 1566-1570.	2.8	67
273	Determination of beef tallow in lard through a multidimensional off-line non-aqueous reversed phaseâ€“argentation LC method coupled to mass spectrometry. <i>Journal of Separation Science</i> , 2006, 29, 567-575.	2.5	33
274	Separation of triacylglycerols in a complex lipidic matrix by using comprehensive two-dimensional liquid chromatography coupled with atmospheric pressure chemical ionization mass spectrometric detection. <i>Journal of Separation Science</i> , 2006, 29, 1146-1154.	2.5	77
275	Optimization of a Comprehensive Two-Dimensional Normal-Phase and Reversed-Phase Liquid Chromatography System. <i>Journal of Chromatographic Science</i> , 2006, 44, 561-565.	1.4	21
276	Silver-ion reversed-phase comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection in lipidic food analysis. <i>Journal of Chromatography A</i> , 2005, 1086, 91-98.	3.7	115
277	Characterization of cold-pressed Mexican dancy tangerine oils. <i>Flavour and Fragrance Journal</i> , 2005, 20, 60-66.	2.6	18
278	Comprehensive two-dimensional GC for the analysis of citrus essential oils. <i>Flavour and Fragrance Journal</i> , 2005, 20, 136-140.	2.6	47
279	Determination of flavor components in Sicilian goat cheese by automated HS-SPME-GC. <i>Flavour and Fragrance Journal</i> , 2005, 20, 659-665.	2.6	46
280	Advanced and innovative chromatographic techniques for the study of citrus essential oils. <i>Flavour and Fragrance Journal</i> , 2005, 20, 249-264.	2.6	24
281	A comparison between different techniques for the isolation of rosemary essential oil. <i>Journal of Separation Science</i> , 2005, 28, 273-280.	2.5	125
282	Determination of triacylglycerols in donkey milk by using high performance liquid chromatography coupled with atmospheric pressure chemical ionization mass spectrometry. <i>Journal of Separation Science</i> , 2005, 28, 1023-1030.	2.5	42
283	Reliable characterization of coffee bean aroma profiles by automated headspace solid phase microextraction-gas chromatography-mass spectrometry with the support of a dual-filter mass spectra library. <i>Journal of Separation Science</i> , 2005, 28, 1101-1109.	2.5	80
284	Determination of flavonoids in citrus juices by micro-HPLC-ESI/MS. <i>Journal of Separation Science</i> , 2005, 28, 1149-1156.	2.5	131
285	Comprehensive two-dimensional gas chromatography in combination with rapid scanning quadrupole mass spectrometry in perfume analysis. <i>Journal of Chromatography A</i> , 2005, 1067, 235-243.	3.7	95
286	Effect of Anthocyanins Contained in a Blackberry Extract on the Circulatory Failure and Multiple Organ Dysfunction Caused by Endotoxin in the Rat. <i>Planta Medica</i> , 2004, 70, 745-752.	1.3	32
287	Fast GC for the Analysis of Citrus Oils. <i>Journal of Chromatographic Science</i> , 2004, 42, 410-416.	1.4	40
288	Evaluation of fast gas chromatography and gas chromatographyâ€“mass spectrometry in the analysis of lipids. <i>Journal of Chromatography A</i> , 2004, 1035, 237-247.	3.7	65

#	ARTICLE	IF	CITATIONS
289	Off-line coupling of non-aqueous reversed-phase and silver ion high-performance liquid chromatography with mass spectrometry for the characterization of rice oil triacylglycerol positional isomers. <i>Journal of Chromatography A</i> , 2004, 1041, 135-142.	3.7	114
290	Comprehensive multidimensional GC for the characterization of roasted coffee beans. <i>Journal of Separation Science</i> , 2004, 27, 442-450.	2.5	76
291	Determination of anthocyanins and related components in red wines by micro- and capillary HPLC. <i>Journal of Separation Science</i> , 2004, 27, 1458-1466.	2.5	17
292	Fast GC analysis with a 50 μ m ID column: theory, practical aspects, and application to a highly complex sample. <i>Journal of Separation Science</i> , 2004, 27, 1149-1156.	2.5	21
293	Comprehensive Two-Dimensional Normal-Phase (Adsorption) Reversed-Phase Liquid Chromatography. <i>Analytical Chemistry</i> , 2004, 76, 2525-2530.	6.5	149
294	Comprehensive two-dimensional chromatography in food analysis. <i>Journal of Chromatography A</i> , 2004, 1054, 3-16.	3.7	91
295	Comprehensive two-dimensional chromatography in food analysis. <i>Journal of Chromatography A</i> , 2004, 1054, 3-16.	3.7	8
296	Fast GC for the analysis of fats and oils. <i>Journal of Separation Science</i> , 2003, 26, 1467-1473.	2.5	29
297	Detailed analysis and group-type separation of natural fats and oils using comprehensive two-dimensional gas chromatography. <i>Journal of Chromatography A</i> , 2003, 1019, 187-196.	3.7	77
298	Characterization of the Anthocyanin Fraction of Sicilian Blood Orange Juice by Micro-HPLC-ESI/MS. <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 1173-1176.	5.2	76
299	Protective effects of cyanidin-3-O-glucoside from blackberry extract against peroxynitrite-induced endothelial dysfunction and vascular failure. <i>Life Sciences</i> , 2003, 73, 1097-1114.	4.3	162
300	Protective Effects of Anthocyanins from Blackberry in a Rat Model of Acute Lung Inflammation. <i>Free Radical Research</i> , 2003, 37, 891-900.	3.3	150
301	Comparison of Fast and Conventional GC Analysis for Citrus Essential Oils. <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 5602-5606.	5.2	50
302	Analysis of oxygen heterocyclic compounds in citrus essential oils by capillary electrochromatography and comparison with HPLC. <i>Chromatographia</i> , 2001, 53, 57-62.	1.3	18
303	Identification of Anthocyanins in Berries by Narrow-Bore High-Performance Liquid Chromatography with Electrospray Ionization Detection. <i>Journal of Agricultural and Food Chemistry</i> , 2001, 49, 3987-3992.	5.2	133
304	Determination of anthocyanins in blood orange juices by HPLC analysis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2000, 23, 191-195.	2.8	57
305	LC-MS for the identification of oxygen heterocyclic compounds in citrus essential oils. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2000, 24, 147-154.	2.8	135
306	High-performance liquid chromatography coupled on-line with high resolution gas chromatography State of the art. <i>Journal of Chromatography A</i> , 1999, 842, 373-390.	3.7	39

#	ARTICLE	IF	CITATIONS
307	Fast HPLC for the Analysis of Oxygen Heterocyclic Compounds of Citrus Essential Oils. Journal of Agricultural and Food Chemistry, 1999, 47, 4237-4239.	5.2	44
308	IDENTIFICATION OF MINOR OXYGEN HETEROCYCLIC COMPOUNDS OF CITRUS ESSENTIAL OILS BY LIQUID CHROMATOGRAPHY-ATMOSPHERIC PRESSURE CHEMICAL IONISATION MASS SPECTROMETRY. Journal of Liquid Chromatography and Related Technologies, 1999, 22, 2991-3005.	1.0	26
309	Rosmarinus officinalisL. (Labiatae) Essential Oils from the South of Brazil and Uruguay. Journal of Essential Oil Research, 1999, 11, 27-30.	2.7	25
310	On the genuineness of citrus essential oils. Part LVII. The composition of distilled lime oil. Flavour and Fragrance Journal, 1998, 13, 93-97.	2.6	14
311	On the genuineness of citrus essential oils. Part LIII. Determination of the composition of the oxygen heterocyclic fraction of lemon essential oils (Citrus limon (L.) Burm. F.) by normal-phase high performance liquid chromatography. Flavour and Fragrance Journal, 1998, 13, 329-334.	2.6	21
312	Multidimensional capillary GC-GC for the analysis of real complex samples. Part II. Enantiomeric distribution of monoterpene hydrocarbons and monoterpene alcohols of cold-pressed and distilled lime oils. Journal of Separation Science, 1998, 10, 203-212.	1.0	40
313	Multidimensional Tandem Capillary Gas Chromatography System for the Analysis of Real Complex Samples. Part I: Development of a Fully Automated Tandem Gas Chromatography System. Journal of Chromatographic Science, 1998, 36, 201-209.	1.4	67
314	Characterization of Cold-Pressed Key and Persian Lime Oils by Gas Chromatography, Gas Chromatography/Mass Spectroscopy, High-Performance Liquid Chromatography, and Physicochemical Indices. Journal of Agricultural and Food Chemistry, 1997, 45, 3608-3616.	5.2	39
315	Italian <i>Citrus</i> Petitgrain Oils. Part IV. Composition of Lemon Petitgrain Oil. Journal of Essential Oil Research, 1997, 9, 495-508.	2.7	14
316	On the Genuineness of Citrus Essential Oils. Part II. Chemical Characterization of the Essential Oil of New Hybrids of Lemon Obtained in Sicily. , 1997, 12, 153-161.		19
317	On the Genuineness of Citrus Essential Oils. Part III. Chemical Characterization of Essential Oil of three Cultivars of <i>Citrus clementine</i> Hort.. , 1997, 12, 163-172.		12
318	Uruguayan essential oils. Part VI. Composition of lemon oil. Flavour and Fragrance Journal, 1997, 12, 247-255.	2.6	15
319	On the Genuineness of Citrus Essential Oils. 51. Oxygen Heterocyclic Compounds of Bitter Orange Oil (<i>Citrus aurantium</i> L.). Journal of Agricultural and Food Chemistry, 1996, 44, 544-549.	5.2	36
320	Rapid Analysis of Polymethoxylated Flavones from Citrus Oils by Supercritical Fluid Chromatography. Journal of Agricultural and Food Chemistry, 1996, 44, 3900-3905.	5.2	24
321	Italian <i>Citrus</i> Petitgrain Oils. Part I. Composition of Bitter Orange Petitgrain Oil. Journal of Essential Oil Research, 1996, 8, 597-609.	2.7	29
322	On-line HPLC-HRGC-MS for the Analysis of Natural Complex Mixtures. Journal of Chromatographic Science, 1996, 34, 174-181.	1.4	18
323	Interactive use of linear retention indices, on polar and apolar columns, with a ms-library for reliable identification of complex mixtures. Journal of Separation Science, 1995, 7, 581-591.	1.0	68
324	Automated HPLC-HRGC: A powerful method for essential oils analysis. Part V. identification of terpene hydrocarbons of bergamot, lemon, mandarin, sweet orange, bitter orange, grapefruit, clementine and mexican lime oils by coupled HPLC-HRGC-MS(ITD). Flavour and Fragrance Journal, 1995, 10, 33-42.	2.6	62

#	ARTICLE	IF	CITATIONS
325	Deterpenation of sweet orange and lemon essential oils with supercritical carbon dioxide using silica gel as an adsorbent. <i>Flavour and Fragrance Journal</i> , 1995, 10, 51-58.	2.6	50
326	Automated LC-GC: A powerful method for essential oils analysis. Part IV. Coupled LC-GC-MS (ITD) for bergamot oil analysis. <i>Journal of Separation Science</i> , 1994, 6, 237-244.	1.0	29
327	Automated HPLC-HRGC: A powerful method for essential oil analysis. Part III. Aliphatic and terpene aldehydes of orange oil. <i>Journal of High Resolution Chromatography</i> , 1994, 17, 312-314.	1.4	21
328	On the genuineness of citrus essential oils. Part XLVI. Polymethoxylated flavones of the non-volatile residue of Italian sweet orange and mandarin essential oils. <i>Flavour and Fragrance Journal</i> , 1994, 9, 105-111.	2.6	18
329	On-line high performance liquid chromatography coupled with high resolution gas chromatography and mass spectrometry (HPLC-HRGC-MS) for the analysis of complex mixtures containing highly volatile compounds. <i>Chromatographia</i> , 1994, 39, 529-538.	1.3	17
330	On the Genuineness of Citrus Essential Oils. Part XXXVI. Detection of Added Reconstituted Lemon Oil in Genuine Cold-Pressed Lemon Essential Oil by High Resolution Gas Chromatography with Chiral Capillary Columns. <i>Journal of Essential Oil Research</i> , 1993, 5, 21-26.	2.7	20
331	On the Genuineness of Citrus Essential Oils. Part XXXV. Detection of Added Reconstituted Mandarin Oil in Genuine Cold-Pressed Mandarin Essential Oil by High Resolution Gas Chromatography with Chiral Capillary Columns. <i>Journal of Essential Oil Research</i> , 1992, 4, 589-594.	2.7	27