

Audrey M Neyrinck

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6910078/publications.pdf>

Version: 2024-02-01

129
papers

25,848
citations

30070

54
h-index

13771

129
g-index

132
all docs

132
docs citations

132
times ranked

25068
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Commentary on: prebiotic effects: metabolic and health benefits. British Journal of Nutrition, 2022, 127, 554-555. | 2.3 | 7 |
| 2 | Physical activity enhances the improvement of body mass index and metabolism by inulin: a multicenter randomized placebo-controlled trial performed in obese individuals. BMC Medicine, 2022, 20, 110. | 5.5 | 21 |
| 3 | Restoring an adequate dietary fiber intake by inulin supplementation: a pilot study showing an impact on gut microbiota and sociability in alcohol use disorder patients. Gut Microbes, 2022, 14, 2007042. | 9.8 | 15 |
| 4 | Microbiota and Metabolite Profiling as Markers of Mood Disorders: A Cross-Sectional Study in Obese Patients. Nutrients, 2022, 14, 147. | 4.1 | 6 |
| 5 | Liver alterations are not improved by inulin supplementation in alcohol use disorder patients during alcohol withdrawal: A pilot randomized, double-blind, placebo-controlled study. EBioMedicine, 2022, 80, 104033. | 6.1 | 7 |
| 6 | Breath volatile metabolome reveals the impact of dietary fibres on the gut microbiota: Proof of concept in healthy volunteers. EBioMedicine, 2022, 80, 104051. | 6.1 | 7 |
| 7 | Chitin-glucan supplementation improved postprandial metabolism and altered gut microbiota in subjects at cardiometabolic risk in a randomized trial. Scientific Reports, 2022, 12, . | 3.3 | 6 |
| 8 | Chitin-Glucan Supplementation Altered Gut Microbiota and Improved Postprandial Metabolism in Subjects at Cardiometabolic Risk. Current Developments in Nutrition, 2022, 6, 331. | 0.3 | 0 |
| 9 | Noninvasive monitoring of fibre fermentation in healthy volunteers by analyzing breath volatile metabolites: lessons from the FiberTAG intervention study. Gut Microbes, 2021, 13, 1-16. | 9.8 | 8 |
| 10 | Improvement of gastrointestinal discomfort and inflammatory status by a synbiotic in middle-aged adults: a double-blind randomized placebo-controlled trial. Scientific Reports, 2021, 11, 2627. | 3.3 | 18 |
| 11 | Multi-compartment metabolomics and metagenomics reveal major hepatic and intestinal disturbances in cancer cachectic mice. Journal of Cachexia, Sarcopenia and Muscle, 2021, 12, 456-475. | 7.3 | 30 |
| 12 | Prebiotic dietary fibre intervention improves fecal markers related to inflammation in obese patients: results from the Food4Gut randomized placebo-controlled trial. European Journal of Nutrition, 2021, 60, 3159-3170. | 3.9 | 46 |
| 13 | Hepatoprotective Effects of Indole, a Gut Microbial Metabolite, in Leptin-Deficient Obese Mice. Journal of Nutrition, 2021, 151, 1507-1516. | 2.9 | 27 |
| 14 | Prebiotic Effect of Berberine and Curcumin Is Associated with the Improvement of Obesity in Mice. Nutrients, 2021, 13, 1436. | 4.1 | 22 |
| 15 | Specific gut microbial, biological, and psychiatric profiling related to binge eating disorders: A cross-sectional study in obese patients. Clinical Nutrition, 2021, 40, 2035-2044. | 5.0 | 30 |
| 16 | Prebiotic effect on mood in obese patients is determined by the initial gut microbiota composition: A randomized, controlled trial. Brain, Behavior, and Immunity, 2021, 94, 289-298. | 4.1 | 35 |
| 17 | Dietary fiber deficiency as a component of malnutrition associated with psychological alterations in alcohol use disorder. Clinical Nutrition, 2021, 40, 2673-2682. | 5.0 | 11 |
| 18 | Biomarkers for assessment of intestinal permeability in clinical practice. American Journal of Physiology - Renal Physiology, 2021, 321, G11-G17. | 3.4 | 65 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 19 | A dynamic association between myosteatosis and liver stiffness: Results from a prospective interventional study in obese patients. <i>JHEP Reports</i> , 2021, 3, 100323. | 4.9 | 24 |
| 20 | Microbiota analysis and transient elastography reveal new extra-hepatic components of liver steatosis and fibrosis in obese patients. <i>Scientific Reports</i> , 2021, 11, 659. | 3.3 | 29 |
| 21 | Inflammation-induced cholestasis in cancer cachexia. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2021, 12, 70-90. | 7.3 | 24 |
| 22 | Nutritional interest of dietary fiber and prebiotics in obesity: Lessons from the MyNewGut consortium. <i>Clinical Nutrition</i> , 2020, 39, 414-424. | 5.0 | 77 |
| 23 | <i>In vitro</i> approach to evaluate the fermentation pattern of inulin-rich food in obese individuals. <i>British Journal of Nutrition</i> , 2020, 123, 472-479. | 2.3 | 3 |
| 24 | Gut Microbiota-Induced Changes in $\hat{1}^2$ -Hydroxybutyrate Metabolism Are Linked to Altered Sociability and Depression in Alcohol Use Disorder. <i>Cell Reports</i> , 2020, 33, 108238. | 6.4 | 87 |
| 25 | Microbiome response to diet: focus on obesity and related diseases. <i>Reviews in Endocrine and Metabolic Disorders</i> , 2020, 21, 369-380. | 5.7 | 28 |
| 26 | Development of a Repertoire and a Food Frequency Questionnaire for Estimating Dietary Fiber Intake Considering Prebiotics: Input from the FiberTAG Project. <i>Nutrients</i> , 2020, 12, 2824. | 4.1 | 8 |
| 27 | Metabolite profiling reveals the interaction of chitin-glucan with the gut microbiota. <i>Gut Microbes</i> , 2020, 12, 1810530. | 9.8 | 31 |
| 28 | Influence of the Mediterranean diet on the production of short-chain fatty acids in women at risk for breast cancer (LIBRE). <i>Proceedings of the Nutrition Society</i> , 2020, 79, . | 1.0 | 2 |
| 29 | Breath volatile compounds and conjugated polyunsaturated fatty acids as metabolic biomarkers reflecting the interaction between chitin-glucan and the gut microbiota.. <i>Proceedings of the Nutrition Society</i> , 2020, 79, . | 1.0 | 0 |
| 30 | Discovery of the gut microbial signature driving the efficacy of prebiotic intervention in obese patients. <i>Gut</i> , 2020, 69, 1975-1987. | 12.1 | 141 |
| 31 | Link between gut microbiota and health outcomes in inulin -treated obese patients: Lessons from the Food4Gut multicenter randomized placebo-controlled trial. <i>Clinical Nutrition</i> , 2020, 39, 3618-3628. | 5.0 | 87 |
| 32 | Milk Polar Lipids in a High-Fat Diet Can Prevent Body Weight Gain: Modulated Abundance of Gut Bacteria in Relation with Fecal Loss of Specific Fatty Acids. <i>Molecular Nutrition and Food Research</i> , 2019, 63, e1801078. | 3.3 | 35 |
| 33 | The Janus Face of Cereals: Wheat-Derived Prebiotics Counteract the Detrimental Effect of Gluten on Metabolic Homeostasis in Mice Fed a High-Fat/High-Sucrose Diet. <i>Molecular Nutrition and Food Research</i> , 2019, 63, e1900632. | 3.3 | 15 |
| 34 | Functional Effects of EPS-Producing Bifidobacterium Administration on Energy Metabolic Alterations of Diet-Induced Obese Mice. <i>Frontiers in Microbiology</i> , 2019, 10, 1809. | 3.5 | 35 |
| 35 | Chitin-glucan and pomegranate polyphenols improve endothelial dysfunction. <i>Scientific Reports</i> , 2019, 9, 14150. | 3.3 | 25 |
| 36 | Effects of a diet based on inulin-rich vegetables on gut health and nutritional behavior in healthy humans. <i>American Journal of Clinical Nutrition</i> , 2019, 109, 1683-1695. | 4.7 | 121 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 37 | Contribution of the gut microbiota to the regulation of host metabolism and energy balance: a focus on the gut-liver axis. <i>Proceedings of the Nutrition Society</i> , 2019, 78, 319-328. | 1.0 | 84 |
| 38 | A Preventive Prebiotic Supplementation Improves the Sweet Taste Perception in Diet-Induced Obese Mice. <i>Nutrients</i> , 2019, 11, 549. | 4.1 | 17 |
| 39 | Microbiota and nonalcoholic fatty liver disease. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2019, 22, 393-400. | 2.5 | 28 |
| 40 | High-fat diet induces depression-like behaviour in mice associated with changes in microbiome, neuropeptide Y, and brain metabolome. <i>Nutritional Neuroscience</i> , 2019, 22, 877-893. | 3.1 | 133 |
| 41 | Efficacy of advanced pace-mapping technology for idiopathic premature ventricular complexes ablation. <i>Journal of Interventional Cardiac Electrophysiology</i> , 2018, 51, 271-277. | 1.3 | 7 |
| 42 | Wheat-derived arabinoxylan oligosaccharides with bifidogenic properties abolishes metabolic disorders induced by western diet in mice. <i>Nutrition and Diabetes</i> , 2018, 8, 15. | 3.2 | 28 |
| 43 | Characterization of fructans and dietary fibre profiles in raw and steamed vegetables. <i>International Journal of Food Sciences and Nutrition</i> , 2018, 69, 682-689. | 2.8 | 33 |
| 44 | Particle size determines the anti-inflammatory effect of wheat bran in a model of fructose over-consumption: Implication of the gut microbiota. <i>Journal of Functional Foods</i> , 2018, 41, 155-162. | 3.4 | 24 |
| 45 | Targeting the gut microbiota with inulin-type fructans: preclinical demonstration of a novel approach in the management of endothelial dysfunction. <i>Gut</i> , 2018, 67, 271-283. | 12.1 | 150 |
| 46 | The DPP-4 inhibitor vildagliptin impacts the gut microbiota and prevents disruption of intestinal homeostasis induced by a Western diet in mice. <i>Diabetologia</i> , 2018, 61, 1838-1848. | 6.3 | 76 |
| 47 | The gut microbiota metabolite indole alleviates liver inflammation in mice. <i>FASEB Journal</i> , 2018, 32, 6681-6693. | 0.5 | 137 |
| 48 | Inulin Improves Postprandial Hypertriglyceridemia by Modulating Gene Expression in the Small Intestine. <i>Nutrients</i> , 2018, 10, 532. | 4.1 | 24 |
| 49 | Towards microbiome-informed dietary recommendations for promoting metabolic and mental health: Opinion papers of the MyNewGut project. <i>Clinical Nutrition</i> , 2018, 37, 2191-2197. | 5.0 | 29 |
| 50 | <i>Klebsiella oxytoca</i> expands in cancer cachexia and acts as a gut pathobiont contributing to intestinal dysfunction. <i>Scientific Reports</i> , 2018, 8, 12321. | 3.3 | 71 |
| 51 | Increased gut permeability in cancer cachexia: mechanisms and clinical relevance. <i>Oncotarget</i> , 2018, 9, 18224-18238. | 1.8 | 90 |
| 52 | The Potential Role of the Dipeptidyl Peptidase-4-Like Activity From the Gut Microbiota on the Host Health. <i>Frontiers in Microbiology</i> , 2018, 9, 1900. | 3.5 | 47 |
| 53 | Implication of trans-11,trans-13 conjugated linoleic acid in the development of hepatic steatosis. <i>PLoS ONE</i> , 2018, 13, e0192447. | 2.5 | 8 |
| 54 | Rhubarb extract prevents hepatic inflammation induced by acute alcohol intake, an effect related to the modulation of the gut microbiota. <i>Molecular Nutrition and Food Research</i> , 2017, 61, 1500899. | 3.3 | 138 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 55 | Ffar2 expression regulates leukaemic cell growth in vivo. <i>British Journal of Cancer</i> , 2017, 117, 1336-1340. | 6.4 | 12 |
| 56 | Fat binding capacity and modulation of the gut microbiota both determine the effect of wheat bran fractions on adiposity. <i>Scientific Reports</i> , 2017, 7, 5621. | 3.3 | 51 |
| 57 | A polyphenolic extract from green tea leaves activates fat browning in high-fat-diet-induced obese mice. <i>Journal of Nutritional Biochemistry</i> , 2017, 49, 15-21. | 4.2 | 64 |
| 58 | Spirulina Protects against Hepatic Inflammation in Aging: An Effect Related to the Modulation of the Gut Microbiota?. <i>Nutrients</i> , 2017, 9, 633. | 4.1 | 49 |
| 59 | Intestinal Sucrase as a Novel Target Contributing to the Regulation of Glycemia by Prebiotics. <i>PLoS ONE</i> , 2016, 11, e0160488. | 2.5 | 27 |
| 60 | Nutritional depletion in $\alpha 3$ PUFA in apoE knock-out mice: A new model of endothelial dysfunction associated with fatty liver disease. <i>Molecular Nutrition and Food Research</i> , 2016, 60, 2198-2207. | 3.3 | 4 |
| 61 | Microbiome and metabolic disorders related to obesity: Which lessons to learn from experimental models?. <i>Trends in Food Science and Technology</i> , 2016, 57, 256-264. | 15.1 | 26 |
| 62 | Synbiotic approach restores intestinal homeostasis and prolongs survival in leukaemic mice with cachexia. <i>ISME Journal</i> , 2016, 10, 1456-1470. | 9.8 | 149 |
| 63 | Lack of anti-inflammatory effect of coenzyme Q10 supplementation in the liver of rodents after lipopolysaccharide challenge. <i>Clinical Nutrition Experimental</i> , 2015, 1, 10-18. | 2.0 | 4 |
| 64 | Gut microorganisms as promising targets for the management of type 2 diabetes. <i>Diabetologia</i> , 2015, 58, 2206-2217. | 6.3 | 220 |
| 65 | Ezetimibe and simvastatin modulate gut microbiota and expression of genes related to cholesterol metabolism. <i>Life Sciences</i> , 2015, 132, 77-84. | 4.3 | 43 |
| 66 | Ability of the gut microbiota to produce PUFA-derived bacterial metabolites: Proof of concept in germ-free versus conventionalized mice. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 1603-1613. | 3.3 | 48 |
| 67 | Inulin-type fructans modulate intestinal Bifidobacterium species populations and decrease fecal short-chain fatty acids in obese women. <i>Clinical Nutrition</i> , 2015, 34, 501-507. | 5.0 | 220 |
| 68 | Non Digestible Oligosaccharides Modulate the Gut Microbiota to Control the Development of Leukemia and Associated Cachexia in Mice. <i>PLoS ONE</i> , 2015, 10, e0131009. | 2.5 | 109 |
| 69 | Gut microbiota controls adipose tissue expansion, gut barrier and glucose metabolism: novel insights into molecular targets and interventions using prebiotics. <i>Beneficial Microbes</i> , 2014, 5, 3-17. | 2.4 | 241 |
| 70 | Gut Microbial Metabolites of Polyunsaturated Fatty Acids Correlate with Specific Fecal Bacteria and Serum Markers of Metabolic Syndrome in Obese Women. <i>Lipids</i> , 2014, 49, 397-402. | 1.7 | 63 |
| 71 | Positive interaction between prebiotics and thiazolidinedione treatment on adiposity in diet-induced obese mice. <i>Obesity</i> , 2014, 22, 1653-1661. | 3.0 | 9 |
| 72 | Intestinal permeability, gut-bacterial dysbiosis, and behavioral markers of alcohol-dependence severity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, E4485-93. | 7.1 | 652 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 73 | Modulation of the Gut Microbiota by Nutrients with Prebiotic and Probiotic Properties. <i>Advances in Nutrition</i> , 2014, 5, 624S-633S. | 6.4 | 92 |
| 74 | Prebiotics supplementation improves the endothelial dysfunction in n-3 PUFA-depleted ApoE ^{-/-} mice. <i>Archives of Public Health</i> , 2014, 72, O5. | 2.4 | 1 |
| 75 | Role of the Lower and Upper Intestine in the Production and Absorption of Gut Microbiota-Derived PUFA Metabolites. <i>PLoS ONE</i> , 2014, 9, e87560. | 2.5 | 67 |
| 76 | Polyphenol-rich extract of pomegranate peel alleviates tissue inflammation and hypercholesterolaemia in high-fat diet-induced obese mice: potential implication of the gut microbiota. <i>British Journal of Nutrition</i> , 2013, 109, 802-809. | 2.3 | 197 |
| 77 | Prebiotic approach alleviates hepatic steatosis: Implication of fatty acid oxidative and cholesterol synthesis pathways. <i>Molecular Nutrition and Food Research</i> , 2013, 57, 347-359. | 3.3 | 90 |
| 78 | Insight into the prebiotic concept: lessons from an exploratory, double blind intervention study with inulin-type fructans in obese women. <i>Gut</i> , 2013, 62, 1112-1121. | 12.1 | 632 |
| 79 | Implication of fermentable carbohydrates targeting the gut microbiota on conjugated linoleic acid production in high-fat-fed mice. <i>British Journal of Nutrition</i> , 2013, 110, 998-1011. | 2.3 | 40 |
| 80 | Gut microbiota and metabolic disorders: how prebiotic can work?. <i>British Journal of Nutrition</i> , 2013, 109, S81-S85. | 2.3 | 148 |
| 81 | Curcuma longa Extract Associated with White Pepper Lessens High Fat Diet-Induced Inflammation in Subcutaneous Adipose Tissue. <i>PLoS ONE</i> , 2013, 8, e81252. | 2.5 | 44 |
| 82 | Wheat-derived arabinoxylan oligosaccharides with prebiotic effect increase satietogenic gut peptides and reduce metabolic endotoxemia in diet-induced obese mice. <i>Nutrition and Diabetes</i> , 2012, 2, e28-e28. | 3.2 | 184 |
| 83 | Restoring Specific Lactobacilli Levels Decreases Inflammation and Muscle Atrophy Markers in an Acute Leukemia Mouse Model. <i>PLoS ONE</i> , 2012, 7, e37971. | 2.5 | 186 |
| 84 | Gut microbiota-derived propionate reduces cancer cell proliferation in the liver. <i>British Journal of Cancer</i> , 2012, 107, 1337-1344. | 6.4 | 238 |
| 85 | Sirtuin inhibition attenuates the production of inflammatory cytokines in lipopolysaccharide-stimulated macrophages. <i>Biochemical and Biophysical Research Communications</i> , 2012, 420, 857-861. | 2.1 | 47 |
| 86 | Role of intestinal permeability and inflammation in the biological and behavioral control of alcohol-dependent subjects. <i>Brain, Behavior, and Immunity</i> , 2012, 26, 911-918. | 4.1 | 237 |
| 87 | The Loss of Metabolic Control on Alcohol Drinking in Heavy Drinking Alcohol-Dependent Subjects. <i>PLoS ONE</i> , 2012, 7, e38682. | 2.5 | 58 |
| 88 | Dietary modulation of clostridial cluster XIVa gut bacteria (<i>Roseburia</i> spp.) by chitin-glucan fiber improves host metabolic alterations induced by high-fat diet in mice. <i>Journal of Nutritional Biochemistry</i> , 2012, 23, 51-59. | 4.2 | 215 |
| 89 | Targeting gut microbiota in obesity: effects of prebiotics and probiotics. <i>Nature Reviews Endocrinology</i> , 2011, 7, 639-646. | 9.6 | 653 |
| 90 | Responses of Gut Microbiota and Glucose and Lipid Metabolism to Prebiotics in Genetic Obese and Diet-Induced Leptin-Resistant Mice. <i>Diabetes</i> , 2011, 60, 2775-2786. | 0.6 | 881 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 91 | Inulin-type fructans with prebiotic properties counteract GPR43 overexpression and PPAR β -related adipogenesis in the white adipose tissue of high-fat diet-fed mice. <i>Journal of Nutritional Biochemistry</i> , 2011, 22, 712-722. | 4.2 | 237 |
| 92 | Modulation of the gut microbiota by nutrients with prebiotic properties: consequences for host health in the context of obesity and metabolic syndrome. <i>Microbial Cell Factories</i> , 2011, 10, S10. | 4.0 | 172 |
| 93 | Involvement of gut microbial fermentation in the metabolic alterations occurring in n-3 polyunsaturated fatty acids-depleted mice. <i>Nutrition and Metabolism</i> , 2011, 8, 44. | 3.0 | 15 |
| 94 | Nicotinamide enhances apoptosis of G(M)-CSF-treated neutrophils and attenuates endotoxin-induced airway inflammation in mice. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2011, 300, L354-L361. | 2.9 | 14 |
| 95 | Prebiotic Effects of Wheat Arabinoxylan Related to the Increase in Bifidobacteria, Roseburia and Bacteroides/Prevotella in Diet-Induced Obese Mice. <i>PLoS ONE</i> , 2011, 6, e20944. | 2.5 | 383 |
| 96 | Hepatic n-3 Polyunsaturated Fatty Acid Depletion Promotes Steatosis and Insulin Resistance in Mice: Genomic Analysis of Cellular Targets. <i>PLoS ONE</i> , 2011, 6, e23365. | 2.5 | 83 |
| 97 | Potential interest of gut microbial changes induced by non-digestible carbohydrates of wheat in the management of obesity and related disorders. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2010, 13, 722-728. | 2.5 | 50 |
| 98 | Changes in Intestinal Bifidobacteria Levels Are Associated with the Inflammatory Response in Magnesium-Deficient Mice. <i>Journal of Nutrition</i> , 2010, 140, 509-514. | 2.9 | 83 |
| 99 | Prebiotic effects: metabolic and health benefits. <i>British Journal of Nutrition</i> , 2010, 104, S1-S63. | 2.3 | 1,745 |
| 100 | Gut microbiota fermentation of prebiotics increases satietogenic and incretin gut peptide production with consequences for appetite sensation and glucose response after a meal. <i>American Journal of Clinical Nutrition</i> , 2009, 90, 1236-1243. | 4.7 | 615 |
| 101 | Changes in gut microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven improvement of gut permeability. <i>Gut</i> , 2009, 58, 1091-1103. | 12.1 | 2,061 |
| 102 | Lipid peroxidation is not a prerequisite for the development of obesity and diabetes in high-fat-fed mice. <i>British Journal of Nutrition</i> , 2009, 102, 462-469. | 2.3 | 27 |
| 103 | Coenzyme Q10 supplementation lowers hepatic oxidative stress and inflammation associated with diet-induced obesity in mice. <i>Biochemical Pharmacology</i> , 2009, 78, 1391-1400. | 4.4 | 145 |
| 104 | Assessment of liver phagocytic activity using EPR spectrometry and imaging. <i>Magnetic Resonance Imaging</i> , 2009, 27, 565-569. | 1.8 | 6 |
| 105 | Critical role of Kupffer cells in the management of diet-induced diabetes and obesity. <i>Biochemical and Biophysical Research Communications</i> , 2009, 385, 351-356. | 2.1 | 91 |
| 106 | Dietary supplementation with chitosan derived from mushrooms changes adipocytokine profile in diet-induced obese mice, a phenomenon linked to its lipid-lowering action. <i>International Immunopharmacology</i> , 2009, 9, 767-773. | 3.8 | 78 |
| 107 | Hepatic steatosis in n-3 fatty acid depleted mice: focus on metabolic alterations related to tissue fatty acid composition. <i>BMC Physiology</i> , 2008, 8, 21. | 3.6 | 42 |
| 108 | Physiological effects of dietary fructans extracted from <i>Agave tequilana</i> and <i>Dasyilirion</i> spp.. <i>British Journal of Nutrition</i> , 2008, 99, 254-261. | 2.3 | 119 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Response to "Comment on: Dietary supplementation with laminarin, a fermentable marine Î² (1-3) glucan, protects against hepatotoxicity induced by LPS in rat by modulating immune response in the hepatic tissue". <i>International Immunopharmacology</i> , 2008, 8, 516-517. | 3.8 | 2 |
| 110 | Immunomodulatory properties of two wheat bran fractions " aleurone-enriched and crude fractions " in obese mice fed a high fat diet. <i>International Immunopharmacology</i> , 2008, 8, 1423-1432. | 3.8 | 27 |
| 111 | Changes in Gut Microbiota Control Metabolic Endotoxemia-Induced Inflammation in High-Fat Diet-Induced Obesity and Diabetes in Mice. <i>Diabetes</i> , 2008, 57, 1470-1481. | 0.6 | 3,897 |
| 112 | Modulation of Glucagon-like Peptide 1 and Energy Metabolism by Inulin and Oligofructose: Experimental Data. <i>Journal of Nutrition</i> , 2007, 137, 2547S-2551S. | 2.9 | 163 |
| 113 | Metabolic Endotoxemia Initiates Obesity and Insulin Resistance. <i>Diabetes</i> , 2007, 56, 1761-1772. | 0.6 | 4,964 |
| 114 | Dietary supplementation with laminarin, a fermentable marine Î² (1-3) glucan, protects against hepatotoxicity induced by LPS in rat by modulating immune response in the hepatic tissue. <i>International Immunopharmacology</i> , 2007, 7, 1497-1506. | 3.8 | 94 |
| 115 | Selective increases of bifidobacteria in gut microflora improve high-fat-diet-induced diabetes in mice through a mechanism associated with endotoxaemia. <i>Diabetologia</i> , 2007, 50, 2374-2383. | 6.3 | 1,507 |
| 116 | Role of apoptotic signaling pathway in metabolic disturbances occurring in liver tissue after cryopreservation: Study on rat precision-cut liver slices. <i>Life Sciences</i> , 2006, 78, 1570-1577. | 4.3 | 17 |
| 117 | Effect on Components of the Intestinal Microflora and Plasma Neuropeptide Levels of Feeding <i>Lactobacillus delbrueckii</i> , <i>Bifidobacterium lactis</i> , and Inulin to Adult and Elderly Rats. <i>Applied and Environmental Microbiology</i> , 2006, 72, 6533-6538. | 3.1 | 55 |
| 118 | Impact of inulin and oligofructose on gastrointestinal peptides. <i>British Journal of Nutrition</i> , 2005, 93, S157-S161. | 2.3 | 248 |
| 119 | Oligofructose Promotes Satiety in Rats Fed a High-Fat Diet: Involvement of Glucagon-Like Peptide-1. <i>Obesity</i> , 2005, 13, 1000-1007. | 4.0 | 326 |
| 120 | Insight into the involvement of Kupffer cell-derived mediators in the hepatoprotective effect of glycine upon inflammation: study on rat precision-cut liver slices. <i>Inflammation Research</i> , 2005, 54, 106-112. | 4.0 | 12 |
| 121 | Kupffer Cell Activity Is Involved in the Hepatoprotective Effect of Dietary Oligofructose in Rats with Endotoxic Shock. <i>Journal of Nutrition</i> , 2004, 134, 1124-1129. | 2.9 | 24 |
| 122 | Potential modulation of plasma ghrelin and glucagon-like peptide-1 by anorexigenic cannabinoid compounds, SR141716A (rimonabant) and oleylethanolamide. <i>British Journal of Nutrition</i> , 2004, 92, 757-761. | 2.3 | 154 |
| 123 | Precision-cut liver slices in culture as a tool to assess the physiological involvement of Kupffer cells in hepatic metabolism. <i>Comparative Hepatology</i> , 2004, 3, S45. | 0.9 | 16 |
| 124 | Kupffer cell-derived prostaglandin E2 is involved in regulation of lipid synthesis in rat liver tissue. <i>Cell Biochemistry and Function</i> , 2004, 22, 327-332. | 2.9 | 15 |
| 125 | Prebiotics: actual and potential effects in inflammatory and malignant colonic diseases. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2003, 6, 581-586. | 2.5 | 25 |
| 126 | Inhibition of Kupffer cell activity induces hepatic triglyceride synthesis in fasted rats, independent of lipopolysaccharide challenge. <i>Journal of Hepatology</i> , 2002, 36, 466-473. | 3.7 | 23 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Are Kupffer cells involved in the metabolic adaptation of the liver to dietary carbohydrates given after fasting?. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2000, 1475, 238-244. | 2.4 | 7 |
| 128 | Modulation of paracetamol metabolism by Kupffer cells: A study on rat liver slices. <i>Life Sciences</i> , 1999, 65, 2851-2859. | 4.3 | 29 |
| 129 | Prebiotics and Lipid Metabolism. , 0, , 183-192. | | 7 |