
H Craig Morton

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6912630/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Performance of triploid Atlantic cod (Gadus morhua L.) in commercial aquaculture. Aquaculture, 2016, 464, 699-709.	3.5	14
2	The frequency of spontaneous triploidy in farmed Atlantic salmon produced in Norway during the period 2007–2014. BMC Genetics, 2015, 16, 37.	2.7	35
3	Risk assessment of the environmental impact of Norwegian Atlantic salmon farming. ICES Journal of Marine Science, 2015, 72, 997-1021.	2.5	299
4	Low-dose exposure to alkylphenols adversely affects the sexual development of Atlantic cod (Gadus) Tj ETQq0 0 female cod. Aquatic Toxicology, 2011, 105, 136-150.	0 rgBT /O 4.0	verlock 10 Tf 33
5	Development of Atlantic cod (Gadus morhua) exposed to produced water during early life stages: Effects on embryos, larvae, and juvenile fish. Marine Environmental Research, 2010, 70, 383-394.	2.5	62
6	Ontogeny of lymphoid organs and development of IgM-bearing cells in Atlantic halibut (Hippoglossus) Tj ETQqO	0 0 rgBT /(Overlock 10 1
7	Effects of alkylphenols on the reproductive system of Atlantic cod (Gadus morhua). Aquatic Toxicology, 2007, 81, 207-218.	4.0	84
8	Oral and systemic administration of \hat{l}^2 -glucan protects against lipopolysaccharide-induced shock and organ injury in rats. Clinical and Experimental Immunology, 2007, 148, 168-177.	2.6	78
9	Fc Receptors for IgA. , 2007, , 90-110.		1
10	Cloning and characterization of equine CD89 and identification of the CD89 gene in chimpanzees and rhesus macaques. Immunology, 2005, 115, 74-84.	4.4	18
11	IgA Fc receptors in cattle and horses. Veterinary Immunology and Immunopathology, 2005, 108, 139-143.	1.2	7
12	Characterization of the Ligand Binding Site of the Bovine IgA Fc Receptor (bFcαR). Journal of Biological Chemistry, 2004, 279, 54018-54022.	3.4	5
13	Cloning and characterization of an immunoglobulin A Fc receptor from cattle. Immunology, 2004, 111, 204-211.	4.4	25
14	Monocyte-like and mature macrophages produce CXCL13 (B cell–attracting chemokine 1) in inflammatory lesions with lymphoid neogenesis. Blood, 2004, 104, 3021-3027.	1.4	219
15	Characteristics of Mucosal B Cells with Emphasis on the Human Secretory Immune System. , 2004, , 223-246.		0
16	Expression and Modulation of the Human Immunoglobulin A Fc Receptor (CD89) and the FcR gamma Chain on Myeloid Cells in Blood and Tissue. Scandinavian Journal of Immunology, 2003, 57, 506-516.	2.7	52
17	Cloning and sequencing of a cDNA encoding the bovine FcR Î ³ chain. Veterinary Immunology and Immunopathology, 2001, 82, 101-106.	1.2	2
18	ldentification of Residues within the Extracellular Domain 1 of Bovine Fcγ2R Essential for Binding Bovine IgG2. Journal of Biological Chemistry, 2001, 276, 47794-47800.	3.4	8

H CRAIG MORTON

#	Article	IF	CITATIONS
19	M cell pockets of human Peyer's patches are specialized extensions of germinal centers. European Journal of Immunology, 2001, 31, 107-117.	2.9	60
20	Regulation of switching and production of IgA in human B cells in donors with duplicated $\hat{l}\pm 1$ genes. European Journal of Immunology, 2001, 31, 3622-3630.	2.9	19
21	From B to A the mucosal way. Nature Immunology, 2001, 2, 1093-1094.	14.5	78
22	Activity of Human IgG and IgA Subclasses in Immune Defense Against <i>Neisseria meningitidis</i> Serogroup B. Journal of Immunology, 2001, 166, 6250-6256.	0.8	125
23	Immunoglobulin A cell distribution in the human small intestine: phenotypic and functional characteristics. Immunology, 2000, 101, 354-363.	4.4	67
24	Human Immunoglobulin A Receptor (FcRI, CD89) Function in Transgenic Mice Requires Both FcR γ Chain and CR3 (CD11b/CD18). Blood, 1999, 93, 4387-4394.	1.4	126
25	Immunoglobulin-binding Sites of Human FcαRI (CD89) and Bovine Fcγ2R Are Located in their Membrane-distal Extracellular Domains. Journal of Experimental Medicine, 1999, 189, 1715-1722.	8.5	62
26	The B ell system of human mucosae and exocrine glands. Immunological Reviews, 1999, 171, 45-87.	6.0	268
27	Characterization of the human myeloid IgA Fc receptor I (CD89) gene in a cosmid clone. Immunogenetics, 1999, 49, 586-589.	2.4	8
28	Human Immunoglobulin A Receptor (FcRI, CD89) Function in Transgenic Mice Requires Both FcR γ Chain and CR3 (CD11b/CD18). Blood, 1999, 93, 4387-4394.	1.4	14
29	Alternatively spliced forms of the human myeloid Fcα receptor (CD89) in neutrophils. Immunogenetics, 1996, 43, 246-247.	2.4	15
30	Alternatively spliced forms of the human myeloid Fc&; receptor (CD89) in neutrophils. Immunogenetics, 1996, 43, 246-247.	2.4	39
31	Functional Association between the Human Myeloid Immunoglobulin A Fc Receptor (CD89) and FcR γChain. Journal of Biological Chemistry, 1995, 270, 29781-29787.	3.4	181