
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6916284/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	THE CONCISE GUIDE TO PHARMACOLOGY 2019/20: G proteinâ€coupled receptors. British Journal of Pharmacology, 2019, 176, S21-S141.	5.4	519
2	Genetic programs expressed in resting and IL-4 alternatively activated mouse and human macrophages: similarities and differences. Blood, 2013, 121, e57-e69.	1.4	426
3	International Union of Basic and Clinical Pharmacology. XCIV. Adhesion G Protein–Coupled Receptors. Pharmacological Reviews, 2015, 67, 338-367.	16.0	392
4	The seven-span transmembrane receptor CD97 has a cellular ligand (CD55, DAF) Journal of Experimental Medicine, 1996, 184, 1185-1189.	8.5	353
5	THE CONCISE GUIDE TO PHARMACOLOGY 2021/22: G protein oupled receptors. British Journal of Pharmacology, 2021, 178, S27-S156.	5.4	337
6	Systematic validation of specific phenotypic markers for in vitro polarized human macrophages. Journal of Immunological Methods, 2012, 375, 196-206.	1.4	324
7	THE CONCISE GUIDE TO PHARMACOLOGY 2017/18: Overview. British Journal of Pharmacology, 2017, 174, S1-S16.	5.4	269
8	Afucosylated IgG characterizes enveloped viral responses and correlates with COVID-19 severity. Science, 2021, 371, .	12.6	244
9	Tissue-resident memory T cells populate the human brain. Nature Communications, 2018, 9, 4593.	12.8	242
10	Sticky Signaling—Adhesion Class G Protein–Coupled Receptors Take the Stage. Science Signaling, 2013, 6, re3.	3.6	226
11	Serine proteases of the human immune system in health and disease. Molecular Immunology, 2010, 47, 1943-1955.	2.2	214
12	Transcriptional profiling of human microglia reveals grey–white matter heterogeneity and multiple sclerosis-associated changes. Nature Communications, 2019, 10, 1139.	12.8	214
13	The epidermal growth factor–like domains of the human EMR2 receptor mediate cell attachment through chondroitin sulfate glycosaminoglycans. Blood, 2003, 102, 2916-2924.	1.4	207
14	Staining of HLA-DR, Iba1 and CD68 in human microglia reveals partially overlapping expression depending on cellular morphology and pathology. Journal of Neuroimmunology, 2017, 309, 12-22.	2.3	189
15	CD27 Defines Phenotypically and Functionally Different Human NK Cell Subsets. Journal of Immunology, 2008, 180, 3739-3745.	0.8	173
16	The Epidermal Growth Factor-Seven Transmembrane (EGF-TM7) Receptor CD97 Is Required for Neutrophil Migration and Host Defense. Journal of Immunology, 2004, 172, 1125-1131.	0.8	136
17	F4/80 and the related adhesionâ€GPCRs. European Journal of Immunology, 2011, 41, 2472-2476.	2.9	132
18	Tissue-resident memory T cells invade the brain parenchyma in multiple sclerosis white matter lesions. Brain, 2020, 143, 1714-1730.	7.6	131

#	Article	IF	CITATIONS
19	Expression of the activation antigen CD97 and its ligand CD55 in rheumatoid synovial tissue. Arthritis and Rheumatism, 1999, 42, 650-658.	6.7	125
20	Phenotyping primary human microglia: Tight regulation of LPS responsiveness. Clia, 2012, 60, 1506-1517.	4.9	122
21	Expression and Regulation of CD97 in Colorectal Carcinoma Cell Lines and Tumor Tissues. American Journal of Pathology, 2002, 161, 1657-1667.	3.8	121
22	The EGF-TM7 family: a postgenomic view. Immunogenetics, 2004, 55, 655-666.	2.4	117
23	EMR1, the human homolog of F4/80, is an eosinophilâ€specific receptor. European Journal of Immunology, 2007, 37, 2797-2802.	2.9	113
24	Characterization of the CD55 (DAF)-binding site on the seven-span transmembrane receptor CD97. European Journal of Immunology, 1998, 28, 1701-1707.	2.9	111
25	GITR Triggering Induces Expansion of Both Effector and Regulatory CD4+ T Cells In Vivo. Journal of Immunology, 2009, 182, 7490-7500.	0.8	110
26	Specific expression of GPR56 by human cytotoxic lymphocytes. Journal of Leukocyte Biology, 2011, 90, 735-740.	3.3	104
27	Ligation of the adhesionâ€GPCR EMR2 regulates human neutrophil function. FASEB Journal, 2008, 22, 741-751.	0.5	101
28	Expression of the Inhibitory CD200 Receptor Is Associated with Alternative Macrophage Activation. Journal of Innate Immunity, 2010, 2, 195-200.	3.8	99
29	Human EMR2, a Novel EGF-TM7 Molecule on Chromosome 19p13.1, Is Closely Related to CD97. Genomics, 2000, 67, 188-200.	2.9	98
30	Gene Expression Profiling of Multiple Sclerosis Pathology Identifies Early Patterns of Demyelination Surrounding Chronic Active Lesions. Frontiers in Immunology, 2017, 8, 1810.	4.8	96
31	Blimpâ€1 homolog Hobit identifies effectorâ€type lymphocytes in humans. European Journal of Immunology, 2015, 45, 2945-2958.	2.9	94
32	Autoantibodies against type I interferons are associated with multi-organ failure in COVID-19 patients. Intensive Care Medicine, 2021, 47, 704-706.	8.2	93
33	Differential expression of HIV-1 interfering factors in monocyte-derived macrophages stimulated with polarizing cytokines or interferons. Scientific Reports, 2012, 2, 763.	3.3	85
34	CD97, but Not Its Closely Related EGF-TM7 Family Member EMR2, Is Expressed on Gastric, Pancreatic, and Esophageal Carcinomas. American Journal of Clinical Pathology, 2002, 118, 699-707.	0.7	84
35	The Adhesion G Protein-Coupled Receptor GPR56/ADGRG1 Is an Inhibitory Receptor on Human NK Cells. Cell Reports, 2016, 15, 1757-1770.	6.4	84
36	Characteristics of differentiated CD8+ and CD4+ T cells present in the human brain. Acta Neuropathologica, 2013, 126, 525-535.	7.7	80

#	Article	IF	CITATIONS
37	Isolation of primary microglia from the human post-mortem brain: effects of ante- and post-mortem variables. Acta Neuropathologica Communications, 2017, 5, 16.	5.2	79
38	Clinical features and prognostic factors in Covid-19: A prospective cohort study. EBioMedicine, 2021, 67, 103378.	6.1	79
39	Chronic Exposure to Glucocorticoids Shapes Gene Expression and Modulates Innate and Adaptive Activation Pathways in Macrophages with Distinct Changes in Leukocyte Attraction. Journal of Immunology, 2014, 192, 1196-1208.	0.8	78
40	Expression of the largest CD97 and EMR2 isoforms on leukocytes facilitates a specific interaction with chondroitin sulfate on B cells. Journal of Leukocyte Biology, 2005, 77, 112-119.	3.3	77
41	Tissue distribution of the human CD97 EGF-TM7 receptor. Tissue Antigens, 2001, 57, 325-331.	1.0	75
42	Selective Upregulation of Scavenger Receptors in and Around Demyelinating Areas in Multiple Sclerosis. Journal of Neuropathology and Experimental Neurology, 2013, 72, 106-118.	1.7	75
43	Mouse Hobit is a homolog of the transcriptional repressor Blimp-1 that regulates NKT cell effector differentiation. Nature Immunology, 2012, 13, 864-871.	14.5	71
44	Analysis of CD97 Expression and Manipulation: Antibody Treatment but Not Gene Targeting Curtails Granulocyte Migration. Journal of Immunology, 2008, 181, 6574-6583.	0.8	70
45	Proteolytic cleavage of the EMR2 receptor requires both the extracellular stalk and the GPS motif. FEBS Letters, 2003, 547, 145-150.	2.8	67
46	Shear Stress–Dependent Downregulation of the Adhesion-G Protein–Coupled Receptor CD97 on Circulating Leukocytes upon Contact with Its Ligand CD55. Journal of Immunology, 2013, 190, 3740-3748.	0.8	67
47	HPA axis activity in multiple sclerosis correlates with disease severity, lesion type and gene expression in normal-appearing white matter. Acta Neuropathologica, 2013, 126, 237-249.	7.7	66
48	Structure of the gene coding for the human early lymphocyte activation antigen CD69: A C-type lectin receptor evolutionarily related with the gene families of natural killer cell-specific receptors. European Journal of Immunology, 1994, 24, 1692-1697.	2.9	62
49	Molecular cloning and characterization of mouse CD97. International Immunology, 2000, 12, 439-448.	4.0	60
50	Adhesion GPCRs in Regulating Immune Responses and Inflammation. Advances in Immunology, 2017, 136, 163-201.	2.2	59
51	Oxidative stress and macrophages: driving forces behind exacerbations of asthma and chronic obstructive pulmonary disease?. American Journal of Physiology - Lung Cellular and Molecular Physiology, 2019, 316, L369-L384.	2.9	55
52	AICL: a new activation-induced antigen encoded by the human NK gene complex. Immunogenetics, 1997, 45, 295-300.	2.4	51
53	Identification of the epidermal growth factor-TM7 receptor EMR2 and its ligand dermatan sulfate in rheumatoid synovial tissue. Arthritis and Rheumatism, 2005, 52, 442-450.	6.7	50
54	Expression of the EGF-TM7 receptor CD97 and its ligand CD55 (DAF) in multiple sclerosis. Journal of Neuroimmunology, 2002, 132, 156-163.	2.3	49

#	Article	IF	CITATIONS
55	CD312, the human adhesion-GPCR EMR2, is differentially expressed during differentiation, maturation, and activation of myeloid cells. Biochemical and Biophysical Research Communications, 2007, 353, 133-138.	2.1	49
56	Adhesion G Protein–Coupled Receptors: From In Vitro Pharmacology to In Vivo Mechanisms. Molecular Pharmacology, 2015, 88, 617-623.	2.3	48
57	The human EGF-TM7 receptor EMR3 is a marker for mature granulocytes. Journal of Leukocyte Biology, 2007, 81, 440-448.	3.3	47
58	Deletion of either CD55 or CD97 ameliorates arthritis in mouse models. Arthritis and Rheumatism, 2010, 62, 1036-1042.	6.7	47
59	Microglia in normal appearing white matter of multiple sclerosis are alerted but immunosuppressed. Glia, 2013, 61, 1848-1861.	4.9	46
60	The prolactin receptor is expressed in macrophages within human carotid atherosclerotic plaques: a role for prolactin in atherogenesis?. Journal of Endocrinology, 2011, 208, 107-117.	2.6	45
61	CD69 Antigen of Human Lymphocytes Is a Calcium-Dependent Carbohydrate-Binding Protein. Biochemical and Biophysical Research Communications, 1995, 208, 68-74.	2.1	44
62	Inactivation of the EGF-TM7 receptor EMR4 after the Pan-Homo divergence. European Journal of Immunology, 2003, 33, 1365-1371.	2.9	44
63	Tie2 Signaling Cooperates with TNF to Promote the Pro-Inflammatory Activation of Human Macrophages Independently of Macrophage Functional Phenotype. PLoS ONE, 2014, 9, e82088.	2.5	44
64	CD97 neutralisation increases resistance to collagen-induced arthritis in mice. Arthritis Research and Therapy, 2006, 8, R155.	3.5	43
65	Stromal cell markers are differentially expressed in the synovial tissue of patients with early arthritis. PLoS ONE, 2017, 12, e0182751.	2.5	43
66	Adhesion GPCRs as Modulators of Immune Cell Function. Handbook of Experimental Pharmacology, 2016, 234, 329-350.	1.8	42
67	The human EGF-TM7 family member EMR2 is a heterodimeric receptor expressed on myeloid cells. Journal of Leukocyte Biology, 2002, 71, 854-62.	3.3	42
68	An unusual mode of concerted evolution of the EGF‶M7 receptor chimera EMR2. FASEB Journal, 2006, 20, 2582-2584.	0.5	41
69	Leukocyte adhesion-GPCR EMR2 is aberrantly expressed in human breast carcinomas and is associated with patient survival. Oncology Reports, 2011, 25, 619-27.	2.6	41
70	Angiopoietin-2 promotes inflammatory activation of human macrophages and is essential for murine experimental arthritis. Annals of the Rheumatic Diseases, 2012, 71, 1402-1417.	0.9	41
71	N-glycosylation of CD97 within the EGF domains is crucial for epitope accessibility in normal and malignant cells as well as CD55 ligand binding. International Journal of Cancer, 2004, 112, 815-822.	5.1	39
72	Enhanced uptake of multiple sclerosis-derived myelin by THP-1 macrophages and primary human microglia. Journal of Neuroinflammation, 2014, 11, 64.	7.2	36

#	Article	IF	CITATIONS
73	Single-cell mass cytometry reveals complex myeloid cell composition in active lesions of progressive multiple sclerosis. Acta Neuropathologica Communications, 2020, 8, 136.	5.2	35
74	Overexpression of CD97 in Intestinal Epithelial Cells of Transgenic Mice Attenuates Colitis by Strengthening Adherens Junctions. PLoS ONE, 2010, 5, e8507.	2.5	35
75	Structure of the Human CD97 Gene: Exon Shuffling Has Generated a New Type of Seven-Span Transmembrane Molecule Related to the Secretin Receptor Superfamily. Genomics, 1996, 32, 144-147.	2.9	34
76	Therapeutic Antibody Targeting of CD97 in Experimental Arthritis: the Role of Antigen Expression, Shedding, and Internalization on the Pharmacokinetics of Anti-CD97 Monoclonal Antibody 1B2. Journal of Immunology, 2009, 183, 4127-4134.	0.8	34
77	Macrophages and HIV-1. Current Opinion in HIV and AIDS, 2011, 6, 385-390.	3.8	30
78	Triggering of the dsRNA Sensors TLR3, MDA5, and RIG-I Induces CD55 Expression in Synovial Fibroblasts. PLoS ONE, 2012, 7, e35606.	2.5	29
79	The Adhesion G Protein-Coupled Receptor GPR97/ADGRG3 Is Expressed in Human Granulocytes and Triggers Antimicrobial Effector Functions. Frontiers in Immunology, 2018, 9, 2830.	4.8	27
80	CD97 is a critical regulator of acute myeloid leukemia stem cell function. Journal of Experimental Medicine, 2019, 216, 2362-2377.	8.5	24
81	Regional sublocalization of the human CD69 gene to chromosome bands 12p12.3-p13.2, the predicted region of the human natural killer cell gene complex. European Journal of Immunology, 1993, 23, 2711-2713.	2.9	23
82	Continuous CD27 triggering <i>in vivo</i> strongly reduces NK cell numbers. European Journal of Immunology, 2010, 40, 1107-1117.	2.9	23
83	Do eosinophils contribute to oxidative stress in mild asthma?. Clinical and Experimental Allergy, 2019, 49, 929-931.	2.9	23
84	Differential role of CD97 in interleukin-8-induced and granulocyte-colony stimulating factor-induced hematopoietic stem and progenitor cell mobilization. Haematologica, 2008, 93, 601-604.	3.5	22
85	Postâ€mortem multiple sclerosis lesion pathology is influenced by single nucleotide polymorphisms. Brain Pathology, 2020, 30, 106-119.	4.1	22
86	The Adhesion GPCR CD97/ADGRE5 inhibits apoptosis. International Journal of Biochemistry and Cell Biology, 2015, 65, 197-208.	2.8	21
87	Detection of alternatively spliced EMR2 mRNAs in colorectal tumor cell lines but rare expression of the molecule in colorectal adenocarcinomas. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2003, 443, 32-37.	2.8	20
88	Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunology Letters, 2010, 129, 64-71.	2.5	20
89	lgG Immune Complexes Break Immune Tolerance of Human Microglia. Journal of Immunology, 2020, 205, 2511-2518.	0.8	20
90	CD97 overexpression in tumor cells at the invasion front in colorectal cancer (CC) is independently regulated of the canonical Wnt pathway. Molecular Carcinogenesis, 2006, 45, 881-886.	2.7	19

#	Article	IF	CITATIONS
91	CD70-Driven Chronic Immune Activation Is Protective against Atherosclerosis. Journal of Innate Immunity, 2010, 2, 344-352.	3.8	19
92	SerpinB9 expression in human renal tubular epithelial cells is induced by triggering of the viral dsRNA sensors TLR3, MDA5 and RIG-I. Nephrology Dialysis Transplantation, 2012, 27, 2746-2754.	0.7	19
93	CD55 deposited on synovial collagen fibers protects from immune complex-mediated arthritis. Arthritis Research and Therapy, 2015, 17, 6.	3.5	19
94	GPCRomics of Homeostatic and Disease-Associated Human Microglia. Frontiers in Immunology, 2021, 12, 674189.	4.8	19
95	CD97 is differentially expressed on murine hematopoietic stem-and progenitor-cells. Haematologica, 2008, 93, 1137-1144.	3.5	18
96	Differential expression of CD97 on human lymphocyte subsets and limited effect of CD97 antibodies on allogeneic T-cell stimulation. Immunology Letters, 2009, 123, 160-168.	2.5	18
97	Viral double-stranded RNA sensors induce antiviral, pro-inflammatory, and pro-apoptotic responses in human renal tubular epithelial cells. Kidney International, 2012, 82, 664-675.	5.2	18
98	Dedicated pinhole SPECT of intestinal neutrophil recruitment in a mouse model of dextran sulfate sodium-induced colitis. Journal of Nuclear Medicine, 2005, 46, 526-31.	5.0	18
99	Silencing the Expression of Ras Family GTPase Homologues Decreases Inflammation and Joint Destruction in Experimental Arthritis. American Journal of Pathology, 2010, 177, 3010-3024.	3.8	17
100	<scp>CD</scp> 97 inhibits cell migration in human fibrosarcoma cells by modulating TIMPâ€2/MT1― MMP/MMPâ€2 activity–Ârole of GPS autoproteolysis and functional cooperation between the N―and Câ€ŧerminal fragments. FEBS Journal, 2014, 281, 4878-4891.	4.7	17
101	Macrophages Do Not Express the Phagocytic Receptor BAI1/ADGRB1. Frontiers in Immunology, 2019, 10, 962.	4.8	17
102	CD97 in Leukocyte Trafficking. Advances in Experimental Medicine and Biology, 2010, 706, 128-137.	1.6	17
103	White matter lesions in multiple sclerosis are enriched for CD20 ^{dim} CD8 ⁺ tissueâ€resident memory T cells. European Journal of Immunology, 2021, 51, 483-486.	2.9	16
104	Absence of B Cells in Brainstem and White Matter Lesions Associates With Less Severe Disease and Absence of Oligoclonal Bands in MS. Neurology: Neuroimmunology and NeuroInflammation, 2021, 8, .	6.0	16
105	Transcriptome analysis of normal-appearing white matter reveals cortisol- and disease-associated gene expression profiles in multiple sclerosis. Acta Neuropathologica Communications, 2019, 7, 60.	5.2	15
106	Multiple Targets for Oxysterols in Their Regulation of the Immune System. Cells, 2021, 10, 2078.	4.1	15
107	CD97 antibody depletes granulocytes in mice under conditions of acute inflammation via a Fc receptor-dependent mechanism. Journal of Leukocyte Biology, 2010, 89, 413-421.	3.3	14
108	A Novel Role for CD55 in Granulocyte Homeostasis and Anti-Bacterial Host Defense. PLoS ONE, 2011, 6, e24431.	2.5	14

#	Article	IF	CITATIONS
109	CDK7/12/13 inhibition targets an oscillating leukemia stem cell network and synergizes with venetoclax in acute myeloid leukemia. EMBO Molecular Medicine, 2022, 14, e14990.	6.9	14
110	Mice overexpressing CD97 in intestinal epithelial cells provide a unique model for mammalian postnatal intestinal cylindrical growth. Molecular Biology of the Cell, 2013, 24, 2256-2268.	2.1	13
111	The Interaction of CD97/ADGRE5 With β-Catenin in Adherens Junctions Is Lost During Colorectal Carcinogenesis. Frontiers in Oncology, 2018, 8, 182.	2.8	13
112	Perivascular tissue resident memory T cells as therapeutic target in multiple sclerosis. Expert Review of Neurotherapeutics, 2020, 20, 835-848.	2.8	13
113	Glucocorticoid receptor haplotypes conferring increased sensitivity (Bcll and N363S) are associated with faster progression of multiple sclerosis. Journal of Neuroimmunology, 2016, 299, 84-89.	2.3	12
114	T-cell surveillance of the human brain in health and multiple sclerosis. Seminars in Immunopathology, 2022, 44, 855-867.	6.1	12
115	Sevenspan transmembrane molecules: novel receptors involved in leukocyte adhesion. Immunology Letters, 1996, 54, 185-187.	2.5	11
116	Skeletal Muscle Expression of the Adhesion-GPCR CD97: CD97 Deletion Induces an Abnormal Structure of the Sarcoplasmatic Reticulum but Does Not Impair Skeletal Muscle Function. PLoS ONE, 2014, 9, e100513.	2.5	11
117	CD55 Is Essential for CD103+ Dendritic Cell Tolerogenic Responses that Protect against Autoimmunity. American Journal of Pathology, 2019, 189, 1386-1401.	3.8	11
118	Physical mapping of EMR1 and CD97 in human Chromosome 19 and assignment of Cd97 to mouse Chromosome 8 suggest an ancient genomic duplication. Mammalian Genome, 1999, 10, 1039-1040.	2.2	10
119	Primary Human Renal-Derived Tubular Epithelial Cells Fail to Recognize and Suppress BK Virus Infection. Transplantation, 2017, 101, 1820-1829.	1.0	10
120	Corticosteroid Withdrawal-Induced Loss of Control in Mild to Moderate Asthma Is Independent of Classic Granulocyte Activation. Chest, 2020, 157, 16-25.	0.8	10
121	CD97-DECAY-ACCELERATING FACTOR INTERACTION IS NOT INVOLVED IN LEUKOCYTE ADHESION TO ENDOTHELIAL CELLS1. Transplantation, 2002, 73, 429-436.	1.0	6
122	Introduction: History of the Adhesion GPCR Field. Handbook of Experimental Pharmacology, 2016, 234, 1-11.	1.8	5
123	The EGF-TM7 family of the rat. Immunogenetics, 2004, 56, 679-681.	2.4	4
124	Activation of Wnt signaling in the intestinal mucosa of Apc+/min mice does not cause overexpression of the receptor tyrosine kinase Met. Cancer Science, 2006, 97, 710-715.	3.9	4
125	CD97 Is a Critical Regulator of Acute Myeloid Leukemia Stem Cell Function. Blood, 2016, 128, 1077-1077.	1.4	3
126	Programmed Cell Death Protein 1–Positive CD8 ⁺ T Cells in Multiple Sclerosis. Neurology: Neuroimmunology and NeuroInflammation, 2022, 9, .	6.0	3

#	Article	IF	CITATIONS
127	Intracellular delivery of poly(I:C) induces apoptosis of fibroblast-like synoviocytes via an unknown dsRNA sensor. Biochemical and Biophysical Research Communications, 2016, 477, 343-349.	2.1	2
128	The Inhibitory Receptor GPR56 (Adgrg1) Is Specifically Expressed by Tissue-Resident Memory T Cells in Mice But Dispensable for Their Differentiation and Function In Vivo. Cells, 2021, 10, 2675.	4.1	2
129	Expression of the inhibitory CD200 receptor is associated with alternative macrophage activation. Cytokine, 2009, 48, 35.	3.2	1
130	Phenotypic comparison of human alveolar macrophages before and after in vivo rhinovirus 16 challenge. European Journal of Immunology, 2021, 51, 2691-2693.	2.9	1
131	HIV-1 infection in polarized primary macrophages. Retrovirology, 2011, 8, .	2.0	0
132	Reply: Tissue-resident CD8+ memory T cells in multiple sclerosis. Brain, 2021, 144, e8-e8.	7.6	0
133	Adhesion Class GPCRs in GtoPdb v.2021.3. IUPHAR/BPS Guide To Pharmacology CITE, 2021, 2021, .	0.2	0
134	CD97 Is Differentially Expressed on Murine Hematopoietic Stem Cells and Progenitor Cells Blood, 2005, 106, 2281-2281.	1.4	0
135	Do eosinophils contribute to oxidative stress in mild asthma?. , 2019, , .		0
136	Adhesion Class GPCRs (version 2019.4) in the IUPHAR/BPS Guide to Pharmacology Database. IUPHAR/BPS Guide To Pharmacology CITE, 2019, 2019, .	0.2	0