
Karin M Andersson

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6929245/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	MicroRNA and interleukin 6 interplay in the adipose tissue of rheumatoid arthritis patients. Clinical and Experimental Rheumatology, 2022, , .	0.8	0
2	Impact of the Uncoupling Protein 1 on Cardiovascular Risk in Patients with Rheumatoid Arthritis. Cells, 2021, 10, 1131.	4.1	4
3	Aberrant expression of USF2 in refractory rheumatoid arthritis and its regulation of proinflammatory cytokines in Th17 cells. Proceedings of the National Academy of Sciences of the United States of America, 2020, 117, 30639-30648.	7.1	25
4	Low serum IGF1 is associated with hypertension and predicts early cardiovascular events in women with rheumatoid arthritis. BMC Medicine, 2019, 17, 141.	5.5	20
5	FRI0059â€SOLUBLE RECEPTOR FOR ADVANCED GLYCATION END PRODUCTS (SRAGE) AND RISK FOR CARDIOVASCULAR DISEASES IN FEMALES WITH RHEUMATOID ARTHRITIS. , 2019, , .		0
6	FRI0028â€LOW SERUM IGF1 IS ASSOCIATED WITH AN INCREASED RISK AND HIGH PREVALENCE OF CARDIOVASCULAR EVENTS IN MIDDLE-AGED FEMALE PATIENTS WITH RA – A 5-YEAR FOLLOW-UP. , 2019, , .		0
7	FRI0105â€EXPRESSION OF UNCOUPLING PROTEIN-1 IN SUBCUTANEOUS FAT IS INCREASED BY TOCILIZUMAB. 2019, , .		0
8	THU0012â€ESTROGEN REGULATES MICRO RNA BIOPROCESSING AND PRODUCTION OF IL9 CYTOKINE WITHIN LEUKOCYTES IN RHEUMATOID ARTHRITIS. , 2019, , .		0
9	Inflammation in the hippocampus affects IGF1 receptor signaling and contributes to neurological sequelae in rheumatoid arthritis. Proceedings of the National Academy of Sciences of the United States of America, 2018, 115, E12063-E12072.	7.1	41
10	Smoking activates cytotoxic CD8+ T cells and causes survivin release in rheumatoid arthritis. Journal of Autoimmunity, 2017, 78, 101-110.	6.5	33
11	Survivin controls biogenesis of microRNA in smokers: A link to pathogenesis of rheumatoid arthritis. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2017, 1863, 663-673.	3.8	15
12	IGF-1R signalling contributes to IL-6 production and T cell dependent inflammation in rheumatoid arthritis. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2017, 1863, 2158-2170.	3.8	36
13	Transcriptional signature of human pro-inflammatory TH17 cells identifies reduced IL10 gene expression in multiple sclerosis. Nature Communications, 2017, 8, 1600.	12.8	93
14	01.13â€lgf1R signalling contributes to T cell dependent inflammation in rheumatoid arthritis. , 2017, , .		0
15	08.29â€Smoking contributes to exhaustion state of cd4 ⁺ t cells in rheumatoid arthritis. , 2017, , .		0
16	High Expression of STAT3 in Subcutaneous Adipose Tissue Associates with Cardiovascular Risk in Women with Rheumatoid Arthritis. International Journal of Molecular Sciences, 2017, 18, 2410.	4.1	7
17	Suppressed diversity of survivin splicing in active rheumatoid arthritis. Arthritis Research and Therapy, 2015, 17, 175.	3.5	18
18	S100A4 regulates the Src-tyrosine kinase dependent differentiation of Th17 cells in rheumatoid arthritis. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2014, 1842, 2049-2059.	3.8	11