

# Fernanda O Novais

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6936796/publications.pdf>

Version: 2024-02-01

19  
papers

1,576  
citations

471509

17  
h-index

794594

19  
g-index

20  
all docs

20  
docs citations

20  
times ranked

1762  
citing authors

#	ARTICLE	IF	CITATIONS
1	Cutaneous leishmaniasis: immune responses in protection and pathogenesis. <i>Nature Reviews Immunology</i> , 2016, 16, 581-592.	22.7	467
2	CD8+ T cell cytotoxicity mediates pathology in the skin by inflammasome activation and IL-1 $\beta$ production. <i>PLoS Pathogens</i> , 2017, 13, e1006196.	4.7	160
3	Cytotoxic T Cells Mediate Pathology and Metastasis in Cutaneous Leishmaniasis. <i>PLoS Pathogens</i> , 2013, 9, e1003504.	4.7	130
4	Genomic Profiling of Human <i>Leishmania braziliensis</i> Lesions Identifies Transcriptional Modules Associated with Cutaneous Immunopathology. <i>Journal of Investigative Dermatology</i> , 2015, 135, 94-101.	0.7	130
5	Human Classical Monocytes Control the Intracellular Stage of <i>Leishmania braziliensis</i> by Reactive Oxygen Species. <i>Journal of Infectious Diseases</i> , 2014, 209, 1288-1296.	4.0	99
6	Cutaneous Leishmaniasis Induces a Transmissible Dysbiotic Skin Microbiota that Promotes Skin Inflammation. <i>Cell Host and Microbe</i> , 2017, 22, 13-24.e4.	11.0	82
7	CD8+ T cells in cutaneous leishmaniasis: the good, the bad, and the ugly. <i>Seminars in Immunopathology</i> , 2015, 37, 251-259.	6.1	72
8	Meta-transcriptome Profiling of the Human- <i>Leishmania braziliensis</i> Cutaneous Lesion. <i>PLoS Neglected Tropical Diseases</i> , 2016, 10, e0004992.	3.0	71
9	Variable gene expression and parasite load predict treatment outcome in cutaneous leishmaniasis. <i>Science Translational Medicine</i> , 2019, 11, .	12.4	63
10	Intermediate Monocytes Contribute to Pathologic Immune Response in <i>Leishmania braziliensis</i> Infections. <i>Journal of Infectious Diseases</i> , 2015, 211, 274-282.	4.0	62
11	Lymphocytic Choriomeningitis Virus Expands a Population of NKG2D+CD8+ T Cells That Exacerbates Disease in Mice Coinfected with <i>Leishmania major</i> . <i>Journal of Immunology</i> , 2015, 195, 3301-3310.	0.8	40
12	Matrix Metalloproteinase 9 Production by Monocytes is Enhanced by TNF and Participates in the Pathology of Human Cutaneous Leishmaniasis. <i>PLoS Neglected Tropical Diseases</i> , 2014, 8, e3282.	3.0	36
13	Granzyme B Produced by Natural Killer Cells Enhances Inflammatory Response and Contributes to the Immunopathology of Cutaneous Leishmaniasis. <i>Journal of Infectious Diseases</i> , 2020, 221, 973-982.	4.0	30
14	Phenotypic and functional characteristics of HLA-DR+ neutrophils in Brazilians with cutaneous leishmaniasis. <i>Journal of Leukocyte Biology</i> , 2017, 101, 739-749.	3.3	25
15	CD8+ T Cells Lack Local Signals To Produce IFN- $\gamma$ in the Skin during <i>Leishmania</i> Infection. <i>Journal of Immunology</i> , 2018, 200, 1737-1745.	0.8	24
16	Glyburide, a NLRP3 Inhibitor, Decreases Inflammatory Response and Is a Candidate to Reduce Pathology in <i>Leishmania braziliensis</i> Infection. <i>Journal of Investigative Dermatology</i> , 2020, 140, 246-249.e2.	0.7	24
17	Granzyme B Inhibition by Tofacitinib Blocks the Pathology Induced by CD8 T Cells in Cutaneous Leishmaniasis. <i>Journal of Investigative Dermatology</i> , 2021, 141, 575-585.	0.7	24
18	Host-Directed Therapies for Cutaneous Leishmaniasis. <i>Frontiers in Immunology</i> , 2021, 12, 660183.	4.8	19

#	ARTICLE	IF	CITATIONS
19	Localized skin inflammation during cutaneous leishmaniasis drives a chronic, systemic IFN- $\gamma$ signature. PLoS Neglected Tropical Diseases, 2021, 15, e0009321.	3.0	17