

# Chien-Feng Li

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6954293/publications.pdf>

Version: 2024-02-01

260  
papers

7,973  
citations

50276

46  
h-index

88630

70  
g-index

266  
all docs

266  
docs citations

266  
times ranked

12315  
citing authors

#	ARTICLE	IF	CITATIONS
1	Pharmacological Inactivation of Skp2 SCF Ubiquitin Ligase Restricts Cancer Stem Cell Traits and Cancer Progression. <i>Cell</i> , 2013, 154, 556-568.	28.9	335
2	The Skp2-SCF E3 Ligase Regulates Akt Ubiquitination, Glycolysis, Herceptin Sensitivity, and Tumorigenesis. <i>Cell</i> , 2012, 149, 1098-1111.	28.9	332
3	Deciphering the transcriptional complex critical for RhoA gene expression and cancer metastasis. <i>Nature Cell Biology</i> , 2010, 12, 457-467.	10.3	190
4	Characterization of FN1â€“FGFR1 and novel FN1â€“FGF1 fusion genes in a large series of phosphaturic mesenchymal tumors. <i>Modern Pathology</i> , 2016, 29, 1335-1346.	5.5	139
5	NAB2â€“STAT6 fusion types account for clinicopathological variations in solitary fibrous tumors. <i>Modern Pathology</i> , 2015, 28, 1324-1335.	5.5	133
6	The critical role of AMPK in driving Akt activation under stress, tumorigenesis and drug resistance. <i>Nature Communications</i> , 2018, 9, 4728.	12.8	125
7	Prognostic and Therapeutic Impact of Argininosuccinate Synthetase 1 Control in Bladder Cancer as Monitored Longitudinally by PET Imaging. <i>Cancer Research</i> , 2014, 74, 896-907.	0.9	122
8	Phosphorylation of PDHA by AMPK Drives TCA Cycle to Promote Cancer Metastasis. <i>Molecular Cell</i> , 2020, 80, 263-278.e7.	9.7	120
9	Cancer/stroma interplay via cyclooxygenase-2 and indoleamine 2,3-dioxygenase promotes breast cancer progression. <i>Breast Cancer Research</i> , 2014, 16, 410.	5.0	119
10	<i>ASS1</i> as a Novel Tumor Suppressor Gene in Myxofibrosarcomas: Aberrant Loss via Epigenetic DNA Methylation Confers Aggressive Phenotypes, Negative Prognostic Impact, and Therapeutic Relevance. <i>Clinical Cancer Research</i> , 2013, 19, 2861-2872.	7.0	118
11	Saikosaponin b2 is a naturally occurring terpenoid that efficiently inhibits hepatitis C virus entry. <i>Journal of Hepatology</i> , 2015, 62, 541-548.	3.7	99
12	Skp2-Dependent Ubiquitination and Activation of LKB1 Is Essential for Cancer Cell Survival under Energy Stress. <i>Molecular Cell</i> , 2015, 57, 1022-1033.	9.7	97
13	Skp2â€“MacroH2A1â€“CDK8 axis orchestrates G2/M transition and tumorigenesis. <i>Nature Communications</i> , 2015, 6, 6641.	12.8	87
14	ALK oncoproteins in atypical inflammatory myofibroblastic tumours: novel RRBP1-ALK fusions in epithelioid inflammatory myofibroblastic sarcoma. <i>Journal of Pathology</i> , 2017, 241, 316-323.	4.5	87
15	Overexpression of Nuclear Protein Kinase CK2 Î± Catalytic Subunit (CK2Î±) as a Poor Prognosticator in Human Colorectal Cancer. <i>PLoS ONE</i> , 2011, 6, e17193.	2.5	86
16	A hypoxia-responsive TRAF6â€“ATMâ€“H2AX signalling axis promotes HIF1Î± activation, tumorigenesis and metastasis. <i>Nature Cell Biology</i> , 2017, 19, 38-51.	10.3	83
17	The DNA Damage Transducer RNF8 Facilitates Cancer Chemoresistance and Progression through Twist Activation. <i>Molecular Cell</i> , 2016, 63, 1021-1033.	9.7	82
18	Non-proteolytic ubiquitination of Hexokinase 2 by HectH9 controls tumor metabolism and cancer stem cell expansion. <i>Nature Communications</i> , 2019, 10, 2625.	12.8	82

#	ARTICLE	IF	CITATIONS
19	KDM8/JMJD5 as a dual coactivator of AR and PKM2 integrates AR/EZH2 network and tumor metabolism in CRPC. <i>Oncogene</i> , 2019, 38, 17-32.	5.9	77
20	± <sub>2</sub> -Adrenoceptor agonist dexmedetomidine protects septic acute kidney injury through increasing BMP-7 and inhibiting HDAC2 and HDAC5. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 303, F1443-F1453.	2.7	76
21	Prognostic factors of myxofibrosarcomas: Implications of margin status, tumor necrosis, and mitotic rate on survival. <i>Journal of Surgical Oncology</i> , 2006, 93, 294-303.	1.7	72
22	TOP2A overexpression as a poor prognostic factor in patients with nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 179-187.	1.8	72
23	Decreased succinate dehydrogenase B in human hepatocellular carcinoma accelerates tumor malignancy by inducing the Warburg effect. <i>Scientific Reports</i> , 2018, 8, 3081.	3.3	67
24	TRAF6 Restricts p53 Mitochondrial Translocation, Apoptosis, and Tumor Suppression. <i>Molecular Cell</i> , 2016, 64, 803-814.	9.7	63
25	The clinicopathological significance of <i>NAB2-STAT6</i> gene fusions in 52 cases of intrathoracic solitary fibrous tumors. <i>Cancer Medicine</i> , 2016, 5, 159-168.	2.8	61
26	Anti-IL-20 Monoclonal Antibody Suppresses Breast Cancer Progression and Bone Osteolysis in Murine Models. <i>Journal of Immunology</i> , 2012, 188, 1981-1991.	0.8	60
27	Cycles of Ubiquitination and Deubiquitination Critically Regulate Growth Factor-Mediated Activation of Akt Signaling. <i>Science Signaling</i> , 2013, 6, ra3.	3.6	59
28	Malignant Diffuse-type Tenosynovial Giant Cell Tumors. <i>American Journal of Surgical Pathology</i> , 2008, 32, 587-599.	3.7	57
29	Vav3-Rac1 Signaling Regulates Prostate Cancer Metastasis with Elevated Vav3 Expression Correlating with Prostate Cancer Progression and Posttreatment Recurrence. <i>Cancer Research</i> , 2012, 72, 3000-3009.	0.9	56
30	Skp2 deficiency restricts the progression and stem cell features of castration-resistant prostate cancer by destabilizing Twist. <i>Oncogene</i> , 2017, 36, 4299-4310.	5.9	56
31	Inhibition of the EGFR/STAT3/CEBPD Axis Reverses Cisplatin Cross-resistance with Paclitaxel in the Urothelial Carcinoma of the Urinary Bladder. <i>Clinical Cancer Research</i> , 2017, 23, 503-513.	7.0	56
32	Metformin promotes apoptosis in hepatocellular carcinoma through the CEBPD-induced autophagy pathway. <i>Oncotarget</i> , 2017, 8, 13832-13845.	1.8	56
33	Upregulated IL-19 in Breast Cancer Promotes Tumor Progression and Affects Clinical Outcome. <i>Clinical Cancer Research</i> , 2012, 18, 713-725.	7.0	54
34	Deficiency in expression and epigenetic DNA Methylation of ASS1 gene in nasopharyngeal carcinoma: negative prognostic impact and therapeutic relevance. <i>Tumor Biology</i> , 2014, 35, 161-169.	1.8	54
35	CDCA5 overexpression is an indicator of poor prognosis in patients with urothelial carcinomas of the upper urinary tract and urinary bladder. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 710-22.	0.0	54
36	The E2F Transcription Factor 1 Transactivates Stathmin 1 in Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2013, 20, 4041-4054.	1.5	53

#	ARTICLE	IF	CITATIONS
37	Angiomatoid fibrous histiocytoma: clinicopathological and molecular characterisation with emphasis on variant histomorphology. <i>Journal of Clinical Pathology</i> , 2014, 67, 210-215.	2.0	53
38	Inhibition of HDAC3- and HDAC6-Promoted Survivin Expression Plays an Important Role in SAHA-Induced Autophagy and Viability Reduction in Breast Cancer Cells. <i>Frontiers in Pharmacology</i> , 2016, 7, 81.	3.5	53
39	Targeting chemotherapy-induced PTX3 in tumor stroma to prevent the progression of drug-resistant cancers. <i>Oncotarget</i> , 2015, 6, 23987-24001.	1.8	51
40	Lysine demethylase 2A promotes stemness and angiogenesis of breast cancer by upregulating Jagged1. <i>Oncotarget</i> , 2016, 7, 27689-27710.	1.8	51
41	Skp2 Overexpression Is Highly Representative of Intrinsic Biological Aggressiveness and Independently Associated with Poor Prognosis in Primary Localized Myxofibrosarcomas. <i>Clinical Cancer Research</i> , 2006, 12, 487-498.	7.0	50
42	Primary urothelial carcinoma of the upper tract: Important clinicopathological factors predicting bladder recurrence after surgical resection. <i>Pathology International</i> , 2009, 59, 642-649.	1.3	50
43	ZBRK1 Acts as a Metastatic Suppressor by Directly Regulating <i>MMP9</i> in Cervical Cancer. <i>Cancer Research</i> , 2010, 70, 192-201.	0.9	50
44	Metabolic Stress-Induced Phosphorylation of KAP1 Ser473 Blocks Mitochondrial Fusion in Breast Cancer Cells. <i>Cancer Research</i> , 2016, 76, 5006-5018.	0.9	50
45	Clinicopathological and genetic heterogeneity of the head and neck solitary fibrous tumours: a comparative histological, immunohistochemical and molecular study of 36 cases. <i>Histopathology</i> , 2016, 68, 492-501.	2.9	49
46	Fatal HHV-8-Associated Hemophagocytic Syndrome in an HIV-Negative Immunocompetent Patient With Plasmablastic Variant of Multicentric Castleman Disease (Plasmablastic Microlymphoma). <i>American Journal of Surgical Pathology</i> , 2006, 30, 123-127.	3.7	47
47	Activin A Enhances Prostate Cancer Cell Migration Through Activation of Androgen Receptor and Is Overexpressed in Metastatic Prostate Cancer. <i>Journal of Bone and Mineral Research</i> , 2009, 24, 1180-1193.	2.8	47
48	Transplantation of autologous endothelial progenitor cells in porous PLGA scaffolds create a microenvironment for the regeneration of hyaline cartilage in rabbits. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 1613-1622.	1.3	47
49	Immunohistochemical Expression of p16INK4A, Ki-67, and Mcm2 Proteins in Gastrointestinal Stromal Tumors: Prognostic Implications and Correlations with Risk Stratification of NIH Consensus Criteria. <i>Annals of Surgical Oncology</i> , 2006, 13, 1633-1644.	1.5	46
50	Overexpression of Nuclear Protein Kinase CK2 $\hat{2}$ Subunit and Prognosis in Human Gastric Carcinoma. <i>Annals of Surgical Oncology</i> , 2010, 17, 1695-1702.	1.5	44
51	Overexpression of a Chromatin Remodeling Factor, RSF-1/HBXAP, Correlates with Aggressive Oral Squamous Cell Carcinoma. <i>American Journal of Pathology</i> , 2011, 178, 2407-2415.	3.8	44
52	Prognostic Impact of Thrombospondin-2 (THBS2) Overexpression on Patients with Urothelial Carcinomas of Upper Urinary Tracts and Bladders. <i>Journal of Cancer</i> , 2016, 7, 1541-1549.	2.5	44
53	Skp2 is an independent prognosticator of gallbladder carcinoma among p27Kip1-interacting cell cycle regulators: an immunohistochemical study of 62 cases by tissue microarray. <i>Modern Pathology</i> , 2007, 20, 497-507.	5.5	43
54	IGFBP-5 overexpression as a poor prognostic factor in patients with urothelial carcinomas of upper urinary tracts and urinary bladder. <i>Journal of Clinical Pathology</i> , 2013, 66, 573-582.	2.0	43

#	ARTICLE	IF	CITATIONS
55	The Prognostic Significance of Inflammation-Associated Blood Cell Markers in Patients with Upper Tract Urothelial Carcinoma. <i>Annals of Surgical Oncology</i> , 2016, 23, 343-351.	1.5	43
56	p53 and p27 as predictors of clinical outcome for rectal-cancer patients receiving neoadjuvant therapy. <i>Surgical Oncology</i> , 2006, 15, 211-216.	1.6	42
57	The combined effects of continuous passive motion treatment and acellular PLGA implants on osteochondral regeneration in the rabbit. <i>Biomaterials</i> , 2012, 33, 3153-3163.	11.4	42
58	CCAAT/Enhancer Binding Protein $\beta$ in Macrophages Contributes to Immunosuppression and Inhibits Phagocytosis in Nasopharyngeal Carcinoma. <i>Science Signaling</i> , 2013, 6, ra59.	3.6	42
59	<i>NAB2</i> and <i>STAT6</i> gene fusion and <i>STAT6</i> immunoreexpression in extrathoracic solitary fibrous tumors: the association between fusion variants and locations. <i>Pathology International</i> , 2016, 66, 288-296.	1.3	42
60	Homozygous Deletion of <i>MTAP</i> Gene as a Poor Prognosticator in Gastrointestinal Stromal Tumors. <i>Clinical Cancer Research</i> , 2009, 15, 6963-6972.	7.0	41
61	CEBPD Reverses RB/E2F1-Mediated Gene Repression and Participates in HMDB-Induced Apoptosis of Cancer Cells. <i>Clinical Cancer Research</i> , 2010, 16, 5770-5780.	7.0	41
62	Heat Shock Protein 90 Overexpression Independently Predicts Inferior Disease-Free Survival with Differential Expression of the $\beta$ 1 and $\beta$ 2 Isoforms in Gastrointestinal Stromal Tumors. <i>Clinical Cancer Research</i> , 2008, 14, 7822-7831.	7.0	40
63	Shiunko and Acetylshikonin Promote Reepithelialization, Angiogenesis, and Granulation Tissue Formation in Wounded Skin. <i>The American Journal of Chinese Medicine</i> , 2008, 36, 115-123.	3.8	40
64	Characterization of Gene Amplification-Driven SKP2 Overexpression in Myxofibrosarcoma: Potential Implications in Tumor Progression and Therapeutics. <i>Clinical Cancer Research</i> , 2012, 18, 1598-1610.	7.0	40
65	Overexpression of stathmin 1 confers an independent prognostic indicator in nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 2619-2629.	1.8	40
66	Fibronectin overexpression is associated with latent membrane protein 1 expression and has independent prognostic value for nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 1703-1712.	1.8	40
67	Prognostic Implication of Ezrin Overexpression in Myxofibrosarcomas. <i>Annals of Surgical Oncology</i> , 2010, 17, 3212-3219.	1.5	39
68	INHBA overexpression indicates poor prognosis in urothelial carcinoma of urinary bladder and upper tract. <i>Journal of Surgical Oncology</i> , 2015, 111, 414-422.	1.7	39
69	Interleukin-20 targets renal mesangial cells and is associated with lupus nephritis. <i>Clinical Immunology</i> , 2008, 129, 277-285.	3.2	38
70	Biological significance of chromosomal imbalance aberrations in gastrointestinal stromal tumors. <i>Journal of Biomedical Science</i> , 2004, 11, 65-71.	7.0	37
71	Annexin-I overexpression is associated with tumour progression and independently predicts inferior disease-specific and metastasis-free survival in urinary bladder urothelial carcinoma. <i>Pathology</i> , 2010, 42, 43-49.	0.6	37
72	HuR cytoplasmic expression is associated with increased cyclin A expression and poor outcome with upper urinary tract urothelial carcinoma. <i>BMC Cancer</i> , 2012, 12, 611.	2.6	37

#	ARTICLE	IF	CITATIONS
73	FGF7 Over Expression is an Independent Prognosticator in Patients with Urothelial Carcinoma of the Upper Urinary Tract and Bladder. <i>Journal of Urology</i> , 2015, 194, 223-229.	0.4	37
74	Positive effects of cell-free porous PLGA implants and early loading exercise on hyaline cartilage regeneration in rabbits. <i>Acta Biomaterialia</i> , 2015, 28, 128-137.	8.3	37
75	SPOCK1 Overexpression Confers a Poor Prognosis in Urothelial Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 467-476.	2.5	37
76	Autophagic reliance promotes metabolic reprogramming in oncogenic KRAS-driven tumorigenesis. <i>Autophagy</i> , 2018, 14, 1481-1498.	9.1	37
77	Effect of S-Phase Kinase-Associated Protein 2 Expression on Distant Metastasis and Survival in Nasopharyngeal Carcinoma Patients. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009, 73, 202-207.	0.8	36
78	Nicotinamide N-methyltransferase overexpression is associated with Akt phosphorylation and indicates worse prognosis in patients with nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2013, 34, 3923-3931.	1.8	35
79	A phase 1 study of ADI-PEG 20 and modified FOLFOX6 in patients with advanced hepatocellular carcinoma and other gastrointestinal malignancies. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 82, 429-440.	2.3	35
80	Loss of ZBRK1 Contributes to the Increase of KAP1 and Promotes KAP1-Mediated Metastasis and Invasion in Cervical Cancer. <i>PLoS ONE</i> , 2013, 8, e73033.	2.5	35
81	<i>C</i> EBPD amplification and overexpression in urothelial carcinoma: a driver of tumor metastasis indicating adverse prognosis. <i>Oncotarget</i> , 2015, 6, 31069-31084.	1.8	35
82	Fatty acid synthase overexpression confers an independent prognosticator and associates with radiation resistance in nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2013, 34, 759-768.	1.8	34
83	SKP2 overexpression is associated with a poor prognosis of rectal cancer treated with chemoradiotherapy and represents a therapeutic target with high potential. <i>Tumor Biology</i> , 2013, 34, 1107-1117.	1.8	34
84	Overexpression of thymidylate synthetase confers an independent prognostic indicator in nasopharyngeal carcinoma. <i>Experimental and Molecular Pathology</i> , 2013, 95, 83-90.	2.1	34
85	Interleukin-19 in Breast Cancer. <i>Clinical and Developmental Immunology</i> , 2013, 2013, 1-9.	3.3	34
86	CSF2 Overexpression Is Associated with STAT5 Phosphorylation and Poor Prognosis in Patients with Urothelial Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 711-721.	2.5	34
87	MCM10 overexpression implicates adverse prognosis in urothelial carcinoma. <i>Oncotarget</i> , 2016, 7, 77777-77792.	1.8	34
88	Associations of Rsf-1 overexpression with poor therapeutic response and worse survival in patients with nasopharyngeal carcinoma. <i>Journal of Clinical Pathology</i> , 2012, 65, 248-253.	2.0	32
89	<i>AMACR</i> Amplification in Myxofibrosarcomas: A Mechanism of Overexpression That Promotes Cell Proliferation with Therapeutic Relevance. <i>Clinical Cancer Research</i> , 2014, 20, 6141-6152.	7.0	32
90	Prognostic implication of MET overexpression in myxofibrosarcomas: an integrative array comparative genomic hybridization, real-time quantitative PCR, immunoblotting, and immunohistochemical analysis. <i>Modern Pathology</i> , 2010, 23, 1379-1392.	5.5	31

#	ARTICLE	IF	CITATIONS
91	The expression and prognostic significance of hepatoma-derived growth factor in oral cancer. <i>Oral Oncology</i> , 2012, 48, 629-635.	1.5	31
92	The prognostic impact of lipid biosynthesis-associated markers, HSD17B2 and HMGCS2, in rectal cancer treated with neoadjuvant concurrent chemoradiotherapy. <i>Tumor Biology</i> , 2015, 36, 7675-7683.	1.8	31
93	The cAMP responsive element binding protein 1 transactivates epithelial membrane protein 2, a potential tumor suppressor in the urinary bladder urothelial carcinoma. <i>Oncotarget</i> , 2015, 6, 9220-9239.	1.8	30
94	Overexpression of the PSAT1 Gene in Nasopharyngeal Carcinoma Is an Indicator of Poor Prognosis. <i>Journal of Cancer</i> , 2016, 7, 1088-1094.	2.5	29
95	BCAT1 overexpression is an indicator of poor prognosis in patients with urothelial carcinomas of the upper urinary tract and urinary bladder. <i>Histopathology</i> , 2016, 68, 520-532.	2.9	29
96	DUSP2 regulates extracellular vesicle-VEGF secretion and pancreatic cancer early dissemination. <i>Journal of Extracellular Vesicles</i> , 2020, 9, 1746529.	12.2	29
97	Differentiation between Glioblastoma Multiforme and Primary Cerebral Lymphoma: Additional Benefits of Quantitative Diffusion-Weighted MR Imaging. <i>PLoS ONE</i> , 2016, 11, e0162565.	2.5	29
98	EGFR Nuclear Import in Gallbladder Carcinoma: Nuclear Phosphorylated EGFR Upregulates iNOS Expression and Confers Independent Prognostic Impact. <i>Annals of Surgical Oncology</i> , 2012, 19, 443-454.	1.5	28
99	Enhancer of Zeste Homolog 2 Overexpression in Nasopharyngeal Carcinoma: An Independent Poor Prognosticator That Enhances Cell Growth. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012, 82, 597-604.	0.8	27
100	Autophagy is involved in endogenous and NVP-AUY922-induced KIT degradation in gastrointestinal stromal tumors. <i>Autophagy</i> , 2013, 9, 220-233.	9.1	27
101	GPX2 underexpression indicates poor prognosis in patients with urothelial carcinomas of the upper urinary tract and urinary bladder. <i>World Journal of Urology</i> , 2015, 33, 1777-1789.	2.2	27
102	Overexpressed Fatty Acid Synthase in Gastrointestinal Stromal Tumors: Targeting a Progression-Associated Metabolic Driver Enhances the Antitumor Effect of Imatinib. <i>Clinical Cancer Research</i> , 2017, 23, 4908-4918.	7.0	27
103	CKS1B Overexpression Implicates Clinical Aggressiveness of Hepatocellular Carcinomas but Not p27Kip1 Protein Turnover: an Independent Prognosticator with Potential p27Kip1-Independent Oncogenic Attributes?. <i>Annals of Surgical Oncology</i> , 2010, 17, 907-922.	1.5	26
104	CCAAT/Enhancer-binding Protein $\beta$ Mediates Tumor Necrosis Factor $\alpha$ -induced Aurora Kinase C Transcription and Promotes Genomic Instability. <i>Journal of Biological Chemistry</i> , 2011, 286, 28662-28670.	3.4	26
105	Anesthetic Propofol Overdose Causes Vascular Hyperpermeability by Reducing Endothelial Glycocalyx and ATP Production. <i>International Journal of Molecular Sciences</i> , 2015, 16, 12092-12107.	4.1	26
106	HAS3 underexpression as an indicator of poor prognosis in patients with urothelial carcinoma of the upper urinary tract and urinary bladder. <i>Tumor Biology</i> , 2015, 36, 5441-5450.	1.8	26
107	The diagnostic ureteroscopy before radical nephroureterectomy in upper urinary tract urothelial carcinoma is not associated with higher intravesical recurrence. <i>World Journal of Surgical Oncology</i> , 2018, 16, 135.	1.9	26
108	Sulfatase-1 overexpression indicates poor prognosis in urothelial carcinoma of the urinary bladder and upper tract. <i>Oncotarget</i> , 2017, 8, 47216-47229.	1.8	26



#	ARTICLE	IF	CITATIONS
109	Ezrin overexpression in gastrointestinal stromal tumors: an independent adverse prognosticator associated with the non-gastric location. <i>Modern Pathology</i> , 2009, 22, 1351-1360.	5.5	25
110	Overexpression of CPS1 is an independent negative prognosticator in rectal cancers receiving concurrent chemoradiotherapy. <i>Tumor Biology</i> , 2014, 35, 11097-11105.	1.8	25
111	Overexpression of DNAJC12 predicts poor response to neoadjuvant concurrent chemoradiotherapy in patients with rectal cancer. <i>Experimental and Molecular Pathology</i> , 2015, 98, 338-345.	2.1	25
112	Oct4 upregulates osteopontin via Egr1 and is associated with poor outcome in human lung cancer. <i>BMC Cancer</i> , 2019, 19, 791.	2.6	25
113	Arginine starvation elicits chromatin leakage and cGAS-STING activation via epigenetic silencing of metabolic and DNA-repair genes. <i>Theranostics</i> , 2021, 11, 7527-7545.	10.0	25
114	Rsf-1 expression in rectal cancer: with special emphasis on the independent prognostic value after neoadjuvant chemoradiation. <i>Journal of Clinical Pathology</i> , 2012, 65, 687-692.	2.0	24
115	DPP4/CD26 overexpression in urothelial carcinoma confers an independent prognostic impact and correlates with intrinsic biological aggressiveness. <i>Oncotarget</i> , 2017, 8, 2995-3008.	1.8	24
116	Recurrent Amplification at 7q21.2 Targets CDK6 Gene in Primary Myxofibrosarcomas and Identifies CDK6 Overexpression as an Independent Adverse Prognosticator. <i>Annals of Surgical Oncology</i> , 2012, 19, 2716-2725.	1.5	23
117	Fibroblast growth factor receptor 2 overexpression is predictive of poor prognosis in rectal cancer patients receiving neoadjuvant chemoradiotherapy. <i>Journal of Clinical Pathology</i> , 2014, 67, 1056-1061.	2.0	23
118	Galectin-1 Dysregulation Independently Predicts Disease Specific Survival in Bladder Urothelial Carcinoma. <i>Journal of Urology</i> , 2015, 193, 1002-1008.	0.4	23
119	Rsf-1/HBXAP overexpression is associated with disease-specific survival of patients with gallbladder carcinoma. <i>Apmis</i> , 2011, 119, 808-814.	2.0	22
120	Transmembrane and Coiled-Coil Domain 1 Impairs the AKT Signaling Pathway in Urinary Bladder Urothelial Carcinoma: A Characterization of a Tumor Suppressor. <i>Clinical Cancer Research</i> , 2017, 23, 7650-7663.	7.0	22
121	Hypoxia-regulated MicroRNA-210 Overexpression is Associated with Tumor Development and Progression in Upper Tract Urothelial Carcinoma. <i>International Journal of Medical Sciences</i> , 2017, 14, 578-584.	2.5	22
122	AMACR amplification and overexpression in primary imatinib-naïve gastrointestinal stromal tumors: a driver of cell proliferation indicating adverse prognosis. <i>Oncotarget</i> , 2014, 5, 11588-11603.	1.8	22
123	Loss of lactate dehydrogenase B subunit expression is correlated with tumour progression and independently predicts inferior disease-specific survival in urinary bladder urothelial carcinoma. <i>Pathology</i> , 2011, 43, 707-712.	0.6	21
124	Role of galectin-1 in urinary bladder urothelial carcinoma cell invasion through the JNK pathway. <i>Cancer Science</i> , 2016, 107, 1390-1398.	3.9	21
125	IL-18-induced interaction between IMP3 and HuR contributes to COX-2 mRNA stabilization in acute myeloid leukemia. <i>Journal of Leukocyte Biology</i> , 2016, 99, 131-141.	3.3	21
126	Over-Expression of CHD4 Is an Independent Biomarker of Poor Prognosis in Patients with Rectal Cancers Receiving Concurrent Chemoradiotherapy. <i>International Journal of Molecular Sciences</i> , 2019, 20, 4087.	4.1	21



#	ARTICLE	IF	CITATIONS
127	High TNFAIP6 level is associated with poor prognosis of urothelial carcinomas. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019, 37, 293.e11-293.e24.	1.6	21
128	Downregulated <i>MTAP</i> expression in myxofibrosarcoma: A characterization of inactivating mechanisms, tumor suppressive function, and therapeutic relevance. <i>Oncotarget</i> , 2014, 5, 11428-11441.	1.8	21
129	MTOR inhibition enhances NVP-ALY922-induced autophagy-mediated KIT degradation and cytotoxicity in imatinib-resistant gastrointestinal stromal tumors. <i>Oncotarget</i> , 2014, 5, 11723-11736.	1.8	21
130	<i>Candida</i> invasion and influences in smoking patients with multiple oral leucoplakias – a retrospective study. <i>Mycoses</i> , 2011, 54, e377-83.	4.0	20
131	Involvement of phorbol-12-myristate-13-acetate-induced protein 1 in goniiothalamine-induced TP53-dependent and -independent apoptosis in hepatocellular carcinoma-derived cells. <i>Toxicology and Applied Pharmacology</i> , 2011, 256, 8-23.	2.8	20
132	Selecting Tyrosine Kinase Inhibitors for Gastrointestinal Stromal Tumor with Secondary KIT Activation-Loop Domain Mutations. <i>PLoS ONE</i> , 2013, 8, e65762.	2.5	20
133	Chromatophagy: Autophagy goes nuclear and captures broken chromatin during arginine-starvation. <i>Autophagy</i> , 2015, 11, 419-421.	9.1	20
134	Role of Microtubule-Associated Protein 1b in Urothelial Carcinoma: Overexpression Predicts Poor Prognosis. <i>Cancers</i> , 2020, 12, 630.	3.7	20
135	Phase 1b study of pegylated arginine deiminase (ADI-PEG 20) plus Pembrolizumab in advanced solid cancers. <i>Oncimmunology</i> , 2021, 10, 1943253.	4.6	20
136	PLCB4 copy gain and PLC $\beta$ 4 overexpression in primary gastrointestinal stromal tumors: Integrative characterization of a lipid-catabolizing enzyme associated with worse disease-free survival. <i>Oncotarget</i> , 2017, 8, 19997-20010.	1.8	20
137	HMDB and 5-AzadC Combination Reverses Tumor Suppressor CCAAT/Enhancer-Binding Protein Delta to Strengthen the Death of Liver Cancer Cells. <i>Molecular Cancer Therapeutics</i> , 2015, 14, 2623-2633.	4.1	19
138	A microtubule inhibitor, ABT-751, induces autophagy and delays apoptosis in Huh-7 cells. <i>Toxicology and Applied Pharmacology</i> , 2016, 311, 88-98.	2.8	19
139	Metformin Resensitizes Sorafenib-Resistant HCC Cells Through AMPK-Dependent Autophagy Activation. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 596655.	3.7	19
140	Clinicopathological significance of HuR expression in gallbladder carcinoma: with special emphasis on the implications of its nuclear and cytoplasmic expression. <i>Tumor Biology</i> , 2013, 34, 3059-3069.	1.8	18
141	Deficiency in asparagine synthetase expression in rectal cancers receiving concurrent chemoradiotherapy: negative prognostic impact and therapeutic relevance. <i>Tumor Biology</i> , 2014, 35, 6823-6830.	1.8	18
142	Overexpression of Transcobalamin 1 is an Independent Negative Prognosticator in Rectal Cancers Receiving Concurrent Chemoradiotherapy. <i>Journal of Cancer</i> , 2017, 8, 1330-1337.	2.5	18
143	Hepatoma-derived growth factor supports the antiapoptosis and profibrosis of pancreatic stellate cells. <i>Cancer Letters</i> , 2019, 457, 180-190.	7.2	18
144	SLC14A1 prevents oncometabolite accumulation and recruits HDAC1 to transrepress oncometabolite genes in urothelial carcinoma. <i>Theranostics</i> , 2020, 10, 11775-11793.	10.0	18

#	ARTICLE	IF	CITATIONS
145	Overexpression of ANXA1 confers independent negative prognostic impact in rectal cancers receiving concurrent chemoradiotherapy. <i>Tumor Biology</i> , 2014, 35, 7755-7763.	1.8	17
146	Downregulation of RNF128 Predicts Progression and Poor Prognosis in Patients with Urothelial Carcinoma of the Upper Tract and Urinary Bladder. <i>Journal of Cancer</i> , 2016, 7, 2187-2196.	2.5	17
147	Matrix metalloproteinase-11 as a marker of metastasis and predictor of poor survival in urothelial carcinomas. <i>Journal of Surgical Oncology</i> , 2016, 113, 700-707.	1.7	17
148	Prediction of progression in skull base meningiomas: additional benefits of apparent diffusion coefficient value. <i>Journal of Neuro-Oncology</i> , 2018, 138, 63-71.	2.9	17
149	YWHAZ amplification/overexpression defines aggressive bladder cancer and contributes to chemo- and radio-resistance by suppressing caspase-mediated apoptosis. <i>Journal of Pathology</i> , 2019, 248, 476-487.	4.5	17
150	Lysine demethylase 2A expression in cancer-associated fibroblasts promotes breast tumour growth. <i>British Journal of Cancer</i> , 2021, 124, 484-493.	6.4	17
151	Disruption of the pentraxin 3/CD44 interaction as an efficient therapy for triple-negative breast cancers. <i>Clinical and Translational Medicine</i> , 2022, 12, e724.	4.0	17
152	Glutamate Decarboxylase 1 Overexpression as a Poor Prognostic Factor in Patients with Nasopharyngeal Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 1716-1723.	2.5	16
153	Overexpression of Transient Receptor Protein Cation Channel Subfamily A Member 1, Confers an Independent Prognostic Indicator in Nasopharyngeal Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 1181-1188.	2.5	16
154	Alpha-methylacyl coenzyme A racemase overexpression in gallbladder carcinoma confers an independent prognostic indicator. <i>Journal of Clinical Pathology</i> , 2012, 65, 309-314.	2.0	15
155	Rsf-1/HBXAP overexpression is independent of gene amplification and is associated with poor outcome in patients with urinary bladder urothelial carcinoma. <i>Journal of Clinical Pathology</i> , 2012, 65, 802-807.	2.0	15
156	Inhibiting Interleukin-19 Activity Ameliorates Esophageal Squamous Cell Carcinoma Progression. <i>PLoS ONE</i> , 2013, 8, e75254.	2.5	15
157	PLA2G2A overexpression is associated with poor therapeutic response and inferior outcome in rectal cancer patients receiving neoadjuvant concurrent chemoradiotherapy. <i>Histopathology</i> , 2015, 66, 991-1002.	2.9	15
158	Overexpression of regulator of G protein signaling 11 promotes cell migration and associates with advanced stages and aggressiveness of lung adenocarcinoma. <i>Oncotarget</i> , 2016, 7, 31122-31136.	1.8	15
159	Subcellular localisation of anillin is associated with different survival outcomes in upper urinary tract urothelial carcinoma. <i>Journal of Clinical Pathology</i> , 2015, 68, 1026-1032.	2.0	14
160	High Expression of EphA4 Predicted Lesser Degree of Tumor Regression after Neoadjuvant Chemoradiotherapy in Rectal Cancer. <i>Journal of Cancer</i> , 2017, 8, 1089-1096.	2.5	14
161	Nuclear KIT induces a NFKB1B-RELA-KIT autoregulatory loop in imatinib-resistant gastrointestinal stromal tumors. <i>Oncogene</i> , 2019, 38, 6550-6565.	5.9	14
162	Downregulation of the cytochrome P450 4B1 protein confers a poor prognostic factor in patients with urothelial carcinomas of upper urinary tracts and urinary bladder. <i>Apms</i> , 2019, 127, 170-180.	2.0	14

#	ARTICLE	IF	CITATIONS
163	Suppression of Extracellular Vesicle VEGF-Ca <sup>2+</sup> -mediated Lymphangiogenesis and Pancreatic Cancer Early Dissemination By a Selective HDAC1/2 Inhibitor. <i>Molecular Cancer Therapeutics</i> , 2021, 20, 1550-1560.	4.1	14
164	Overexpression of REG4 confers an independent negative prognosticator in rectal cancers receiving concurrent chemoradiotherapy. <i>Journal of Surgical Oncology</i> , 2014, 110, 1002-1010.	1.7	13
165	Ductal activation of oncogenic KRAS alone induces sarcomatoid phenotype. <i>Scientific Reports</i> , 2015, 5, 13347.	3.3	13
166	Complement Component 1, s Subcomponent Overexpression is an Independent Poor Prognostic Indicator in Patients with Urothelial Carcinomas of the Upper Urinary Tract and Urinary Bladder. <i>Journal of Cancer</i> , 2016, 7, 1396-1405.	2.5	13
167	High Expression of Aldolase B Confers a Poor Prognosis for Rectal Cancer Patients Receiving Neoadjuvant Chemoradiotherapy. <i>Journal of Cancer</i> , 2017, 8, 1197-1204.	2.5	13
168	High Expression of UBE2B as a Poor Prognosis Factor in Patients With Rectal Cancer Following Chemoradiotherapy. <i>Anticancer Research</i> , 2020, 40, 6305-6317.	1.1	13
169	Adult NTRK-rearranged spindle cell neoplasms of the viscera: with an emphasis on rare locations and heterologous elements. <i>Modern Pathology</i> , 2022, 35, 911-921.	5.5	13
170	Overexpression of JAK2: a predictor of unfavorable prognosis for nasopharyngeal carcinoma. <i>Future Oncology</i> , 2016, 12, 1887-1896.	2.4	12
171	Amplification-driven BCL6-suppressed cytostasis is mediated by transrepression of FOXO3 and post-translational modifications of FOXO3 in urinary bladder urothelial carcinoma. <i>Theranostics</i> , 2020, 10, 707-724.	10.0	12
172	BMI1-KLF4 axis deficiency improves responses to neoadjuvant concurrent chemoradiotherapy in patients with rectal cancer. <i>Radiotherapy and Oncology</i> , 2020, 149, 249-258.	0.6	12
173	O6-methylguanine-DNA methyltransferase modulates cisplatin-induced DNA double-strand breaks by targeting the homologous recombination pathway in nasopharyngeal carcinoma. <i>Journal of Biomedical Science</i> , 2021, 28, 2.	7.0	12
174	Transcriptomic reappraisal identifies <i>MGLL</i> overexpression as an unfavorable prognosticator in primary gastrointestinal stromal tumors. <i>Oncotarget</i> , 2016, 7, 49986-49997.	1.8	12
175	The prognostic impact of RAP2A expression in patients with early and locoregionally advanced nasopharyngeal carcinoma in an endemic area. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 912-21.	0.0	12
176	Preprogramming therapeutic response of PI3K/mTOR dual inhibitor via the regulation of EHMT2 and p27 in pancreatic cancer. <i>American Journal of Cancer Research</i> , 2018, 8, 1812-1822.	1.4	12
177	Loss of epithelial membrane protein-2 expression confers an independent prognosticator in nasopharyngeal carcinoma: a cohort study. <i>BMJ Open</i> , 2012, 2, e000900.	1.9	11
178	A rare malignant tumor of scalp in a 3-month-old Taiwanese infancy: case report of primitive myxoid mesenchymal tumor of infancy with molecular study. <i>Medical Molecular Morphology</i> , 2013, 46, 109-113.	1.0	11
179	Ectopic hamartomatous thymoma is distinct from lipomatous pleomorphic adenoma in lacking <i>PLAG</i> aberration. <i>Histopathology</i> , 2013, 62, 518-522.	2.9	11
180	Necdin Overexpression Predicts Poor Prognosis in Patients with Urothelial Carcinomas of the Upper Urinary Tract and Urinary Bladder. <i>Journal of Cancer</i> , 2016, 7, 304-313.	2.5	11

#	ARTICLE	IF	CITATIONS
181	Prognostic Value of Leptin Receptor Overexpression in Upper Tract Urothelial Carcinomas in Taiwan. <i>Clinical Genitourinary Cancer</i> , 2017, 15, e653-e659.	1.9	11
182	Deciphering the Molecular Mechanism Underlying the Inhibitory Efficacy of Taiwanese Local Pomegranate Peels against Urinary Bladder Urothelial Carcinoma. <i>Nutrients</i> , 2018, 10, 543.	4.1	11
183	Utility of EFEMP1 in the Prediction of Oncologic Outcomes of Urothelial Carcinoma. <i>Genes</i> , 2021, 12, 872.	2.4	11
184	DDR2 overexpression in urothelial carcinoma indicates an unfavorable prognosis: a large cohort study. <i>Oncotarget</i> , 2016, 7, 78918-78931.	1.8	11
185	Flow Cytometric Analysis of DNA Ploidy and S-Phase Fraction in Primary Localized Myxofibrosarcoma: Correlations with Clinicopathological Factors, Skp2 Expression, and Patient Survival. <i>Annals of Surgical Oncology</i> , 2008, 15, 2239-2249.	1.5	10
186	Overexpression of CDC28 protein kinase regulatory subunit 1B confers an independent prognostic factor in nasopharyngeal carcinoma. <i>Apmis</i> , 2014, 122, 206-214.	2.0	10
187	Aspiration cytology of an ectopic cervical thymoma misinterpreted as a lymphoproliferative lesion of the thyroid: A case report. <i>Oncology Letters</i> , 2015, 10, 1255-1258.	1.8	10
188	PTP4A3 Independently Predicts Metastasis and Survival in Upper Tract Urothelial Carcinoma Treated with Radical Nephroureterectomy. <i>Journal of Urology</i> , 2015, 194, 1449-1455.	0.4	10
189	Is preoperative anemia a risk factor for upper tract urothelial carcinoma following radical nephroureterectomy?. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016, 34, 337.e1-337.e9.	1.6	10
190	Inflammatory myofibroblastic tumor with RANBP2 and ALK gene rearrangement with bland cytological features mimicking desmoid-type fibromatosis: A case report and review of the literature. <i>Oncology Letters</i> , 2016, 11, 1429-1434.	1.8	10
191	Higher nuclear EGFR expression is a better predictor of survival in rectal cancer patients following neoadjuvant chemoradiotherapy than cytoplasmic EGFR expression. <i>Oncology Letters</i> , 2018, 17, 1551-1558.	1.8	10
192	Low thrombospondin 2 expression is predictive of low tumor regression after neoadjuvant chemoradiotherapy in rectal cancer. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 2423-32.	0.0	10
193	BCL6 overexpression is associated with decreased p19ARF expression and confers an independent prognosticator in gallbladder carcinoma. <i>Tumor Biology</i> , 2014, 35, 1417-1426.	1.8	9
194	Cyclin-dependent kinase 4 overexpression is mostly independent of gene amplification and constitutes an independent prognosticator for nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 7209-7216.	1.8	9
195	Laparoscopic partial nephrectomy without intracorporeal suturing. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2016, 30, 1585-1591.	2.4	9
196	Hydroxysteroid 11-Beta Dehydrogenase 1 Overexpression with Copy-Number Gain and Missense Mutations in Primary Gastrointestinal Stromal Tumors. <i>Journal of Clinical Medicine</i> , 2018, 7, 408.	2.4	9
197	High FRMD3 expression is prognostic for worse survival in rectal cancer patients treated with CCRT. <i>International Journal of Clinical Oncology</i> , 2021, 26, 1689-1697.	2.2	9
198	Angiogenesis Driven by the CEBPDâ€“hsa-miR-429â€“VEGFA Signaling Axis Promotes Urothelial Carcinoma Progression. <i>Cells</i> , 2022, 11, 638.	4.1	9

#	ARTICLE	IF	CITATIONS
199	Gastrointestinal stromal tumor (GIST) in southern Taiwan: A clinicopathologic study of 93 resected cases. <i>Pathology Research and Practice</i> , 2005, 201, 1-9.	2.3	8
200	Expression of cyclin-dependent kinase 2-associated protein 1 confers an independent prognosticator in nasopharyngeal carcinoma: a cohort study. <i>Journal of Clinical Pathology</i> , 2012, 65, 795-801.	2.0	8
201	The Effect of Osteochondral Regeneration Using Polymer Constructs and Continuous Passive Motion Therapy in the Lower Weight-Bearing Zone of Femoral Trochlear Groove in Rabbits. <i>Annals of Biomedical Engineering</i> , 2013, 41, 385-397.	2.5	8
202	Draft genome sequence of CTX-M-type $\beta$ -lactamase-producing <i>Klebsiella quasipneumoniae</i> subsp. <i>similipneumoniae</i> isolated from a Box turtle. <i>Journal of Global Antimicrobial Resistance</i> , 2018, 12, 235-236.	2.2	8
203	Therapeutic benefit of second-look transurethral resection of bladder tumors for newly diagnosed T1 bladder cancer: a single-center experience. <i>International Urology and Nephrology</i> , 2019, 51, 1335-1342.	1.4	8
204	MIC $\beta$ 6 is essential for promoting glucose metabolic reprogramming and tumor growth in triple-negative breast cancer. <i>EMBO Reports</i> , 2021, 22, e50781.	4.5	8
205	Overexpression of Pyruvate Dehydrogenase Kinase-3 Predicts Poor Prognosis in Urothelial Carcinoma. <i>Frontiers in Oncology</i> , 2021, 11, 749142.	2.8	8
206	Molecular Characterization of Dermatofibrosarcoma Protuberans. <i>American Journal of Surgical Pathology</i> , 2022, 46, 942-955.	3.7	8
207	HuR cytoplasmic expression is associated with increased cyclin A expression and inferior disease-free survival in patients with gastrointestinal stromal tumours (<sc>GIST</sc>s). <i>Histopathology</i> , 2013, 63, 445-454.	2.9	7
208	Galectin-1 overexpression in nasopharyngeal carcinoma: effect on survival. <i>Acta Oto-Laryngologica</i> , 2014, 134, 536-542.	0.9	7
209	AMACR overexpression as a poor prognostic factor in patients with nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 7983-7991.	1.8	7
210	Prognostic significance of DSG3 in rectal adenocarcinoma treated with preoperative chemoradiotherapy. <i>Future Oncology</i> , 2016, 12, 1457-1467.	2.4	7
211	Anti-IL-20 monoclonal antibody inhibited tumor growth in hepatocellular carcinoma. <i>Scientific Reports</i> , 2017, 7, 17609.	3.3	7
212	Proteomic Identification of the Galectin-1-Involved Molecular Pathways in Urinary Bladder Urothelial Carcinoma. <i>International Journal of Molecular Sciences</i> , 2018, 19, 1242.	4.1	7
213	Deduction of Novel Genes Potentially Involved in Upper Tract Urothelial Carcinoma Using Next-Generation Sequencing and Bioinformatics Approaches. <i>International Journal of Medical Sciences</i> , 2019, 16, 93-105.	2.5	7
214	Prognostic Utility of FBLN2 Expression in Patients With Urothelial Carcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 570340.	2.8	7
215	High SPINK4 Expression Predicts Poor Outcomes among Rectal Cancer Patients Receiving CCRT. <i>Current Oncology</i> , 2021, 28, 2373-2384.	2.2	7
216	Characterization and Prognostic Significance of Methylthioadenosine Phosphorylase Deficiency in Nasopharyngeal Carcinoma. <i>Medicine (United States)</i> , 2015, 94, e2271.	1.0	6

#	ARTICLE	IF	CITATIONS
217	Malignant Peripheral Nerve Sheath Tumor of Prostate: A Rare Case Report and Literature Review. Case Reports in Urology, 2016, 2016, 1-3.	0.3	6
218	Overexpression of PTP4A3 is associated with metastasis and unfavorable prognosis in bladder cancer. World Journal of Urology, 2016, 34, 835-846.	2.2	6
219	Periostin overexpression is associated with worse prognosis in nasopharyngeal carcinoma from endemic area: a cohort study. OncoTargets and Therapy, 2018, Volume 11, 3205-3213.	2.0	6
220	Expression and prognostic utility of SSX2IP in patients with nasopharyngeal carcinoma. Apmis, 2020, 128, 287-297.	2.0	6
221	Upregulated MUC2 Is an Unfavorable Prognostic Indicator for Rectal Cancer Patients Undergoing Preoperative CCRT. Journal of Clinical Medicine, 2021, 10, 3030.	2.4	6
222	Inhibition of the formation of autophagosome but not autolysosome augments ABT-751-induced apoptosis in TP53-deficient Hep3B cells. Journal of Cellular Physiology, 2019, 234, 9551-9563.	4.1	5
223	A biosensor of high-density lipoprotein of human serum on a liquid crystal and polymer composite film. Proceedings of SPIE, 2013, , .	0.8	4
224	High Expression of 17 $\beta$ -hydroxysteroid Dehydrogenase Type 2 is Associated with a Better Prognosis in Urothelial Carcinoma of the Urinary Tract. Journal of Cancer, 2016, 7, 2221-2230.	2.5	4
225	Myoepithelial carcinoma of tibia mimic giant cell tumor: a case report with emphasis on MR features. Skeletal Radiology, 2019, 48, 1637-1641.	2.0	4
226	RSF1 requires CEBP $\beta$ and hSNF2H to promote IL-1 $\beta$ -mediated angiogenesis: the clinical and therapeutic relevance of RSF1 overexpression and amplification in myxofibrosarcomas. Angiogenesis, 2021, 24, 533-548.	7.2	4
227	Complete and Durable Response to Nivolumab in Recurrent Poorly Differentiated Pancreatic Neuroendocrine Carcinoma with High Tumor Mutational Burden. Current Oncology, 2021, 28, 4587-4596.	2.2	4
228	Myoepithelial and oral intracranial myxoid mesenchymal tumor-like neoplasms as diagnostic considerations of the ever-expanding extracranial myxocollagenous tumors harboring FET-CREB fusions. Pathology Research and Practice, 2021, 229, 153700.	2.3	4
229	High Level of Aristolochic Acid Detected With a Unique Genomic Landscape Predicts Early UTUC Onset After Renal Transplantation in Taiwan. Frontiers in Oncology, 2021, 11, 828314.	2.8	4
230	Biological significance of MYC and CEBPD coamplification in urothelial carcinoma: Multilayered genomic, transcriptional and posttranscriptional positive feedback loops enhance oncogenic glycolysis. Clinical and Translational Medicine, 2021, 11, e674.	4.0	4
231	Identification of phenothiazine as an ETV1-targeting agent in gastrointestinal stromal tumors using the Connectivity Map. International Journal of Oncology, 2019, 55, 536-546.	3.3	3
232	Metformin Resensitizes Sorafenib-Resistant HCC Cells Through AMPK-Dependent Autophagy Activation. Frontiers in Cell and Developmental Biology, 2021, 8, .	3.7	3
233	Utility of Acute Physiology and Chronic Health Evaluation (APACHE II) in Predicting Mortality in Patients with Pyogenic Liver Abscess: A Retrospective Study. Journal of Clinical Medicine, 2021, 10, 2644.	2.4	3
234	CTSE Overexpression Is an Adverse Prognostic Factor for Survival among Rectal Cancer Patients Receiving CCRT. Life, 2021, 11, 646.	2.4	3



#	ARTICLE	IF	CITATIONS
235	EMP2 induces cytostasis and apoptosis via the TGF $\beta$ <sup>2</sup> /SMAD/SP1 axis and recruitment of P2RX7 in urinary bladder urothelial carcinoma. <i>Cellular Oncology (Dordrecht)</i> , 2021, 44, 1133-1150.	4.4	3
236	Prognostic Significance of ROR2 Expression in Patients with Urothelial Carcinoma. <i>Biomedicines</i> , 2021, 9, 1054.	3.2	3
237	The protective effects of glutamine in a rat model of ventilator-induced lung injury. <i>Journal of Thoracic Disease</i> , 2014, 6, 1704-13.	1.4	3
238	Open the Technical Black Box of Tumor Mutational Burden (TMB): Factors Affecting Harmonization and Standardization of Panel-Based TMB. <i>International Journal of Molecular Sciences</i> , 2022, 23, 5097.	4.1	3
239	Clinical Aggressiveness of Myxofibrosarcomas Associates with Down-Regulation of p12CDK2AP1: Prognostic Implication of a Putative Tumor Suppressor that Induces Cell Cycle Arrest and Apoptosis Via Mitochondrial Pathway. <i>Annals of Surgical Oncology</i> , 2014, 21, 711-720.	1.5	2
240	An Effective and Well Tolerated Strategy of Bladder Preservation Therapy in Cisplatin-Ineligible Patients With Muscle-Invasive Bladder Cancer. <i>Clinical Genitourinary Cancer</i> , 2016, 14, e67-e74.	1.9	2
241	Expression of prothymosin $\beta$ in lung cancer is associated with squamous cell carcinoma and smoking. <i>Oncology Letters</i> , 2019, 17, 5740-5746.	1.8	2
242	Treatment of chronic hepatitis C regimens containing with recombinant interferon in patients with sustained virological response predicts risk of hepatocellular carcinoma. <i>Medicine (United States)</i> , 2020, 99, e22435.	1.0	2
243	Identification of potential genes in upper tract urothelial carcinoma using next-generation sequencing with bioinformatics and in vitro analyses. <i>PeerJ</i> , 2021, 9, e11343.	2.0	2
244	High G protein subunit beta 4 protein level is correlated to poor prognosis of urothelial carcinoma. <i>Medical Molecular Morphology</i> , 2021, 54, 356-367.	1.0	2
245	A phase II study of ADI-PEG 20 and FOLFOX6 in patients (pts) with advanced hepatocellular carcinoma (HCC).. <i>Journal of Clinical Oncology</i> , 2019, 37, TPS477-TPS477.	1.6	2
246	Role of second-look transurethral resection of bladder tumors for newly diagnosed T1 bladder cancer: Experience at a single center. <i>Urological Science</i> , 2018, 29, 95.	0.6	2
247	Loss of epithelial membrane protein-2 expression confers an independent prognosticator in gallbladder carcinoma. <i>Biomarkers and Genomic Medicine</i> , 2013, 5, 31-38.	0.2	1
248	Molecular biology of urothelial carcinoma. <i>Journal of Cancer Research and Practice</i> , 2021, 8, 1.	0.2	1
249	S100P as a marker for poor survival and advanced stage in gallbladder carcinoma. <i>Annals of Diagnostic Pathology</i> , 2021, 52, 151736.	1.3	1
250	Advances in molecular genetics of early-stage urothelial carcinoma. <i>Translational Cancer Research</i> , 2016, 5, S1126-S1130.	1.0	1
251	High Stromal SFRP2 Expression in Urothelial Carcinoma Confers an Unfavorable Prognosis. <i>Frontiers in Oncology</i> , 2022, 12, 834249.	2.8	1
252	Characterization of Aberrations in DNA Damage Repair Pathways in Gastrointestinal Stromal Tumors: The Clinicopathologic Relevance of $\beta$ -H2AX and 53BP1 in Correlation with Heterozygous Deletions of CHEK2, BRCA2, and RB1. <i>Cancers</i> , 2022, 14, 1787.	3.7	1



#	ARTICLE	IF	CITATIONS
253	Discrimination of Methicillin-resistant Staphylococcus aureus by MALDI-TOF Mass Spectrometry with Machine Learning Techniques in Patients with Staphylococcus aureus Bacteremia. Pathogens, 2022, 11, 586.	2.8	1
254	High MT2A Expression Predicts Worse Prognosis in Patients with Urothelial Carcinoma. Oncology, 2022, 100, 485-497.	1.9	1
255	E2F transcription factor 1 overexpression as a poor prognostic factor in patients with nasopharyngeal carcinomas. Biomarkers and Genomic Medicine, 2013, 5, 23-30.	0.2	0
256	Dedifferentiated liposarcoma can induce a leukemoid reaction. Formosan Journal of Surgery, 2014, 47, 42-48.	0.2	0
257	MIG-6 Is Essential for Promoting Glucose Metabolic Reprogramming and Tumor Growth in Triple-Negative Breast Cancer. FASEB Journal, 2021, 35, .	0.5	0
258	High glycosyltransferase-8 domain containing two protein levels contribute to poor prognosis in urothelial carcinoma. International Journal of Urology, 2021, 28, 1178-1187.	1.0	0
259	ADI-PEG 20 and FOLFOX6: A phase 1 study in pts (pts) with advanced hepatocellular carcinoma (HCC).. Journal of Clinical Oncology, 2018, 36, 4085-4085.	1.6	0
260	Abstract 1452: Hypoxia-induced MIG-6 promotes glucose metabolic reprogramming and tumorigenesis in triple-negative breast cancer. , 2020, , .		0