

Lucy C Robinson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/695564/publications.pdf>

Version: 2024-02-01

26
papers

1,540
citations

394421

19
h-index

552781

26
g-index

26
all docs

26
docs citations

26
times ranked

1247
citing authors

#	ARTICLE	IF	CITATIONS
1	Reg1 and Snf1 regulate stress-induced relocalization of protein phosphatase-1 to cytoplasmic granules. <i>FEBS Journal</i> , 2021, 288, 4833-4848.	4.7	5
2	Î±-Synuclein inhibits Snx3-retromer retrograde trafficking of the conserved membrane-bound proprotein convertase Kex2 in the secretory pathway of <i>Saccharomyces cerevisiae</i> . <i>Human Molecular Genetics</i> , 2021, , .	2.9	2
3	SDS22 selectively recognizes and traps metal-deficient inactive PP1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 20472-20481.	7.1	28
4	<i>Saccharomyces cerevisiae</i> Mhr1 can bind Xho I-induced mitochondrial DNA double-strand breaks in vivo. <i>Mitochondrion</i> , 2018, 42, 23-32.	3.4	5
5	New ubiquitin-dependent mechanisms regulating the Aurora B-Protein Phosphatase 1 balance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2018, 131, .	2.0	2
6	Evidence for double-strand break mediated mitochondrial DNA replication in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2017, 45, 7760-7773.	14.5	20
7	Molecular mechanics and dynamics characterization of an <i>in silico</i> mutated protein: A stand-alone lab module or support activity for <i>in vivo</i> and <i>in vitro</i> analyses of targeted proteins. <i>Biochemistry and Molecular Biology Education</i> , 2013, 41, 402-408.	1.2	11
8	Suppressors of <i>ipl1-2</i> in Components of a Glc7 Phosphatase Complex, Cdc48 AAA ATPase, TORC1, and the Kinetochore. <i>G3: Genes, Genomes, Genetics</i> , 2012, 2, 1687-1701.	1.8	10
9	Temperature-sensitive <i>ipl1-2/Aurora</i> B mutation is suppressed by mutations in TOR complex 1 via the Glc7/PP1 phosphatase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 3994-3999.	7.1	27
10	GABA acts as a ligand chaperone in the early secretory pathway to promote cell surface expression of GABAA receptors. <i>Brain Research</i> , 2010, 1346, 1-13.	2.2	39
11	A Molecular Genetics Laboratory Course Applying Bioinformatics and Cell Biology in the Context of Original Research. <i>CBE Life Sciences Education</i> , 2008, 7, 410-421.	2.3	27
12	Novel suppressors of Î±-synuclein toxicity identified using yeast. <i>Human Molecular Genetics</i> , 2008, 17, 3784-3795.	2.9	58
13	Glc7-Reg1 Phosphatase Signals to Yck1,2 Casein Kinase 1 to Regulate Transport Activity and Glucose-Induced Inactivation of <i>Saccharomyces</i> Maltose Permease. <i>Genetics</i> , 2006, 172, 1427-1439.	2.9	38
14	Akr1p-dependent Palmitoylation of Yck2p Yeast Casein Kinase 1 Is Necessary and Sufficient for Plasma Membrane Targeting. <i>Journal of Biological Chemistry</i> , 2004, 279, 27138-27147.	3.4	59
15	Constitutive GABAA Receptor Endocytosis Is Dynamin-mediated and Dependent on a Dilucine AP2 Adaptin-binding Motif within the Î²2 Subunit of the Receptor. <i>Journal of Biological Chemistry</i> , 2003, 278, 24046-24052.	3.4	83
16	Use of green fluorescent protein in living yeast cells. <i>Methods in Enzymology</i> , 2002, 351, 661-683.	1.0	13
17	Plasma membrane localization of the Yck2p yeast casein kinase 1 isoform requires the C-terminal extension and secretory pathway function. <i>Journal of Cell Science</i> , 2002, 115, 4957-4968.	2.0	50
18	The Yck2 Yeast Casein Kinase 1 Isoform Shows Cell Cycle-specific Localization to Sites of Polarized Growth and Is Required for Proper Septin Organization. <i>Molecular Biology of the Cell</i> , 1999, 10, 1077-1092.	2.1	63

#	ARTICLE	IF	CITATIONS
19	Functional characterization and visualization of a GABAA receptor-GFP chimera expressed in <i>Xenopus</i> oocytes. <i>Molecular Brain Research</i> , 1998, 59, 165-177.	2.3	29
20	Activation of Protein Kinase C Induces \hat{I}^3 -Aminobutyric Acid Type A Receptor Internalization in <i>Xenopus</i> Oocytes. <i>Journal of Biological Chemistry</i> , 1998, 273, 32595-32601.	3.4	101
21	Casein Kinase \hat{I}^3 Subfamily.. <i>Journal of Biological Chemistry</i> , 1995, 270, 12717-12724.	3.4	99
22	TFS1: A suppressor of <i>cdc25</i> mutations in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1991, 230, 241-250.	2.4	52
23	CDC25: a component of the RAS-adenylate cyclase pathway in <i>Saccharomyces cerevisiae</i> . <i>Science</i> , 1987, 235, 1218-1221.	12.6	258
24	Mapping of the <i>Saccharomyces cerevisiae</i> CDC3, CDC25, and CDC42 genes to chromosome XII by chromosome blotting and tetrad analysis. <i>Yeast</i> , 1987, 3, 243-253.	1.7	35
25	RAS2 of <i>Saccharomyces cerevisiae</i> is required for gluconeogenic growth and proper response to nutrient limitation.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1985, 82, 3785-3789.	7.1	192
26	Mammalian and yeast ras gene products: biological function in their heterologous systems. <i>Science</i> , 1985, 228, 179-184.	12.6	234