

Thomas Miedaner

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6960763/publications.pdf>

Version: 2024-02-01

179
papers

6,568
citations

66234

42
h-index

91712

69
g-index

179
all docs

179
docs citations

179
times ranked

4083
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Molecular mapping of QTLs for Fusarium head blight resistance in spring wheat. II. Resistance to fungal penetration and spread. <i>Theoretical and Applied Genetics</i> , 2003, 107, 503-508. | 1.8 | 285 |
| 2 | Developments in breeding cereals for organic agriculture. <i>Euphytica</i> , 2008, 163, 323. | 0.6 | 285 |
| 3 | Involvement of trichothecenes in fusarioses of wheat, barley and maize evaluated by gene disruption of the trichodiene synthase (Tri5) gene in three field isolates of different chemotype and virulence. <i>Molecular Plant Pathology</i> , 2006, 7, 449-461. | 2.0 | 266 |
| 4 | Marker-Assisted Selection for Disease Resistance in Wheat and Barley Breeding. <i>Phytopathology</i> , 2012, 102, 560-566. | 1.1 | 223 |
| 5 | Revealing the genetic architecture of FHB resistance in hexaploid wheat (<i>Triticum aestivum</i> L.) by QTL meta-analysis. <i>Molecular Breeding</i> , 2009, 23, 473-488. | 1.0 | 203 |
| 6 | Molecular mapping of Fusarium head blight resistance in the winter wheat population Dream/Lynx. <i>Theoretical and Applied Genetics</i> , 2005, 111, 747-756. | 1.8 | 137 |
| 7 | Stacking quantitative trait loci (QTL) for Fusarium head blight resistance from non-adapted sources in an European elite spring wheat background and assessing their effects on deoxynivalenol (DON) content and disease severity. <i>Theoretical and Applied Genetics</i> , 2006, 112, 562-569. | 1.8 | 133 |
| 8 | Effects of genotype and genotype-environment interaction on deoxynivalenol accumulation and resistance to Fusarium head blight in rye, triticale, and wheat. <i>Plant Breeding</i> , 2001, 120, 97-105. | 1.0 | 128 |
| 9 | Mapping QTLs with main and epistatic effects underlying grain yield and heading time in soft winter wheat. <i>Theoretical and Applied Genetics</i> , 2011, 123, 283-292. | 1.8 | 124 |
| 10 | A European Database of Fusarium graminearum and F. culmorum Trichothecene Genotypes. <i>Frontiers in Microbiology</i> , 2016, 7, 406. | 1.5 | 124 |
| 11 | Detection of segregation distortion loci in triticale (x <i>Triticosecale</i> Wittmack) based on a high-density DArT marker consensus genetic linkage map. <i>BMC Genomics</i> , 2011, 12, 380. | 1.2 | 113 |
| 12 | Biology, Genetics, and Management of Ergot (<i>Claviceps</i> spp.) in Rye, Sorghum, and Pearl Millet. <i>Toxins</i> , 2015, 7, 659-678. | 1.5 | 111 |
| 13 | Population Genetics of Three Important Head Blight Pathogens <i>Fusarium graminearum</i> , <i>F. pseudograminearum</i> and <i>F. culmorum</i> . <i>Journal of Phytopathology</i> , 2008, 156, 129-139. | 0.5 | 108 |
| 14 | Genetic Mapping of Pathogenicity and Aggressiveness of <i>Gibberella zeae</i> (<i>Fusarium graminearum</i>) Toward Wheat. <i>Phytopathology</i> , 2004, 94, 520-526. | 1.1 | 93 |
| 15 | Inheritance of resistance to Fusarium head blight in three European winter wheat populations. <i>Theoretical and Applied Genetics</i> , 2008, 117, 1119-1128. | 1.8 | 91 |
| 16 | Genome-Wide Association Study Identifies Novel Candidate Genes for Aggressiveness, Deoxynivalenol Production, and Azole Sensitivity in Natural Field Populations of <i>Fusarium graminearum</i> . <i>Molecular Plant-Microbe Interactions</i> , 2016, 29, 417-430. | 1.4 | 89 |
| 17 | Comparison of phenotypic and marker-based selection for Fusarium head blight resistance and DON content in spring wheat. <i>Molecular Breeding</i> , 2007, 19, 357-370. | 1.0 | 86 |
| 18 | Multiple-trait- and selection indices-genomic predictions for grain yield and protein content in rye for feeding purposes. <i>Theoretical and Applied Genetics</i> , 2016, 129, 273-287. | 1.8 | 86 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | A comparison of aggressiveness and deoxynivalenol production between Canadian <i>Fusarium graminearum</i> isolates with 3-acetyl and 15-acetyldeoxynivalenol chemotypes in field-grown spring wheat. <i>European Journal of Plant Pathology</i> , 2010, 127, 407-417. | 0.8 | 84 |
| 20 | Effect of Dwarfing <i>Rht</i> Genes on <i>Fusarium</i> Head Blight Resistance in Two Sets of Near-Isogenic Lines of Wheat and Check Cultivars. <i>Crop Science</i> , 2008, 48, 2115-2122. | 0.8 | 76 |
| 21 | Association mapping for <i>Fusarium</i> head blight resistance in European soft winter wheat. <i>Molecular Breeding</i> , 2011, 28, 647-655. | 1.0 | 70 |
| 22 | Climate change will influence disease resistance breeding in wheat in Northwestern Europe. <i>Theoretical and Applied Genetics</i> , 2021, 134, 1771-1785. | 1.8 | 70 |
| 23 | Genetics of Resistance and Pathogenicity in the Maize/ <i>Setosphaeria turcica</i> Pathosystem and Implications for Breeding. <i>Frontiers in Plant Science</i> , 2017, 8, 1490. | 1.7 | 69 |
| 24 | The accuracy of prediction of genomic selection in elite hybrid rye populations surpasses the accuracy of marker-assisted selection and is equally augmented by multiple field evaluation locations and test years. <i>BMC Genomics</i> , 2014, 15, 556. | 1.2 | 68 |
| 25 | Relatedness severely impacts accuracy of marker-assisted selection for disease resistance in hybrid wheat. <i>Heredity</i> , 2014, 112, 552-561. | 1.2 | 67 |
| 26 | Genetic architecture of complex agronomic traits examined in two testcross populations of rye (<i>Secale cereale</i> L.). <i>BMC Genomics</i> , 2012, 13, 706. | 1.2 | 66 |
| 27 | Genetic architecture of resistance to <i>Septoria tritici</i> blotch in European wheat. <i>BMC Genomics</i> , 2013, 14, 858. | 1.2 | 62 |
| 28 | Genetic Variation for Resistance to Ear Rots and Mycotoxins Contamination in Early European Maize Inbred Lines. <i>Crop Science</i> , 2009, 49, 2019-2028. | 0.8 | 60 |
| 29 | Diversity in genetic structure and chemotype composition of <i>Fusarium graminearum sensu stricto</i> populations causing wheat head blight in individual fields in Germany. <i>European Journal of Plant Pathology</i> , 2011, 131, 39-48. | 0.8 | 57 |
| 30 | Comparative mapping of DNA sequences in rye (<i>Secale cereale</i> L.) in relation to the rice genome. <i>Theoretical and Applied Genetics</i> , 2009, 118, 371-384. | 1.8 | 56 |
| 31 | Mapping of genes for male-fertility restoration in â€™Pampaâ€™ CMS winter rye (<i>Secale cereale</i> L.). <i>Theoretical and Applied Genetics</i> , 2000, 101, 1226-1233. | 1.8 | 53 |
| 32 | Deoxynivalenol (DON) Content and <i>Fusarium</i> Head Blight Resistance in Segregating Populations of Winter Rye and Winter Wheat. <i>Crop Science</i> , 2003, 43, 519. | 0.8 | 53 |
| 33 | Genetic variation of aggressiveness in individual field populations of <i>Fusarium graminearum</i> and <i>Fusarium culmorum</i> tested on young plants of winter rye. <i>European Journal of Plant Pathology</i> , 1996, 102, 823-830. | 0.8 | 51 |
| 34 | Estimation of deoxynivalenol (DON) content by symptom rating and exoantigen content for resistance selection in wheat and triticale. <i>Euphytica</i> , 2004, 139, 123-132. | 0.6 | 50 |
| 35 | Accuracy of within- and among-family genomic prediction for <i>Fusarium</i> head blight and <i>Septoria tritici</i> blotch in winter wheat. <i>Theoretical and Applied Genetics</i> , 2019, 132, 1121-1135. | 1.8 | 50 |
| 36 | Establishment of introgression libraries in hybrid rye (<i>Secale cereale</i> L.) from an Iranian primitive accession as a new tool for rye breeding and genomics. <i>Theoretical and Applied Genetics</i> , 2008, 117, 641-652. | 1.8 | 49 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Effect of the <i>Rht1</i> dwarfing locus on <i>Fusarium</i> head blight rating in three segregating populations of winter wheat. <i>Plant Breeding</i> , 2008, 127, 333-339. | 1.0 | 49 |
| 38 | Population parameters for resistance to <i>Fusarium graminearum</i> and <i>Fusarium verticillioides</i> ear rot among large sets of early, mid-late and late maturing European maize (<i>Zea mays</i> L.) inbred lines. <i>Theoretical and Applied Genetics</i> , 2010, 120, 1053-1062. | 1.8 | 49 |
| 39 | REML approach for adjusting the <i>Fusarium</i> head blight rating to a phenological date in inoculated selection experiments of wheat. <i>Theoretical and Applied Genetics</i> , 2008, 117, 65-73. | 1.8 | 48 |
| 40 | Sources of resistance to <i>Fusarium</i> head blight within Syrian durum wheat landraces. <i>Plant Breeding</i> , 2011, 130, 398-400. | 1.0 | 47 |
| 41 | Genomics-assisted breeding for ear rot resistances and reduced mycotoxin contamination in maize: methods, advances and prospects. <i>Theoretical and Applied Genetics</i> , 2019, 132, 2721-2739. | 1.8 | 45 |
| 42 | Colocalization of QTL for <i>Gibberella</i> Ear Rot Resistance and Low Mycotoxin Contamination in Early European Maize. <i>Crop Science</i> , 2011, 51, 1935-1945. | 0.8 | 44 |
| 43 | Aggressiveness and mycotoxin production of eight isolates each of <i>Fusarium graminearum</i> and <i>Fusarium verticillioides</i> for ear rot on susceptible and resistant early maize inbred lines. <i>European Journal of Plant Pathology</i> , 2010, 127, 113-123. | 0.8 | 43 |
| 44 | Agronomic and Quality Performance of Winter Wheat Backcross Populations Carrying Non-Adapted <i>Fusarium</i> Head Blight Resistance QTL. <i>Crop Science</i> , 2010, 50, 2283-2290. | 0.8 | 43 |
| 45 | Quantitative Trait Loci for Adult-Plant Resistance to <i>Mycosphaerella graminicola</i> in Two Winter Wheat Populations. <i>Phytopathology</i> , 2011, 101, 1209-1216. | 1.1 | 43 |
| 46 | Testcross performance of rye introgression lines developed by marker-assisted backcrossing using an Iranian accession as donor. <i>Theoretical and Applied Genetics</i> , 2009, 118, 1225-1238. | 1.8 | 42 |
| 47 | Broad-spectrum resistance loci for three quantitatively inherited diseases in two winter wheat populations. <i>Molecular Breeding</i> , 2012, 29, 731-742. | 1.0 | 42 |
| 48 | Marker-based introduction of three quantitative-trait loci conferring resistance to <i>Fusarium</i> head blight into an independent elite winter wheat breeding population. <i>Theoretical and Applied Genetics</i> , 2008, 117, 29-35. | 1.8 | 41 |
| 49 | Marker selection for <i>Fusarium</i> head blight resistance based on quantitative trait loci (QTL) from two European sources compared to phenotypic selection in winter wheat. <i>Euphytica</i> , 2009, 166, 219-227. | 0.6 | 41 |
| 50 | Global warming and increasing maize cultivation demand comprehensive efforts in disease and insect resistance breeding in north-western Europe. <i>Plant Pathology</i> , 2021, 70, 1032-1046. | 1.2 | 40 |
| 51 | Development of PCR-based markers linked to dominant genes for male-fertility restoration in Pampa CMS of rye (<i>Secale cereale</i> L.). <i>Theoretical and Applied Genetics</i> , 2003, 106, 1184-1190. | 1.8 | 35 |
| 52 | Hybrid rye performance under natural drought stress in Europe. <i>Theoretical and Applied Genetics</i> , 2013, 126, 475-482. | 1.8 | 35 |
| 53 | Mycotoxin accumulation and corresponding ear rot rating in three maturity groups of European maize inoculated by two <i>Fusarium</i> species. <i>Euphytica</i> , 2010, 174, 153-164. | 0.6 | 34 |
| 54 | 8 Biology, Diversity, and Management of FHB-Causing <i>Fusarium</i> Species in Small-Grain Cereals. , 2013, , 199-241. | | 34 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Genetic architecture is more complex for resistance to <i>Septoria tritici</i> blotch than to <i>Fusarium</i> head blight in Central European winter wheat. <i>BMC Genomics</i> , 2015, 16, 430. | 1.2 | 34 |
| 56 | Integration of genotypic, hyperspectral, and phenotypic data to improve biomass yield prediction in hybrid rye. <i>Theoretical and Applied Genetics</i> , 2020, 133, 3001-3015. | 1.8 | 34 |
| 57 | Competition Effects Among Isolates of <i>Fusarium culmorum</i> Differing in Aggressiveness and Mycotoxin Production on Heads of Winter Rye. <i>European Journal of Plant Pathology</i> , 2004, 110, 63-70. | 0.8 | 33 |
| 58 | Segregation for aggressiveness and deoxynivalenol production of a population of <i>Gibberella zeae</i> causing head blight of wheat. <i>European Journal of Plant Pathology</i> , 2004, 110, 789-799. | 0.8 | 33 |
| 59 | Molecular mapping of quantitative trait loci for field resistance to <i>Fusarium</i> head blight in a European winter wheat population. <i>Plant Breeding</i> , 2008, 127, 459-464. | 1.0 | 33 |
| 60 | Effect of a rye dwarfing gene on plant height, heading stage, and <i>Fusarium</i> head blight in triticale (<i>Å-Triticosecale</i> Wittmack). <i>Theoretical and Applied Genetics</i> , 2014, 127, 1527-1536. | 1.8 | 33 |
| 61 | Genomic prediction and GWAS of <i>Gibberella</i> ear rot resistance traits in dent and flint lines of a public maize breeding program. <i>Euphytica</i> , 2018, 214, 1. | 0.6 | 32 |
| 62 | QTL mapping and comparative genome analysis of agronomic traits including grain yield in winter rye. <i>Theoretical and Applied Genetics</i> , 2017, 130, 1801-1817. | 1.8 | 31 |
| 63 | Choice of models for QTL mapping with multiple families and design of the training set for prediction of <i>Fusarium</i> resistance traits in maize. <i>Theoretical and Applied Genetics</i> , 2016, 129, 431-444. | 1.8 | 30 |
| 64 | Copy number variation of Ppd-B1 is the major determinant of heading time in durum wheat. <i>BMC Genetics</i> , 2019, 20, 64. | 2.7 | 30 |
| 65 | Impact of genotype, harvest time and chemical composition on the methane yield of winter rye for biogas production. <i>Biomass and Bioenergy</i> , 2011, 35, 4316-4323. | 2.9 | 29 |
| 66 | Covariation between line and testcross performance for reduced mycotoxin concentrations in European maize after silk channel inoculation of two <i>Fusarium</i> species. <i>Theoretical and Applied Genetics</i> , 2011, 122, 925-934. | 1.8 | 29 |
| 67 | Genetic architecture of plant height in winter rye introgression libraries. <i>Plant Breeding</i> , 2011, 130, 209-216. | 1.0 | 28 |
| 68 | Genetic Architecture of <i>Fusarium</i> Head Blight Resistance in Four Winter Triticale Populations. <i>Phytopathology</i> , 2015, 105, 334-341. | 1.1 | 28 |
| 69 | Dynamic quantitative trait loci (QTL) for plant height predict biomass yield in hybrid rye (<i>Secale</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 | 2.9 | 28 |
| 70 | An experimental approach for estimating the genomic selection advantage for <i>Fusarium</i> head blight and <i>Septoria tritici</i> blotch in winter wheat. <i>Theoretical and Applied Genetics</i> , 2019, 132, 2425-2437. | 1.8 | 28 |
| 71 | Genomics-Assisted Breeding for Quantitative Disease Resistances in Small-Grain Cereals and Maize. <i>International Journal of Molecular Sciences</i> , 2020, 21, 9717. | 1.8 | 28 |
| 72 | Comparative Quantitative Trait Loci Mapping for <i>Gibberella</i> Ear Rot Resistance and Reduced Deoxynivalenol Contamination across Connected Maize Populations. <i>Crop Science</i> , 2012, 52, 32-43. | 0.8 | 27 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Genetic variation and covariation for aggressiveness, deoxynivalenol production and fungal colonization among progeny of <i>Gibberella zeae</i> in wheat. <i>Plant Pathology</i> , 2004, 53, 446-453. | 1.2 | 26 |
| 74 | Effectiveness and environmental stability of quantitative powdery mildew (<i>Blumeria graminis</i>) resistance among winter wheat cultivars. <i>Plant Breeding</i> , 2007, 126, 553-558. | 1.0 | 26 |
| 75 | Virulence phenotypes in powdery mildew (<i>Blumeria graminis</i>) populations and resistance genes in triticale (x <i>Triticosecale</i>). <i>European Journal of Plant Pathology</i> , 2013, 137, 463-476. | 0.8 | 26 |
| 76 | Genetic variation for resistance to <i>Fusarium</i> head blight in winter durum material. <i>Crop and Pasture Science</i> , 2014, 65, 46. | 0.7 | 26 |
| 77 | Rht24 reduces height in the winter wheat population "SolitÃr"–"Bussard"™ without adverse effects on <i>Fusarium</i> head blight infection. <i>Theoretical and Applied Genetics</i> , 2018, 131, 1263-1272. | 1.8 | 26 |
| 78 | Estimates of additive and dominance effects for <i>Fusarium</i> head blight resistance of winter triticale. <i>Plant Breeding</i> , 2004, 123, 525-530. | 1.0 | 25 |
| 79 | Within-Field Variation of <i>Fusarium graminearum</i> Isolates for Aggressiveness and Deoxynivalenol Production in Wheat Head Blight. <i>Phytopathology</i> , 2012, 102, 128-134. | 1.1 | 25 |
| 80 | First insights into the genotype–phenotype map of phenotypic stability in rye. <i>Journal of Experimental Botany</i> , 2015, 66, 3275-3284. | 2.4 | 25 |
| 81 | Use of non-adapted quantitative trait loci for increasing <i>Fusarium</i> head blight resistance for breeding semi-dwarf wheat. <i>Plant Breeding</i> , 2019, 138, 140-147. | 1.0 | 25 |
| 82 | Combining ability of non-adapted sources for male-fertility restoration in Pampa CMS of hybrid rye*. <i>Plant Breeding</i> , 2005, 124, 39-43. | 1.0 | 24 |
| 83 | Variation and Transgression of Aggressiveness Among Two <i>Gibberella zeae</i> Crosses Developed from Highly Aggressive Parental Isolates. <i>Phytopathology</i> , 2010, 100, 904-912. | 1.1 | 24 |
| 84 | The potential of genomic-assisted breeding to improve <i>Fusarium</i> head blight resistance in winter durum wheat. <i>Plant Breeding</i> , 2017, 136, 610-619. | 1.0 | 24 |
| 85 | High accuracy of predicting hybrid performance of <i>Fusarium</i> head blight resistance by mid-parent values in wheat. <i>Theoretical and Applied Genetics</i> , 2017, 130, 461-470. | 1.8 | 24 |
| 86 | Genomic predictions for <i>Fusarium</i> head blight resistance in a diverse durum wheat panel: an effective incorporation of plant height and heading date as covariates. <i>Euphytica</i> , 2020, 216, 1. | 0.6 | 24 |
| 87 | Association between line per se and testcross performance for eight agronomic and quality traits in winter rye. <i>Theoretical and Applied Genetics</i> , 2014, 127, 33-41. | 1.8 | 23 |
| 88 | Fine mapping of the restorer gene Rfp3 from an Iranian primitive rye (<i>Secale cereale</i> L.). <i>Theoretical and Applied Genetics</i> , 2017, 130, 1179-1189. | 1.8 | 23 |
| 89 | Editorial: Management of <i>Fusarium</i> Species and their Mycotoxins in Cereal Food and Feed. <i>Frontiers in Microbiology</i> , 2017, 8, 1543. | 1.5 | 23 |
| 90 | Hybrid Breeding in Rye (<i>Secale cereale</i> L.). , 2019, , 343-372. | | 23 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Molecular Variation and Genetic Structure in Field Populations of Fusarium Species Causing Head Blight in Wheat. <i>Cereal Research Communications</i> , 1997, 25, 549-554. | 0.8 | 23 |
| 92 | Genetic variation for resistance to ergot (<i>Claviceps purpurea</i> [Fr.] Tul.) among full-sib families of five populations of winter rye (<i>Secale cereale</i> L.). <i>Theoretical and Applied Genetics</i> , 2008, 118, 85-90. | 1.8 | 22 |
| 93 | Association of single nucleotide polymorphic sites in candidate genes with aggressiveness and deoxynivalenol production in <i>Fusarium graminearum</i> causing wheat head blight. <i>BMC Genetics</i> , 2012, 13, 14. | 2.7 | 22 |
| 94 | Quantitative-genetic analysis of leaf-rust resistance in seedling and adult-plant stages of inbred lines and their testcrosses in winter rye. <i>Plant Breeding</i> , 2002, 121, 475-479. | 1.0 | 21 |
| 95 | Covariation of Ergot Severity and Alkaloid Content Measured by HPLC and One ELISA Method in Inoculated Winter Rye across Three Isolates and Three European Countries. <i>Toxins</i> , 2020, 12, 676. | 1.5 | 21 |
| 96 | Selection for Fusarium head blight resistance in early generations reduces the deoxynivalenol (DON) content in grain of winter and spring wheat. <i>Plant Breeding</i> , 2006, 125, 96-98. | 1.0 | 20 |
| 97 | Means and variances for Fusarium head blight resistance of F2-derived bulks from winter triticale and winter wheat crosses. <i>Euphytica</i> , 2006, 152, 405-411. | 0.6 | 20 |
| 98 | Genetic variation for ergot (<i>Claviceps purpurea</i>) resistance and alkaloid concentrations in cytoplasmic-male sterile winter rye under pollen isolation. <i>Euphytica</i> , 2010, 173, 299-306. | 0.6 | 20 |
| 99 | Genetic variation of winter rye cultivars for their ergot (<i>Claviceps purpurea</i>) reaction tested in a field design with minimized interplot interference. <i>Plant Breeding</i> , 2010, 129, 58-62. | 1.0 | 20 |
| 100 | Correlation between Fusarium head blight severity and DON content in triticale as revealed by phenotypic and molecular data. <i>Plant Breeding</i> , 2016, 135, 31-37. | 1.0 | 20 |
| 101 | Low validation rate of quantitative trait loci for <i>Gibberella</i> ear rot resistance in European maize. <i>Theoretical and Applied Genetics</i> , 2017, 130, 175-186. | 1.8 | 20 |
| 102 | Early Detection of <i>Zymoseptoria tritici</i> in Winter Wheat by Infrared Thermography. <i>Agriculture (Switzerland)</i> , 2019, 9, 139. | 1.4 | 20 |
| 103 | Genome-wide association mapping and genomic prediction of Fusarium head blight resistance, heading stage and plant height in winter rye (<i>Secale cereale</i>). <i>Plant Breeding</i> , 2020, 139, 508-520. | 1.0 | 20 |
| 104 | Breeding progress of disease resistance and impact of disease severity under natural infections in winter wheat variety trials. <i>Theoretical and Applied Genetics</i> , 2021, 134, 1281-1302. | 1.8 | 19 |
| 105 | A model calculation approach towards the optimization of a standard scheme of seed-parent line development in hybrid rye breeding. <i>Plant Breeding</i> , 2008, 127, 433-440. | 1.0 | 18 |
| 106 | Identification of genomic regions carrying QTL for agronomic and quality traits in rye (<i>Secale</i>) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50 1 | 1.0 | 18 |
| 107 | Inheritance of resistance to <i>Gibberella</i> ear rot and deoxynivalenol contamination in five flint maize crosses. <i>Plant Breeding</i> , 2012, 131, 28-32. | 1.0 | 18 |
| 108 | Exploiting genetic diversity in two European maize landraces for improving <i>Gibberella</i> ear rot resistance using genomic tools. <i>Theoretical and Applied Genetics</i> , 2021, 134, 793-805. | 1.8 | 18 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Geography and end use drive the diversification of worldwide winter rye populations. <i>Molecular Ecology</i> , 2016, 25, 500-514. | 2.0 | 17 |
| 110 | Breeding Strategies for Improving Plant Resistance to Diseases. , 2016, , 561-599. | | 17 |
| 111 | Genetic architecture of yellow and stem rust resistance in a durum wheat diversity panel. <i>Euphytica</i> , 2019, 215, 1. | 0.6 | 17 |
| 112 | Molecular tracking of multiple disease resistance in a winter wheat diversity panel. <i>Theoretical and Applied Genetics</i> , 2020, 133, 419-431. | 1.8 | 17 |
| 113 | Comparison of rye, triticale, durum wheat and bread wheat genotypes for Fusarium head blight resistance and deoxynivalenol contamination. <i>Plant Breeding</i> , 2020, 139, 251-262. | 1.0 | 17 |
| 114 | Snow mold of winter cereals: a complex disease and a challenge for resistance breeding. <i>Theoretical and Applied Genetics</i> , 2021, 134, 419-433. | 1.8 | 17 |
| 115 | Prediction of hybrid performance for Fusarium head blight resistance in triticale (Triticosecale) Tj ETQq1 1 0.784314 rgBT /Overloc | 0.6 | 16 |
| 116 | Correlated effects of exotic pollen fertility restorer genes on agronomic and quality traits of hybrid rye. <i>Plant Breeding</i> , 2017, 136, 224-229. | 1.0 | 16 |
| 117 | Genome-wide association study for an efficient selection of Fusarium head blight resistance in winter triticale. <i>Euphytica</i> , 2019, 215, 1. | 0.6 | 16 |
| 118 | Dwarfing gene Rht24 does not affect Fusarium head blight resistance in a large European winter wheat diversity panel. <i>Euphytica</i> , 2022, 218, 1. | 0.6 | 16 |
| 119 | Diversity, spatial variation, and temporal dynamics of virulences in the German leaf rust (<i>Puccinia</i>) Tj ETQq1 1 0.784314 rgBT /Overloc | 0.8 | 15 |
| 120 | Head-blighting populations of <i>Fusarium culmorum</i> from Germany, Russia, and Syria analyzed by microsatellite markers show a recombining structure. <i>European Journal of Plant Pathology</i> , 2013, 137, 743-752. | 0.8 | 15 |
| 121 | Prediction of deoxynivalenol and zearalenone concentrations in <i>Fusarium graminearum</i> inoculated backcross populations of maize by symptom rating and infrared spectroscopy. <i>Plant Breeding</i> , 2015, 134, 529-534. | 1.0 | 15 |
| 122 | Genes for wheat stem rust resistance postulated in German cultivars and their efficacy in seedling and adult plant field tests. <i>Plant Breeding</i> , 2018, 137, 301-312. | 1.0 | 15 |
| 123 | Early prediction of biomass in hybrid rye based on hyperspectral data surpasses genomic predictability in less-related breeding material. <i>Theoretical and Applied Genetics</i> , 2021, 134, 1409-1422. | 1.8 | 15 |
| 124 | Multi-parent QTL mapping reveals stable QTL conferring resistance to <i>Gibberella</i> ear rot in maize. <i>Euphytica</i> , 2021, 217, 1. | 0.6 | 15 |
| 125 | Rye introgression lines as source of alleles for pollen fertility restoration in Pampa CMS. <i>Plant Breeding</i> , 2009, 128, 528-531. | 1.0 | 14 |
| 126 | Amino acid digestibility of different rye genotypes in caecectomised laying hens. <i>Archives of Animal Nutrition</i> , 2016, 70, 470-487. | 0.9 | 14 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Candidate gene based association mapping in <i>Fusarium culmorum</i> for field quantitative pathogenicity and mycotoxin production in wheat. <i>BMC Genetics</i> , 2017, 18, 49. | 2.7 | 14 |
| 128 | Candidate Genes for Aggressiveness in a Natural <i>Fusarium culmorum</i> Population Greatly Differ between Wheat and Rye Head Blight. <i>Journal of Fungi (Basel, Switzerland)</i> , 2018, 4, 14. | 1.5 | 14 |
| 129 | Ergot infection in winter rye hybrids shows differential contribution of male and female genotypes and environment. <i>Euphytica</i> , 2020, 216, 1. | 0.6 | 13 |
| 130 | Intercontinental trials reveal stable QTL for Northern corn leaf blight resistance in Europe and in Brazil. <i>Theoretical and Applied Genetics</i> , 2021, 134, 63-79. | 1.8 | 13 |
| 131 | Selection strategies for the development of rye introgression libraries. <i>Theoretical and Applied Genetics</i> , 2009, 119, 595-603. | 1.8 | 12 |
| 132 | Resistance to Ergot in Self-incompatible Germplasm Resources of Winter Rye. <i>Journal of Phytopathology</i> , 2009, 157, 350-355. | 0.5 | 12 |
| 133 | Genetic Variation in Testcrosses and Relationship between Line per se and Testcross Performance for Resistance to Gibberella Ear Rot in Maize. <i>Crop Science</i> , 2010, 50, 1691-1696. | 0.8 | 12 |
| 134 | Identification of quantitative trait loci in rye introgression lines carrying multiple donor chromosome segments. <i>Theoretical and Applied Genetics</i> , 2013, 126, 49-58. | 1.8 | 12 |
| 135 | Analysis of Covariation of Grain Yield and Dry Matter Yield for Breeding Dual Use Hybrid Rye. <i>Bioenergy Research</i> , 2014, 7, 424-429. | 2.2 | 12 |
| 136 | Stability of Adult-plant Resistance to <i>Septoria tritici</i> blotch in 24 European Winter Wheat Varieties Across Nine Field Environments. <i>Journal of Phytopathology</i> , 2011, 159, no-no. | 0.5 | 11 |
| 137 | Quantitative genetic parameters for selection of biomass yield in hybrid rye. <i>Plant Breeding</i> , 2012, 131, 100-103. | 1.0 | 11 |
| 138 | Toward a Selection of Broadly Adapted Germplasm for Yield Stability of Hybrid Rye under Normal and Managed Drought Stress Conditions. <i>Crop Science</i> , 2015, 55, 1026-1034. | 0.8 | 11 |
| 139 | Significance of host complexity and diversity for race-specific leaf-rust resistance in self-fertile synthetic rye populations. <i>Plant Breeding</i> , 2006, 125, 225-230. | 1.0 | 10 |
| 140 | Phenotypic selection for high resistance to <i>Fusarium</i> head blight after introgression of quantitative trait loci (QTL) from exotic spring wheat and verification by simple sequence repeat markers <i>a posteriori</i> . <i>Plant Breeding</i> , 2008, 127, 217-221. | 1.0 | 10 |
| 141 | Analyzing Genetic Diversity for Virulence and Resistance Phenotypes in Populations of Stem Rust (<i>Puccinia graminis</i> f. sp. <i>secalis</i>) and Winter Rye (<i>Secale cereale</i>). <i>Phytopathology</i> , 2016, 106, 1335-1343. | 1.1 | 10 |
| 142 | Hyperspectral Reflectance Data and Agronomic Traits Can Predict Biomass Yield in Winter Rye Hybrids. <i>Bioenergy Research</i> , 2020, 13, 168-182. | 2.2 | 10 |
| 143 | Ergot Alkaloid Contents in Hybrid Rye are Reduced by Breeding. <i>Agriculture (Switzerland)</i> , 2021, 11, 526. | 1.4 | 9 |
| 144 | The Development of a Host-Pathogen System for Evaluating <i>Fusarium</i> Resistance in Early Growth Stages of Wheat. <i>Journal of Phytopathology</i> , 1988, 121, 150-158. | 0.5 | 8 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Test Systems for Evaluating Quantitative Resistance Against Fusarium Foot Rot in Inbred Lines of Winter Rye. <i>Plant Breeding</i> , 1992, 108, 274-282. | 1.0 | 8 |
| 146 | Inheritance of Foot Rot Resistance in Winter Rye. <i>Crop Science</i> , 1995, 35, 388. | 0.8 | 8 |
| 147 | Lack of Association between Fusarium Foot Rot and Head Blight Resistance in Winter Rye. <i>Crop Science</i> , 1997, 37, 327-331. | 0.8 | 8 |
| 148 | Correlation between per se and Testcross Performance in Rye (<i>Secale cereale</i> L.) Introgression Lines Estimated with a Bivariate Mixed Linear Model. <i>Crop Science</i> , 2010, 50, 1863-1873. | 0.8 | 8 |
| 149 | Combined inoculation of wheat pathogens <i>Zymoseptoria tritici</i> and <i>Fusarium culmorum</i> as a tool for increasing selection intensity in resistance breeding. <i>Plant Breeding</i> , 2014, 133, 543-547. | 1.0 | 8 |
| 150 | Genomics-Based Hybrid Rye Breeding. , 2019, , 329-348. | | 8 |
| 151 | Mapping Stem Rust (<i>Puccinia graminis</i> f. sp. <i>secalis</i>) Resistance in Self-Fertile Winter Rye Populations. <i>Frontiers in Plant Science</i> , 2020, 11, 667. | 1.7 | 8 |
| 152 | Two unnecessary powdery mildew resistance genes in a synthetic rye population are neutral on fitness. <i>Euphytica</i> , 1995, 81, 163-170. | 0.6 | 7 |
| 153 | Variation and covariation for quantitative resistance to head blight (<i>Fusarium culmorum</i>) in two testcross series of S2 lines in winter rye. <i>Plant Breeding</i> , 1995, 114, 155-159. | 1.0 | 7 |
| 154 | Competitive Aggressiveness in Binary Mixtures of <i>Fusarium graminearum</i> and <i>F. culmorum</i> Isolates Inoculated on Spring Wheat with Highly Effective Resistance QTL. <i>Journal of Phytopathology</i> , 2010, 159, no-no. | 0.5 | 7 |
| 155 | Variation and covariation for <i>Gibberella</i> ear rot resistance and agronomic traits in testcrosses of doubled haploid maize lines. <i>Euphytica</i> , 2012, 185, 441-451. | 0.6 | 7 |
| 156 | Detection of donor effects in a rye introgression population with genome-wide prediction. <i>Plant Breeding</i> , 2015, 134, 406-415. | 1.0 | 7 |
| 157 | A multiple disease test for field-based phenotyping of resistances to <i>Fusarium</i> head blight, yellow rust and stem rust in wheat. <i>European Journal of Plant Pathology</i> , 2018, 151, 451-461. | 0.8 | 7 |
| 158 | Multi-parental QTL mapping of resistance to white spot of maize (<i>Zea mays</i>) in southern Brazil and relationship to QTLs of other foliar diseases. <i>Plant Breeding</i> , 2021, 140, 801. | 1.0 | 7 |
| 159 | Effective Pollen-Fertility Restoration Is the Basis of Hybrid Rye Production and Ergot Mitigation. <i>Plants</i> , 2022, 11, 1115. | 1.6 | 7 |
| 160 | Genetic variation for resistance and mycotoxin content of European maize inoculated with <i>Fusarium graminearum</i> and <i>F. verticillioides</i> . <i>Cereal Research Communications</i> , 2008, 36, 45-48. | 0.8 | 6 |
| 161 | Genetic variation for foot-rot and <i>Fusarium</i> head-blight resistances among full-sib families of a self-incompatible winter rye (<i>Secale cereale</i> L.) population. <i>Theoretical and Applied Genetics</i> , 1995, 91-91, 862-868. | 1.8 | 5 |
| 162 | Genotypic correlations and QTL correspondence between line per se and testcross performance in sugar beet (<i>Beta vulgaris</i> L.) for the three agronomic traits beet yield, potassium content, and sodium content. <i>Molecular Breeding</i> , 2014, 34, 205-215. | 1.0 | 5 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | Perennial Rye: Genetics of Perenniality and Limited Fertility. <i>Plants</i> , 2021, 10, 1210. | 1.6 | 5 |
| 164 | Hybrid Rye Breeding. <i>Compendium of Plant Genomes</i> , 2021, , 13-41. | 0.3 | 5 |
| 165 | Genotypic and environmental variation in grain protein components and their relations to beta-amylase and beta-glucanase activity in malting barley. <i>Cereal Research Communications</i> , 2008, 36, 125-134. | 0.8 | 4 |
| 166 | Genetic Architecture of Cereal Leaf Beetle Resistance in Wheat. <i>Plants</i> , 2020, 9, 1117. | 1.6 | 4 |
| 167 | Mapping and validating stem rust resistance genes directly in self-incompatible genetic resources of winter rye. <i>Theoretical and Applied Genetics</i> , 2021, 134, 1989-2003. | 1.8 | 4 |
| 168 | Genome-wide association study for deoxynivalenol production and aggressiveness in wheat and rye head blight by resequencing 92 isolates of <i>Fusarium culmorum</i> . <i>BMC Genomics</i> , 2021, 22, 630. | 1.2 | 4 |
| 169 | Selection strategies in hybrid rye with special consideration of fungal disease resistances. <i>Burleigh Dodds Series in Agricultural Science</i> , 2019, , 223-246. | 0.1 | 4 |
| 170 | Quantitative-Genetic Evaluation of Resistances to Five Fungal Diseases in A Large Triticale Diversity Panel (Å—Triticosecale). <i>Crops</i> , 2022, 2, 218-232. | 0.6 | 4 |
| 171 | Biometrical analysis of alternative plot types for selection in rye. <i>Euphytica</i> , 1991, 57, 141-150. | 0.6 | 3 |
| 172 | Genome-wide prediction methods for detecting genetic effects of donor chromosome segments in introgression populations. <i>BMC Genomics</i> , 2014, 15, 782. | 1.2 | 3 |
| 173 | Be flexible and adapt easily—The great role of plasticity relative to genetic variation for aggressiveness of <i>Fusarium culmorum</i> isolates. <i>Journal of Phytopathology</i> , 2020, 168, 162-174. | 0.5 | 3 |
| 174 | Aggressiveness of <i>Fusarium culmorum</i> isolates for head blight symptoms is highly stable across four cereal crops. <i>Journal of Phytopathology</i> , 2021, 169, 387-392. | 0.5 | 3 |
| 175 | Variance Components and Correlations between Doubled Haploid Lines from Two European Flint Landraces and Their Corresponding Testcrosses for Gibberella Ear Rot Resistance, Silking Time, and Plant Height in Maize. <i>Agronomy</i> , 2021, 11, 1039. | 1.3 | 3 |
| 176 | Appreciable genetic correlation between inbred lines and testcrosses facilitates breeding for resistance to <i>Fusarium</i> head blight in hybrid rye (<i>Secale cereale</i>). <i>Plant Breeding</i> , 0, , . | 1.0 | 1 |
| 177 | Maternal differences for the reaction to ergot in unfertilized hybrid rye (<i>Secale cereale</i>). <i>European Journal of Plant Pathology</i> , 2022, 163, 181-191. | 0.8 | 1 |
| 178 | Editorial: Genetics Became to be Genomics. , 2019, , xvii-xxii. | | 0 |
| 179 | Heiliges Feuer, Abtreibung und Psycho-Droge. , 2017, , 53-76. | | 0 |